

FOLLOW THE DIRECTIONS CAREFULLY
 This product is designed for the control of many weeds in Field Corn, Small Grains, and other crops. It is also useful for other crops. The use directions which appear on the label must accompany this container.
 NEVER MIX WITH OTHER PESTICIDES.
 NEVER MIX WITH FERTILIZERS.
 NEVER MIX WITH OTHER LIQUIDS.
 NEVER MIX WITH OTHER SOLIDS.

CONTAINER DISPOSAL

Do not reuse this container. Wash thoroughly with water and discard.

LIMIT OF WARRANTY AND LIABILITY

The purchaser, user, or buyer and the limit of the liability of this company for the use of this product shall be the purchase price paid by the user or buyer for the quantity of this product. There is no warranty and this company and any other seller disavow any liability for damages arising from combining substances or from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

RECEIVED
 JUN 1 1973

**CAUTION
 KEEP OUT OF REACH
 OF CHILDREN**

See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL
Banvel 2S
 HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, IL 60611

90

Continued from page 88/89
**FENCEROWS AND SIMILAR
 NONCROPLAND
 (NOT PASTURE AND RANGELAND)**

USE	WEED
Fencerows Non Cropland	hedstraw field bindweed blackberry bluebell bracken fern pricklypear (cactus) hop clover dewberry grape Carolina geranium wild honeysuckle horsemint horseweed hutsache Russian knapweed kudzu
	bull nettle poison oak running liver plantain (turbidella) pokeweed leafy spurge sumac Canada thistle sowthistle Dalmatian toadflax vetch white lupine wild plum water hemlock willow yucca
	bracken fern

Continued on page 92

91

**IMPORTANT
 OBSERVE ALL PRECAUTIONS ON
 PAGES 4, 5, 6 AND 113**

DOSAGE PER ACRE	APPLICATION
2 to 4 gallons (4 to 8 pounds dicamba acid equivalent) For spot treatment mix 3 to 6 tablespoons BANVEL 2S Herbicide with 1 gallon water to treat 1 square rod (272 square feet)	For control or suppression of listed weeds, apply BANVEL 2S when weeds are actively growing
2 to 4 gallons (4 to 8 pounds dicamba acid equivalent)	Apply as a preemergence application before frond emergence

Continued on page 92

FOLLOW THE DIRECTIONS CAREFULLY

BANVEL 2S herbicide is used for the control of many weeds in Field Corn, Small Grains, Sorghum, Hay, Pasture Grass and for other crops. The use directions which are on a separate book must accompany this container.

NOT AN AGRICULTURAL PESTICIDE. NOT SPECIFIED ON THE LABEL. Use directions leaf pamphlet No. 0718A 100AA attached. For additional information write the manufacturer.

CONTAINER DISPOSAL

Use empty container. Wash thoroughly with water and detergent. Discard.

LIMIT OF WARRANTY AND LIABILITY

The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all injury, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty, and this company and any other seller disavow all liability for any loss, injury or damages arising from combining substances or from any use of this product not recommended in label directions or arising from misuse, improper handling, or improper storage.

THIS IS A A

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL
Banvel 2S
HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

92

Continued from page 90/91

FENCEROWS AND SIMILAR NONCROPLAND (NOT PASTURE AND RANGELAND)

USE	WEED
Fencerows and Cropland	Eastern whiteclover
	McCartney Rose (undisturbed)
	McCartney Rose (disturbed—plants that have been mowed or bush hogged)

93

IMPORTANT OBSERVE ALL PRECAUTIONS ON PAGES 4, 5, 6 AND 83

DOSAGE PER ACRE	APPLICATION
Prepare a spray mix using 2 to 4 gallons (4 to 8 pounds dicamba acid equivalent) in 100 gals. of water	Apply to ground under tree as a basal treatment using 1/4 to 1/2 pint of spray solution per inch diameter of the plant. May also be used as a stem foliage treatment with sufficient water to give good coverage.
6 quarts (3 pounds dicamba acid equivalent) or 4 quarts (2 pounds dicamba acid equivalent) plus 1.5 to 2 pounds (acid equivalent) of 2,4-D amine	Mix required amount of BANVEL 2S herbicide alone or tank mix BANVEL 2S + 2,4-D amine with 100 gallons of water per acre. Treat all plant foliage and stems to run off and spray ground area around plants.
4 to 6 quarts (2 to 3 pounds dicamba acid equivalent) plus 1.5 to 2.0 pounds (acid equivalent) 2,4-D amine	Tank mix required amount of BANVEL 2S plus 2,4-D amine with 100 gallons of water per acre. Treat all plant foliage and stems to run-off and spray ground area around plants.

FOLLOW THE DIRECTIONS CAREFULLY
 BANVEL 2S is aicide used for the control of many weeds in Field Corn, Sorghum, Grain Sorghum, and Cleopatra. Pasture Grass and for other crops. The use directions which are on a separate booklet must accompany this container.
 DO NOT USE IN ANY MANNER NOT SPECIFIED ON THIS LABEL.
 For use directions, see pamphlet No. 0-765-100AA attached.
 For additional copies, write to the manufacturer.

CONTAINER DISPOSAL
 Do not reuse this container. Wash thoroughly with water and detergent. Discard in place.

LIMIT OF WARRANTY AND LIABILITY
 The exclusive remedy of the user or buyer and the limit of the liability of this company or other seller for any and all losses, injuries or damages resulting from the use of herbicide of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller shall have no liability for any losses, personal injuries or damages arising from combining substances or from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

0-765-100AA

APPROVED
 1976

CAUTION
KEEP OUT OF REACH
OF CHILDREN

See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, IL 60611

56

TREE KILL BY INJECTION
DIRECTIONS FOR USE

USE	WEED
Tree kill by injection	Alder (red) Ash (white) Aspen Basswood Beech Birch (yellow paper) Dogwood Hickory Huckleberry Maple (red sugar) Oak (blackjack Post) Persimmon Pine (white) Sweetgum Blackgum
	Oak (black chestnut red white) Pine (shortleaf)

57

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4 THRU 6

DOSAGE	APPLICATION DIRECTIONS
Mix 1 part BANVEL 2S to 3 part water or use BANVEL 2S undiluted. Apply 1.0 to 2.0 milliliters (ml) per injection. Overlap cuts or space cuts up to 2 inches apart from edge to edge.	May be applied anytime during the year. To obtain satisfactory kill the cut must penetrate the bark and the cambium layer (sapwood). Application may be made by special designed injector that meters out desired quantity of chemical or cuts may be made with an axe and chemical applied with an oil can or other suitable applicator. Symptoms of injury will be noted within a few weeks but kill may take several months.
Mix 2 parts BANVEL 2S to 4 parts water. Apply 0.5 to 1.0 milliliter (ml) of mix per injection. Space cuts up to 3 inches apart from edge to edge.	

FOLLOW THE DIRECTIONS CAREFULLY

BANVEL 2S herbicide is used for the control of many weeds in Field Corn, Small Grains, Sorghum, Non-Cropland, Pasture Grass and for other crops. The use instructions which appear on the label must accompany this container.

DO NOT USE IN ANY MANNER NOT SPECIFIED ON THIS LABEL.
See Back Panel for Additional Cautions.

CONTAINER DISPOSAL

Do not reuse this container. Wash thoroughly with water and detergent. Dispose of properly.

LIMIT OF WARRANTY AND LIABILITY

The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all claims, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances or from any use of this product not recommended in label directions or arising from the use of improper handling or improper storage.

© 1978 AA

RECEIVED
JUN 10 1978

**CAUTION
KEEP OUT OF REACH
OF CHILDREN**

**See Back Panel
for Additional Cautions**

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL

Banvel 2S
HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

USE DIRECTIONS

VELSICOL CHEMICAL CORPORATION

341 Ohio Street, Chicago, Illinois 60611

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

**Do not mix with pesticides by using them
carefully. Read and follow label directions.**

© 1978 AA

MAIZE: Some people grow the traditional maize weeds in their own backyards. Grains of sorghum, millet, and finger millet are also grown. The use of these grains as a separate crop must accompany this maize.

CONTAINER NO. 608

7-1-68

[illegible]

1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 26

E.P.A. REG. NO 076-255-AA
E.P.A. EST NO

Banvel® 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60601

INERT INGREDIENTS

Wash thoroughly after handling in case of contact with eyes. Flush with water for 15 minutes and seek medical attention.

Do not apply except as recommended. Do not spray on plants or parts of plants where chemicals may be washed or moved to parts with young plants.

Prevent injury to product: When applying by ground or air to desirable plants, particularly beans, melons, fruit trees, grasses, ornamentals, peas, potatoes, snapbeans, sunflowers, tobacco, tomatoes and other broadleaf crops. Do not spray on or near plants to be killed.

Do not apply by airplane when sensitive crops are nearby. Consult your State or local authorities for possible aerial application or other application restrictions. When using ground equipment use sufficient water to give good coverage.

Do not contaminate irrigation ditches or water used for domestic purposes. To avoid injury to desirable plants, equipment used for BANVEL 2S Herbicide should be thoroughly cleaned and before reuse to handle or apply any other agricultural chemicals.

All uses of BANVEL 2S are intended for a normal growing interval between plants and

[illegible]

FOLLOW THE DIRECTIONS CAREFULLY
 Banvel 2S is a selective herbicide for the control of many weeds in the corn, soybean, and cotton crops. It is a systemic herbicide and should be applied to the soil in the row between the plants.

VELSICOL
 BANVEL 2S
 HERBICIDE

CONTAINER DISPOSAL

Wash the container thoroughly with water and dispose of it as directed.

LIMIT OF WARRANTY AND LIABILITY

The user assumes the responsibility for the use of this product. The company does not warrant that the product will perform under all conditions. The company does not warrant that the product will be free from defects. The company does not warrant that the product will be free from damage. The company does not warrant that the product will be free from injury. The company does not warrant that the product will be free from loss. The company does not warrant that the product will be free from damage. The company does not warrant that the product will be free from injury. The company does not warrant that the product will be free from loss.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

USE DIRECTIONS

VELSICOL

DEC 18 1978
Banvel 2S

HERBICIDE

Contains 21.23% 3,6-dichloro-p-aminic acid (dicamba) or 2 pounds per gallon

ACTIVE INGREDIENTS:	
Sodium salt of dicamba (3,6-dichloro-p-aminic acid)	23.32%
Sodium salts of related acids	3.18%
INERT INGREDIENTS:	73.50%
TOTAL	100.00%

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN

READ CAREFULLY
DISPOSAL
NO LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

EPA REG. NO 876-255-AA
EPA EST. NO

VELSICOL
Banvel 2S
HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORP
341 E. OHIO ST., CHICAGO, ILL.

CONTENTS

General Cautions Covering All Uses of BANVEL 2S Herbicide	4-6
Field Corn	
Pre- and Early Post Emergence (BANVEL 2S + Lasso Tank-Mix)	7-13
Early Postemergence (BANVEL 2S + Atrazine Tank-Mix)	14-15
Postemergence (BANVEL 2S Alone or BANVEL 2S + 2,4-D Tank-Mix)	16-19
Small Grains (Not Underseeded to Legumes)	
Spring Seeded Wheat and Oats, Spring Seeded Barley, Fall Seeded Barley, Oats and Wheat (BANVEL 2S Alone or BANVEL 2S + 2,4-D or MCPA Tank-Mix)	20-25

CONTENTS

Spring and Fall Seeded Wheat (BANVEL 2S + Bromoxynil Tank-Mix)	26-27
Spot Application to Cropland Rotated to Wheat	28-31
Grain Sorghum	32-35
Postemergence	36-37
Harvest Aid Treatment	
Pasture and Rangeland Grasses and Non-cropland	38-81
BANVEL 2S Alone	40-55
BANVEL 2S + 2,4-D or 2,4,5-T Tank Mix	56-81
Ferns and Similar Noncropland	82-93
Grass Introduction	94-95
Tree Introduction	96-97

NOTICE: Read "LIMIT OF WARRANTY AND LIABILITY" on the container before buying or using. If terms are not acceptable, return at once unopened

FOLLOW THE DIRECTIONS CAREFULLY

Read the entire label carefully before using this product. The label contains important information which must be read and understood before using this product. The label also contains information which must be read and understood before using this product. The label also contains information which must be read and understood before using this product.

CONTAINER DISPOSAL

Do not reuse or refill this container. Dispose of this container in accordance with the instructions on the label.

LIMIT OF WARRANTY AND LIABILITY

The exclusive responsibility for the proper use of this product and the proper disposal of this product lies with the user. The user is responsible for the proper use of this product and the proper disposal of this product. The user is responsible for the proper use of this product and the proper disposal of this product.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. CHIO ST., CHICAGO, ILL. 60611

THESE PRECAUTIONS APPLY TO ALL USES OF BANVEL® 2S HERBICIDE

Additional precautions and restrictions applying to specific uses appear with the use directions under the heading **IMPORTANT**.

CAUTION

Hazard to Humans and Domestic Animals.

Caution: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

IMPORTANT

Do not apply (except as recommended) on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots.

Apply to weeds and brush in pastures, fields, and other areas where they are unwanted. Do not apply to crops or other desirable plants. Do not apply to areas where children or animals may be present. Do not apply to areas where water may be present. Do not apply to areas where fire may be a hazard. Do not apply to areas where there is a risk of contamination of water or soil. Do not apply to areas where there is a risk of contamination of air. Do not apply to areas where there is a risk of contamination of food or feed. Do not apply to areas where there is a risk of contamination of clothing or equipment. Do not apply to areas where there is a risk of contamination of other property. Do not apply to areas where there is a risk of contamination of the environment. Do not apply to areas where there is a risk of contamination of the atmosphere. Do not apply to areas where there is a risk of contamination of the hydrosphere. Do not apply to areas where there is a risk of contamination of the lithosphere. Do not apply to areas where there is a risk of contamination of the biosphere. Do not apply to areas where there is a risk of contamination of the geosphere. Do not apply to areas where there is a risk of contamination of the atmosphere. Do not apply to areas where there is a risk of contamination of the hydrosphere. Do not apply to areas where there is a risk of contamination of the lithosphere. Do not apply to areas where there is a risk of contamination of the biosphere. Do not apply to areas where there is a risk of contamination of the geosphere.

[illegible]

1. The first part of the document is a list of names and addresses, which appears to be a directory or a list of contacts. The names are written in a cursive script, and the addresses are listed below them. The list includes names such as "Mr. J. H. Smith", "Mr. W. H. Jones", and "Mr. R. L. Brown".

The exclusion of liability for the use of the product by the user or buyer, or for damages arising from the use of the product, is not a condition for the sale of the product, and the user or buyer is not obliged to accept it. The user or buyer is not obliged to accept it. The user or buyer is not obliged to accept it.

• • • • •

E.P.A. REG. NO 876-255-AA
E.P.A. EST. NO

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11, ILL.

All tank mixes including those with fluid fertilizers must be properly agitated and mixed according to recommended rates and gallonages to insure compatibility. Add BANVEL 2S Herbicide to spray tank prior to adding 2,4-D or MCPA.

DO NOT USE IN ANY MANNER NOT SPECIFIED ON THE LABEL

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

1084

ACKNOWLEDGENCE

Mixture

Mont.

OBSERVE ALL PRECAUTIONS
Observe all precautions
and instructions.

Only 1000 copies of the
first issue of the
second volume of the

15. $\frac{1}{2} \log 2$

...the ...
...the ...
...the ...

APR 1968

1. The first step is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

The express warranty is made by the seller and the manufacturer of this computer and other related products. It is a promise or damages resulting from the use or handling of this product. It is the responsibility of the buyer to pay for the quantity of the product and the quality of the workmanship and any other seller's disclaimer of liability. The seller is not responsible for damages arising from combining substances from any of the products recommended by the directions or arising from this seller's computer being used in a hazardous environment.

100

E.P.A. REG. NO 876-255-AA
E.P.A. EST NO

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST., CHICAGO II 60611

Avoid incorporation of BANVEL 2S plus LASSO into the soil to minimize potential corn injury. Soil disturbance after planting tends to pile treated soil into press wheel track. If it is necessary to use a rotary hoe, delay rotary hoeing until corn has spiked and do not disturb soil more than one half (1/2) inch deep. Apply BANVEL 2S plus LASSO far enough behind equipment to avoid incorporation by the planter wheel or other covering device.

Consult the LASSO label concerning the method of application, general information and use precautions for LASSO as a pre- and early post-emergence application to field and silage corn.

Do not graze treated areas. Do not harvest or feed immature corn forage to cattle until at least 12 weeks following treatment with BANVEL 2S plus LASSO or prior to ensilage (milk stage), whichever is later.

* ASSO is a registered trademark of Monsanto Company.

RECEIVED
FEB 21 1968

[illegible][illegible]

SP 1 M.L.D.

FOLLOW THE DIRECTIONS CAREFULLY

Read the entire label carefully before using this product. Do not use if the label is torn, missing, or faded. Do not use if the product is not in the original container. Do not use if the product is not in the original container. Do not use if the product is not in the original container.

CONTAINER DISPOSAL

Do not reuse this container. Do not burn or incinerate this container. Do not dump this container into the ground or water.

LIMIT OF WARRANTY AND LIABILITY

This product is sold as a herbicide and the user assumes the responsibility for its use. The user should read the label carefully and follow the directions. The user should not use this product if the label is torn, missing, or faded. The user should not use this product if the product is not in the original container. The user should not use this product if the product is not in the original container.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

- 10 **FIELD CORN**
PRE- AND EARLY POST-EMERGENCE
Continued from page 9
- 3 Add the proper amount of LASSO to the tank. Continue agitation.
 - 4 Fill the rest of the tank with water, agitating continuously.
 - 5 The compatibility of BANVEL 2S plus LASSO 4EC tank mixtures with sprayable fluid fertilizers must be determined by a jar test (pre-emergence use only).

JAR TEST

- a Add one pint of fertilizer solution to a quart jar.
 - b Add two teaspoons of BANVEL 2S and five teaspoons of LASSO 4EC.
 - c Shake jar vigorously and let stand for two hours.
 - d If herbicide does not ball up or form oily film on top of fertilizer solution, mixture is compatible and may be used together providing agitation is maintained in spray tank.
- If a precipitate forms or the solution flakes, the herbicide and the fluid fertilizer should not be sprayed in the same tank.

Keep spray mixture in suspension at all times, either by mechanical means or with bypass or jet agitation from the pump. If spray mixture is allowed to settle, thorough agitation will be required to suspend the mixture before resuming spraying.

Keep bypass line on or near bottom of tank to minimize foaming.

FOR THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

E.P.A. REG. NO 876-255-AA
E.P.A. EST. NO

Banvel® 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST., CHICAGO, ILL 60611

**FIELD CORN
PRE- AND EARLY POST-EMERGENCE
DIRECTIONS FOR USE**
NOTE Before applying, read IMPORTANT section
carefully

WEED	Dosage/Area Broadcast/Application
Broadleaf Weeds Barnyardgrass Crabgrass Fall Pannum Fall Paspalum Goosegrass Green foxtail Yellow foxtail Other annual grass weeds	COARSE SOIL BANVEL 2S Herbicide plus LASSO 4EC is not recommended on sandy textured soils
Broadleaf Weeds Annual Ragwort Big Bluestem Cockspurweed Climbing Mistletoe Cocklebur Common Ragweed Hairy Vetch Lamb's-quarters Lady Saurge Pigweed/Cornflower Purple Spill (Grounds) Purslane - seedling stage Smartweed Sunflower Velvetleaf Water Hyacinth Wild Blueberry Wild Cottonwood Wild Mustard Yarrow	MEDIUM SOILS Lawn and SW Lawns with 2% or less Organic Matter: 0.50 quart BANVEL 2S (4 oz dicamba acid equivalent) tank mixed with 1.5 quarts of LASSO 4EC (equivalent to 1 1/2 lbs Alachlor)
Lawn and Site Lawns with greater than 2% Organic Matter: 1.0 quart BANVEL 2S (8 oz dicamba acid equivalent) tank mixed with 2.0 quarts of LASSO 4EC (equivalent to 2.0 lbs Alachlor)	

**FIELD CORN
PRE- AND EARLY POST-EMERGENCE
DIRECTIONS FOR USE**

NOTE Before applying, read **IMPORTANT** section carefully.

Dosage/Acre Broadcast/Application	Dosage/Acre Band Application
FINES SOILS Silty Clay Loam, Clay Loam, Silty Clay, and Clay Soils 1 Quart BANVEL 2S16 or 4 amba acid equivalent tons mixed with 2.5 quarts of LASSO 4EC equivalent to 2.5 lbs Alachlor)	*To determine rates of BANVEL, 2S and LASSO for band application: Divide band width (in inches) by the number of inches between crop rows and multiply by the recommended broadcast rate in quarts for each herbicide (for the proper soil type) as shown by the formula below: $\frac{\text{Width of Band (Inches)}}{\text{Row Spacing (Inches)}} \times \text{Broadcast Rate (Qts./Acre)} = \text{Quarts Per Acre in Band}$

FD

... ..

THE USER SHALL BE RESPONSIBLE FOR ANY DAMAGE TO THE USER'S EQUIPMENT OR TO THE EQUIPMENT OF OTHERS, INCLUDING BUT NOT LIMITED TO, THE EQUIPMENT OF THE USER'S EMPLOYER, ARISING FROM THE USE OF THE SOFTWARE. THE USER SHALL BE RESPONSIBLE FOR ANY DAMAGE TO THE USER'S EQUIPMENT OR TO THE EQUIPMENT OF OTHERS, INCLUDING BUT NOT LIMITED TO, THE EQUIPMENT OF THE USER'S EMPLOYER, ARISING FROM THE USE OF THE SOFTWARE.

EPA REG NO 876-255-AA
EPA EST NO

Banvel 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

IMPORTANT

Observe all precautions on labeling of all products used in tank mixtures.

Do not graze or harvest for dairy or beef feed prior to the ensilage stage (milk stage). Consult the Atrazine 80 W label concerning instructions, method of application and use precautions for Atrazine.

DOSEAGE / ADMINISTRATION

APPLICATION DIRECTIONS

Take the **BANVEL 25** plus Aflazine and make application every 3 weeks after planting (when there is already some emergence stages) and before grass is 15 inches tall.

It may be necessary to calculate at day-by time to remove surviving weeds or to adjust the

NEW APPLICATION

To determine rates of **BANVEL 25** plus Aflazine for new application

Divide band width by the inches between crop rows and multiply the answer by the recommended band cast rate for each herbicide as shown by the formula below

width of band (inches) × Broadcast rate (in pounds/acre) for **BANVEL 25** or the
new spacing (inches) rate for Aflazine (lb/acre)

= Amount/acre in band (in pounds for **BANVEL 25** or pounds for Aflazine lb/acre)

FOLLOW THE DIRECTIONS CAREFULLY

Always use Banvel 2S in accordance with the directions on the label. Do not use Banvel 2S on crops or in areas where it is not recommended. Do not use Banvel 2S on crops or in areas where it is not recommended. Do not use Banvel 2S on crops or in areas where it is not recommended.

CONTAINER DISPOSAL

Do not reuse or refill this container. Wash thoroughly with water and detergent. Do not use for food or feed.

LIMIT OF WARRANTY AND LIABILITY

The seller warrants that the product is as labeled and that it is free from adulteration. The seller does not warrant that the product will perform under all conditions. The seller does not warrant that the product will be free from defects. The seller does not warrant that the product will be free from defects.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL
Banvel 2S
 HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

16

FIELD CORN — POST-EMERGENCE

BANVEL® 2S ALONE OR BANVEL® 2S PLUS 2,4-D
TANK MIXTURE

IMPORTANT

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6

Observe all precautions on labeling of all products used in tank mixture.

Do not graze or harvest for dairy or beef feed prior to ensilage stage (milk stage).

BANVEL 2S SPRAY DRIFT: Soybeans, tobacco and other broadleaf crops are sensitive to BANVEL 2S as it requires only a small amount of chemical per acre for symptoms to appear. Symptoms generally appear about 2 weeks after drift occurs. Soybeans generally outgrow these initial symptoms without any adverse effect on soybean yields. Affected tobacco leaves will be permanently malformed.

To minimize BANVEL 2S spray drift into soybeans and other sensitive crops, observe the following precautions:

1. Use coarse sprays. Use 20 gallons or more of water per acre. Keep the spray pressure below 20 psi. For broadcast application, use TK5 tips (or equivalent).

17

2. Spray with the nozzles as close to the weeds as is practical for good coverage. When soybeans, tobacco or other susceptible broadleaf crops are nearby, apply BANVEL 2S Herbicide alone or BANVEL 2S Herbicide plus 2,4-D with drop nozzles and direct spray onto the soil and weeds beneath the level of the lower corn leaves when the corn is 8 inches or more in height. When susceptible crops are nearby, do not apply BANVEL 2S or BANVEL 2S plus 2,4-D after corn is 24 inches tall. (When BANVEL 2S plus 2,4-D is used on corn 8 inches or more in height, drop nozzles must be used at all times.)

3. Do not apply BANVEL 2S Herbicide when the daily temperature is expected to be over 85° F.

4. Make application when there is some air movement but wind is less than 5 miles per hour. If any wind over 5 mph is blowing in the direction of sensitive crops, do not spray.

Do not apply BANVEL 2S Herbicide in the vicinity of soybeans if any one of the following conditions exists:

- 1. The corn is more than 24 inches tall.
- 2. Soybeans in the vicinity are more than 10 inches tall.
- 3. Soybeans in the vicinity have begun to bloom.

Continued on page 18

1. Follow the directions to carry out

1. The first part of the document is a list of names and titles, including "The Hon. Mr. Justice" and "The Hon. Mr. Justice".

other than the use of the product. The user or damages resulting from the use or handling of the product shall be the responsibility of the user or the person who handles the product. The company shall not be liable for damages resulting from the use or handling of the product if the user or the person who handles the product has not followed the instructions for use or handling of the product.

1999

EPA REG. NO 876-255-AA
EPA EST. NO

Banvel® 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST., CHICAGO II 60611

FIELD CORN – POSTEMERGENCE

USE	WEED	Dosage/Acre Banvel® 25 Herbicide
Field Corn not reg- istered for use on seed corn, sweet corn or popcorn)	Canada Thistle (seedling) Clover (annual) Cocklebur Morningglory (common) Lambquarters Mustard Pigweed Poa Sage (prostrate) Ragweed Smartweed Spanish nettle Sunflower Velvetleaf Waterhemp	Broadleaf 4 to 1 pint (2 to 4 oz diamba acid equivs lent) 12 gallons (real lb to 32 oz) BANVEL 25 may also be tank mixed with 4 to 8 oz (acid equivalent) of 2,4-D amine for broader spectrum weed control Rate See application direc- tions

APPLICATION DIRECTIONS

BAVULI 25 Herbicide may be applied topically to conventional maximum till or mold board corn. Corn should be 3" inches tall or until 15 days before tassels emergence whenever occurs first. Do not apply BAVULI 25 after this height or growth stage.

WATER TO USE: Waters are water which contain less than 100 ppm and it is suggested that the lowest rate of BAVULI 25 Herbicide may be used. Two pounds per acre (less than 2 inches tall) and the higher rate may be used (more than 2 inches tall) on water 100 ppm or less. Water may only be supplemented with water and/or a water reducer or drought stresser for ground application. Apply 10-60 gallons per acre.

If herbicides like ACC or other volatile crops are nearby, use drift precautions or other water.

BAVULI 25 Herbicide may be applied once or twice. This may be applied as a topical spray on corn less than 3 inches in height. The crop is in more water in height. Direct spray using crop machines once and water. Monitor the level of the water crop leaves. This and provide other weed coverage and reduce drift. After applying BAVULI 25 Herbicide, there are crop residues not used only over herbicide tobacco or other sensitive crops are nearby.

Do not exceed more than one post-emergence application of BAVULI 25 Herbicide per season.

For Bond Applications Use
To determine rates of BANVEL 25 or BANVEL 25 plus 240 for bond applications divide bond width by the inches between the clip rows and multiply the answer by the recommended broadcast rate as shown by the formula below:

width of band
rings
--- = Breadth
New spacing Rate (points) - Amount (points) for each
inch. New spacing - New-line per inch in band

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

20

21

SMALL GRAINS (Not Underseeded to Legumes)

DIRECTIONS FOR USE BANVEL 2S ALONE OR IN MIX WITH MCPA OR 2,4-D

USE	WEED
Spring seeded wheat and oats	Canada thistle (seedling or top growth) Corn chamomile Cow cockle
	Canada thistle (top growth control) Cocklebur Corn chamomile Corn cockle Cow cockle Dandelion Dragonhead American Field pennycress Field pennycress (Frenchweed)
	Smartweed Sowthistle (annual) Wild buckwheat
	Pigweed Puncturevine Ragwort Russian thistle Smartweed (annual) Sunflower Vetch Wild buckwheat Wild radish

Continued on page 22

IMPORTANT

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6

- Observe all precautions on labeling of all products used in mixtures.
- Do not use treated areas or harvest for grain until crop maturity.

Rate/Acre
 2S Herbicide
 2,4-Dicamba
 1 pint

1 pint (2 oz.
 equivalent)
 1 pint (4 to 6
 equivalent) of
 4-D

1 lb/gal for

APPLICATION DIRECTIONS

CAUTION: Do not use the tank mix of BANVEL 2S plus 2,4-D on oats.
 Make application as a foliar spray when wheat or oats are in the 2 to 5 leaf stage.
 For ground application, apply 10-40 gals. of water per acre.

Continued on page 23

FOLLOW THE DIRECTIONS CAREFULLY
 Read the entire label carefully before using this product. Do not use if the label is torn, missing, or if the directions are not clear. Do not use if the product is not in the original container. Do not use if the product is not in the original container. Do not use if the product is not in the original container.

CONTAINER DISPOSAL

Do not reuse this container. Do not burn or incinerate this container. Do not use this container for any other purpose.

LIMIT OF WARRANTY AND LIABILITY

The purchaser of this product warrants that the product is as labeled and that it is suitable for the uses stated on the label. The purchaser warrants that the product is as labeled and that it is suitable for the uses stated on the label. The purchaser warrants that the product is as labeled and that it is suitable for the uses stated on the label.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

22 Continued from Page 20-21

SMALL GRAINS

(Not Underseeded to Legumes)

DIRECTIONS FOR USE

BANVEL 2S ALONE OR TANK MIX WITH MCPA OR 2,4-D

USE	WEED
Spring seeded barley, Montana and North Dakota	Smartweed Wild buckwheat
Fall seeded barley, oats and wheat	Canada thistle (seedling or top growth) Corn chamomile Corn cockle Low Locke Anemone Moss Lambquarters Pigweed Ragweed Smartweed Sowthistle Annual Wild buckwheat

Continued on page 24

23

IMPORTANT

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6

Observe all precautions on labeling of all products used in tank mixtures.

Do not graze treated areas or harvest for dairy feed until crop maturity.

Acres	Herbicide	APPLICATION DIRECTIONS
1/4	BANVEL	Make application when barley is at the 2 to 3 leaf stage. May be tank mixed with 4 to 6 oz. (acid equivalent) of MCPA at the 2 to 3 leaf stage or with 4 oz. (acid equivalent) of MCPA at the 5 leaf stage. Make only one application per season. Do not use more than the recommended rate. For ground application apply in 10-40 gallons of water per acre.
1/4	BANVEL	DO NOT USE ON FALL SEEDED BARLEY IN CALIFORNIA OR ARIZONA. Make application immediately after winter dormancy and before grain begins to joint. For ground application apply in 10-40 gallons of water per acre.

Continued on page 25

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

24 Continued from page 22, 23

25

SMALL GRAINS
 (Not Underseeded to Legumes)

DIRECTIONS FOR USE
 BANVEL 2S ALONE OR TANK MIX WITH MCPA OR 2,4-D

USE	WEED		
Fall seeded barley, oats and wheat	Canada thistle (top growth control) Cocklebur Corn chamomile Cow cockle Dandelion Dragonhead American Field penny- cress (Frenchweed)	Field peppergrass Goatsbeard Hempnettle Knapweed (German mess) Kochia Lodyslumb Lambsquarters Mayweed Mustards (annual) Perennial southwistle (top growth control)	Pigweed Puncture- Ragweed Russian Smartweed Stinkweed Sunflower Verch Wild buck Wild radish
Fall seeded wheat (Pacific Northwest only)	Canada thistle (seedling or top growth) Corn chamomile Cow cockle	Fiddleneck Groundwell Knapweed (German mess) Lambsquarters	Pigweed Ragweed Smartweed Southwistle (annual) Wild Buckwheat

IMPORTANT

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6

Observe all precautions on labeling of all products
 used in tank mixtures.
 Do not graze treated areas or harvest for dairy
 feed prior to crop maturity.

Rate/Acre BANVEL 2S Herbicide	APPLICATION DIRECTIONS
Tank mix 1/2 pint (2 oz. dicamba acid equivalent) with 1/2 to 1/4 pint (4 to 6 oz. acid equivalent) of MCPA or 2,4-D. *Based on 4 lb/gal for mixtures.	CAUTION: Do not use the tank mix of BANVEL 2S plus 2,4-D on oats. DO NOT USE ON FALL SEEDING BARLEY IN CALIFORNIA OR ARIZONA. Make application immediately after winter dormancy and before grain be- gins to joint. For ground application, apply in 10-40 gallons of water per acre.
Tank mix 1/2 pint (2 oz. dicamba acid equivalent) with 1/2 to 1/4 lbs (acid equivalent) of 2,4-D (LV ester).	Make application immediately after winter dormancy and before wheat begins to joint. For ground application, apply in 10-40 gallons of water per acre.

**See Back Panel
for Additional Cautions**

EPA REG NO 076-255-AA
EPA EST NO

Banvel 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11, ILL.

YANKEE. S
- SE AIR -
WATER. WAT N. ARA H.

SUTHERLAND
- P. M.
JANUARY
WASHINGTON
AARMS
- P. M.
JANUARY
SUTHERLAND

NEED

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6
 Observe all precautions on labeling of all products.
 Do not tank mixtures.
 Do not graze treated areas or harvest for 30 days
 prior to crop maturity.

RATE OF APPLICATION/ACRE	APPLICATION DIRECTIONS
1/2 pint BANVEL, 25 (1/2 or dicamba acid equivalent) plus 1 to 1 1/2 units Bromine or Butrictin (14 to 1.0% acid equivalent of bromoxynil)	Apply BANVEL 2S plus bromoxynil as a tank mix. Make application when weeds are small (early seedling) and when wheat is from 3 leaf up to joint stage
1/2 pint BANVEL, 2S (1/2 dicamba acid equivalent) plus 1 to 1 1/2 pints Bromine or Butrictin (14 to 1.0% acid equivalent of bromoxynil)	Apply BANVEL 2S plus bromoxynil as a tank mix. Make application when weeds are small (early seedling) and when wheat is from 3-leaf up to joint stage May be applied in fall, winter, or spring.

MIXING DIRECTIONS: Spray volume of BANVEL 25 plus bromoxynil will vary in certain areas. Unless recommendations specify otherwise, use at least 10 gallons of water per acre for ground applications. Water should be added to the spray tank first, followed by BANVEL 25 plus bromoxynil.

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

EPA REG. NO. 876-255-AA
EPA EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, IL 60611

28

29

**DIRECTION FOR USE —
FOR SPOT APPLICATION ONLY**

TO PERENNIAL WEEDS IN CROPLAND
ROTATED TO WHEAT

IMPORTANT

OBSERVE ALL PRECAUTIONS ON
PAGE 4 THRU 6

For intervals between BANVEL 2S treatment
wheat planting, see application directions for
specific states

Barley, oats, corn, sorghum, annual and perennial
grass crops may be planted in treated areas one year
after applications

Do not plant any crop grown for seed (other than
perennial grass seed) in treated areas until 3 years
after application

Do not plant broadleaf crops (alfalfa, beans, peas,
potatoes and sugarbeets) on treated areas until 2
years after application

Do not treat submerged cropland or where soil
remains saturated with water throughout the year

MAKE ONLY ONE APPLICATION OF DICAMBA PER
YEAR

Some injury to wheat and rotated barley may occur
when BANVEL 2S is applied to soils containing less
than 1% organic matter or sandy soils when planting
takes place after the shortest interval

Continued on page 30

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

30 Continued from page 29

**DIRECTIONS FOR USE —
FOR SPOT APPLICATION ONLY
TO PERENNIAL WEEDS IN CROPLAND
ROTATED TO WHEAT**

STATE	WEED
Idaho Montana Nebraska Oregon Utah Washington	Butter dock Black knapweed Canada thistle Curly dock Field bindweed (morningglory) Leafy spurge Russian knapweed Tansy ragwort See Note below
Colorado Iowa Kansas Minnesota Missouri Nebraska North Dakota South Dakota Wyoming	Canada thistle Field bindweed Leafy spurge Russian knapweed See Note below

*In most cases, these treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

**OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 28 AND 29**

Dosage/Acre BANVEL 2S Herbicide	APPLICATION DIRECTIONS
6 to 12 quarts (4 to 6 lbs. dicamba acid equivalent)	Spot application may be made to fallow land, wheat stubble or land to be rotated to wheat. Application can be made in mid summer to the fall of the year when weeds are actively growing. Wheat may be planted one month after application. BANVEL 2S applied at rates of 6 lbs. acre (dicamba acid equivalent) may cause some wheat injury.
1 to 4 quarts (1 to 1 lb. dicamba acid equivalent)	Spot application may be made to fallow land, wheat stubble or land to be rotated to wheat. Application should be made when weeds are actively growing. Application must be made prior to planting.

...the

...and the fact that the system is not yet fully operational, the Commission has decided to postpone the final decision on the matter until the end of 1990.

... ..

[illegible]

E.P.A. REG. NO 076-255-AA
E.P.A. EST NO

Banvel® 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11 60611

GRAIN SORGHUM - POSTEMERGENCE

Do not graze or feed treated sorghum, hay or silage prior to mature grain stage

BANVEL 2S SPRAY DRIFT—Soybeans, tobacco and other broadleaf crops are sensitive to BANVEL 2S as it requires only a small amount of chemical per acre for symptoms to appear. Soybeans generally outgrow these initial symptoms without any adverse effect on soybean yields. Affected tobacco leaves will be permanently malformed. While cotton and other sensitive broadleaf crops are more tolerant than soybeans and tobacco, similar precautions against drift should be exercised.

Do not apply BANVEL 2S to sorghum grown for seed production

Under certain conditions sorghum may show temporary effects from treatment such as onion leafing or flattening of the plants. Affected plants will normally recover within 10 to 14 days without ill effects.

Continued on page 34

1. The first step in the process of the investigation is the identification of the problem. This is done by the investigator who is assigned to the case. The investigator will then gather information about the problem and the people involved. This information will be used to determine the cause of the problem and to develop a plan to solve it.

1. Washington, D.C. 20540

the seller is not liable for the loss of the goods if the loss is caused by the fire or other cause which is not the fault of the seller. The seller is liable for the loss of the goods if the loss is caused by the fault of the seller. The seller is liable for the loss of the goods if the loss is caused by the fault of the seller.

EPA REG NO 876-255-AA
EPA EST NO

Banvel 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST CHICAGO IL 60611

USE DIRECTIONS

GRAIN SORGHUM — POSTEMERGENCE

**OBSERVE ALL PRECAUTIONS ON
PAGES 4 5, 6 32 AND 33**

APPLICATION DIRECTIONS

HANVY[®] herbicide may be used on both irrigated and non-irrigated grain sorghum and may be applied over the top of the sorghum as a directed application for most effective weed control when weeds are small. As weeds become larger they are harder to kill but will be suppressed when treated with HANVY.

Make postemergence application only using 16.25 gallons of water per acre for ground application.

Make application from 10-25 days after emergence of grain sorghum from the ground. Applications made later than 25 days after emergence of the sorghum from the ground and up through the flowering stage may reduce sorghum grain yield.

Make no more than one application per season.

For Band Application Use:
To determine rates of BANVEL 2S for band application. Divide the band width by the inches between the crop rows and multiply the answer by the recommended broadcast rate as shown by the formula below.

$$\frac{\text{width of band (inches)} \times \text{broadcast rate (pints)} \times \text{row spacing (inches)}}{43,560} = \text{Amount (pint) per acre in band}$$

separately as a "discovery" in the field of

...the ...

1. The National Security Council, in its report to the President, dated 14 October 1950, stated that the United States had no intention of using atomic weapons against the Soviet Union.

Buyer shall not be liable for damages arising from the use of the product in any manner not intended by the Seller or from the use of the product in any manner not intended by the Seller or from the use of the product in any manner not intended by the Seller.

E.P.A. REG. NO 076-255-AA
E.P.A. EST. NO

Banvel 2S

2

FOR USE ONLY IN THE STATES
OF TEXAS AND OKLAHOMA

Weed	Dosage/Acre BANVEL 2S Herbicide
Carthagenweed (Bigweed)	1 pint (4 oz dicamba acid equivalent)
Cocklebur	
Kochia	2 gallons BANVEL 2S "real"
Ambriguars	16 acres
Morning glory (annual)	
Redroot pigweed	
Sunflower	
and	
Other annual broadleaf weeds	

DO NOT GRAZE OR FEED TREATED SORGHUM: FORAGE OR Silage PRIOR TO MATURITY
GRAIN SIlAGE

HARVEST AND TREATMENT IS LIMITED TO ONE APPLICATION PER CROP SEASON
DO NOT USE BANVEL 25 FOR HARVEST-AID TREATMENT*
DICAMBA HAS BEEN APPLIED EARLIER IN THE SEASON

Application may be made over the top of the sorghum or as a direct application.

FOLLOW THE DIRECTIONS CAREFULLY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

The user of this product assumes all liability for any damage to property or persons resulting from the use of this product. The user of this product assumes all liability for any damage to property or persons resulting from the use of this product. The user of this product assumes all liability for any damage to property or persons resulting from the use of this product.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11, ILL. 60611

8

PASTURE AND RANGELAND GRASSES AND NONCROPLAND

IMPORTANT

OBSERVE ALL PRECAUTIONS ON PAGES 4 THRU 6
OBSERVE ALL PRECAUTIONS ON LABELING OF ALL
PRODUCTS USED IN TANK MIXTURES

For aerial application use 3 or more gallons of water per
acre. For ground application use sufficient water to give
good coverage.

Pasture and rangeland grasses and grasses grown for
seed may be grazed for dairy feed.

Do not graze meat animals in treated fields within 30
days of slaughter. Do not use seed from treated grasses
for feed or food purposes.

FOR DAIRY ANIMALS OBSERVE DOSAGE RATES AND
DAYS OF DELAY BETWEEN TREATMENT AND GRAZING.
THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND
GRAZING FOR ANIMALS OTHER THAN DAIRY ANIMALS WHEN
USING BANVEL 2S ALONE. WHEN USING BANVEL 2S PLUS
2,4-D OR BANVEL 2S PLUS 2,4,5-T SEE RESTRICTIONS BELOW.

FEEDING RESTRICTIONS When using BANVEL 2S alone
Interval Between BANVEL 2S Treatment and Grazing of Dairy
Animals

Dosage/Acre BANVEL 2S Herbicide	Days Delay*
Up to 2 pts (8 oz a.i.)**	graze after 7 days
Up to 2 qts (16 oz a.i.)	graze after 21 days
Up to 1 gal (32 oz a.i.)	graze after 40 days
Up to 4 qts (128 oz a.i.)	graze after 90 days

* There is no waiting period between treatment and grazing for
animals other than dairy animals when applying BANVEL 2S
alone.
**Picamba acid equivalent.

39

Treated grasses may be harvested for hay.
OBSERVE DOSAGE RATES AND DAYS OF DELAY BETWEEN TREATMENT
AND HARVESTING FOR HAY.

BANVEL 2S Herbicide Dosage/Acre	Days Delay
Up to 2 pts (8 oz a.i.)**	Harvest after 37 days
Up to 2 qts (16 oz a.i.)	Harvest after 51 days
Up to 1 gal (32 oz a.i.)	Harvest after 71 days
Up to 4 qts (128 oz a.i.)	Harvest after 90 days

**Picamba acid equivalent.

When using BANVEL 2S plus 2,4-D or 2,4,5-T

Do not graze meat animals or treated areas within 30 days
of slaughter.

BANVEL 2S Herbicide Dosage/Acre	2,4-D or 2,4,5-T Dosage/Acre	Days Delay
Up to 1 gallon (32 oz a.i.)	2 lb a.i.	40 days
Up to 4 gallons (128 oz a.i.)	4 lbs a.i.	60 days

**Picamba acid equivalent.

OBSERVE DOSAGE RATES AND DAYS OF DELAY BETWEEN TREATMENT
AND HARVESTING FOR HAY.

BANVEL 2S Herbicide Dosage/Acre	Days Delay
Up to 2 pts (8 oz a.i.)**	Harvest after 3 days
Up to 2 qts (16 oz a.i.)	Harvest after 5 days
Up to 1 gal (32 oz a.i.)	Harvest after 7 days
Up to 4 qts (128 oz a.i.)	Harvest after 9 days

**Picamba acid equivalent.

Continued on page 40

FOLLOW THE DIRECTIONS CAREFULLY

Use only as directed. Do not use if the container is empty or if the label is missing. Do not use if the product is not in the original container. Do not use if the product is not in the original container. Do not use if the product is not in the original container.

CONTAINER DISPOSAL

Do not reuse. Do not burn. Do not dump. Do not pour. Do not throw away. Do not use as a receptacle for other materials. Do not use as a receptacle for other materials. Do not use as a receptacle for other materials.

LIMIT OF WARRANTY AND LIABILITY

The seller warrants that the product is as labeled and that the label directions are true and correct. The seller does not warrant that the product will perform as advertised. The seller does not warrant that the product will be free from defects. The seller does not warrant that the product will be free from defects. The seller does not warrant that the product will be free from defects.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

EPA REG. NO. 876-255-AA
 EPA EST. NO.

VELSICOL

Banvel 2S

HEPBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

40 Continued from page 39

**PASTURE AND RANGELAND
 GRASSES AND NONCROPLAND**

USE	WEED
Pasture and Rangeland Grasses and Noncropland areas such as fence rows, roadways, woodland and similar areas	bloodroot wild buckhorn sweet clover hobbit clover cow tickle corn cockle cocklebur dog fennel (mayweed) corn chamomile barnyard (German knotgrass) barnyard lambquarters mustard field pennycress redroot pigweed tumble pigweed pearson common ragweed rabbitbrush red sorrel (sheep sorrel) smartweed, annual Spanish nettle sowthistle prostrate spurge stinkweed waterhemp

Continued on page

41

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Dosage/Acre	Application Directions
1 pint BANVEL 2S Herbicide (4 oz. di- camba acid equivalent) For spot treatment mix 1/6 teaspoon BANVEL 2S with 1 gallon water to treat 1 square rod (272 square feet)	For control or suppression of listed weeds, apply BANVEL 2S when weeds are actively growing. For ground equipment use 10 to 20 gal- lons of water per acre when treating an- nual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. Rates of BANVEL 2S Herbicide in ex- cess of 4 pounds per acre (dicamba acid equivalent) may cause temporary injury to sensitive grass species. For waiting period between treatment and grazing or harvest of treated grass see IMPORTANT section.

Continued on page 43

...the ... of ...

1. The first of these is the fact that the majority of the population of the United States is now living in urban areas. This is a result of the process of urbanization, which has been going on since the beginning of the 20th century. The population of the United States has increased from about 100 million in 1900 to over 200 million in 1950, and the majority of this increase has been in urban areas. This has led to a concentration of population in a few large cities, which has in turn led to a number of problems, such as overcrowding, pollution, and traffic congestion.

There is no doubt that the buyer's failure to inspect the car prior to purchase was a contributing factor in the damages resulting from the fire or theft of the car. The court found that the buyer's failure to inspect the car prior to purchase was a contributing factor in the damages resulting from the fire or theft of the car. The court found that the buyer's failure to inspect the car prior to purchase was a contributing factor in the damages resulting from the fire or theft of the car.

1. *Chlorophyll a* (Chl *a*)

EPA REG NO 876-255-AA
EPA EST NO

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

**PASTURE AND RANGELAND
GRASSES AND NONCROPLAND**

USE	WEED
Pasture and Rangeland Grasses and Noncropland areas such as inclosures, roadways, wasteland and similar areas	bitter melon buffaloe burdock chickweed chicory creole cudweed early dock hackberry mossy-stemmed annual puncturewort loose ragwort ragwort giant ragweed galium velvetleaf shepherdspurse leaves vertical wormwood
	Tan Grass Control Canada thistle Russian thistle field bindweed black nightshade P. v. blackhead "hot" burr perennial sorrel and other perennial broadleaf weeds

Continued on page

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Doseage/Acre	Application Directions
4 pints BANVEL 25 Herbicide 18 oz. dicamba acid equivalent For 100' treatment mix : 2 teaspoons BANVEL 25 Herbicide with 1 gal. for water to treat 1 square mile (2 1/2 square mile)	For control or suppression of listed weeds apply BANVEL 25 when weeds are actively growing. For ground equipment use 10 to 20 gallons of water per acre when treating annual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. Rate of BANVEL 25 Herbicide in excess of 4 pounds per acre (dicamba acid equivalent) may cause temporary injury to sensitive grass species. For waiting period between treatment and grazing or harvest of treated grass see IMPORTANT section

Continued on page 45

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

EPA REG. NO. 876-255-AA
 EPA EST. NO.

VELSICOL
Banvel 2S
 HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

Continued from page 44, 45
PASTURE AND RANGELAND
GRASSES AND NONCROPLAND

USE	WEED
Pasture and Rangeland Grasses and Noncropland areas such as fence rows roadways wasteland and similar areas	blueweed
	buckbrush
	wild carrot
	cottonwood seedlings
	creosotebush
	evening primrose
	franseria
	legged knawweed
	ole
	western whorled milkweed
	climbing milkweed
	stinging nettle
	silverleaf nightshade
	pepperweed (tall) whitetop
	pinque
	poison ivy
	hairy ragweed
	lance ragwort (mature stage)
	redvine
	sagebrush
	smartweed perennial
	shadows
	woodsorrel
	musk thistle
	trumpet creeper
	yellow
	vaupen

Continued on page 46

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Dosage/Acre	Application Directions
4 quarts BANVEL 2S Herbicide (2 lbs. d- camba acid equivalent) For spot treatment mix 1 1/2 tablespoons BAN- VEL 2S Herbicide with 1 gallon water to treat square foot (7.75 square feet)	For control or suppression of listed weeds apply BANVEL 2S when weeds are actively growing. For ground equipment use 10 to 20 gal- lons of water per acre when treating an- nual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. Rates of BANVEL 2S Herbicide in ex- cess of 4 pounds per acre (d-camba acid equivalent) may cause temporary injury to sensitive grass species. For waiting period between treatment and grazing or harvest of treated grass see IMPORTANT section.

Continued on page 46

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG NO 876-255-AA
 E.P.A. EST NO

VELSICOL
Banvel 2S
 HERBICIDE

Net Contents: 1 Gallon
 VELSCOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO 11, ILL. 60611

Continued from page 46/47

**PASTURE AND RANGELAND
 GRASSES AND NONCROPLAND**

IMPORTANT

OBSERVE ALL PRECAUTIONS (D-1)
 PAGES 4, 5, 6, 38 AND 39

Use	Weed	Dosage/Acre	Application Directions
Pasture and Rangeland Grasses and Noncropland areas such as fencerows, roadways, wasteland and similar areas	budweed field bindweed blackberry bluebell buckeye fern hog clover dewberry grass Carolina geranium wild honeysuckle horseweed horseweed Russian knapweed budzu buckwheat running vine plantain turbinella broad plantain silver oak blackweed leafy spurge velvet Canada thistle smoothstole Delonixia seedling velvet white lupine wild plum waterhemp willow yucca	1 to 4 gallons BANVEL 2S Herbicide (4 in P 10% dicamba acid equivalent) For spot treatment mix 3 to 6 tablespoons BANVEL 2S Herbicide with 1 gallon water to treat 1 square rod (27.5 square feet)	For control or suppression of listed weeds, apply BANVEL 2S when weeds are actively growing. For ground equipment use 10 to 20 gal- lons of water per acre when treating ar- nual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. Rates of BANVEL 2S Herbicide in ex- cess of 4 pounds per acre (dicamba acid equivalent) may cause temporary injury to sensitive grass species. For waiting period between treatment and grazing or harvest of treated grass see IMPORTANT section.

Continued on page 50

Continued on page 51

LIMIT OF WARRANTY AND LIABILITY

the product is not intended for use in the quantities in which it is being used. The product is not intended for use in the quantities in which it is being used. The product is not intended for use in the quantities in which it is being used.

1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 26

EPA REG. NO 876-255-AA
EPA EST NO

Banvel 2S

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

50 Continued from page 48/49

**PASTURE AND RANGELAND
GRASSES AND NONCROPLAND**

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Use	Wood	Dosage/Acre	Application Directions
Pasture and Rangeland Grasses and Noncropland areas such as fencerows, roadways, wasteland and similar areas	bracken fern	2 to 4 gallons BANVEL 25 Herbicide (4 to 8 lbs. dicamba acid equivalent)	Apply as a preemergence application before emergence of the fronds.
	Eastern persimmon	Prepare a spray mix using 2 to 4 gallons BANVEL 25 Herbicide (4 to 8 lbs. dicamba acid equivalent) in 100 gallons of water.	Apply to ground under tree as basal treatment using 1/4 to 1/2 pint of spray solution per inch diameter of the plant. May also be used as a stem foliage treatment with sufficient water to give good coverage.

Continued on page 52

Continued on Page 53

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

E.P.A. REG. NO. 876-255-AA
E.P.A. EST. NO.

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611

58

Continued from page 56/57

**PASTURE AND RANGELAND
GRASSES AND NONCROPLAND
FOR USE ONLY IN THE STATES OF
TEXAS, OKLAHOMA AND NEW MEXICO**

Use	Weed
Pasture and Rangeland Grasses and Noncropland areas such as fencerows, roadways, wasteland and similar areas.	brunner's campion buffalo herb burdock chickweed chicory clover curly dock lactuca morning glory annual sunflower lance ragwort (rattle stage) giant ragwort rattlebush sesbania shepherd's purse sorghum velvetleaf wormwood
	For Growth Control: Canada thistle Russian thistle field bindweed black bindweed Russian bindweed leafy spurge perennial spurge and other perennial broadleaf weeds.

Continued on page 60

59

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Dosage/Acre	Application Directions
Apply mix. 2 Pints BAN- VEL 2S Herbicide (8 oz. kamba acid equivalent) with up to a maximum of 2 lbs. acre acid equivalent 2,4-D or 4.5 T amine or LV ester.	Do not use this mixture on bent, susceptible grass pastures and clover and alfalfa pastures. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas after heading begins. For control or suppression of winter weeds, apply BANVEL 2S when weeds are actively growing. For ground equipment use 10 to 100 gallons of water per acre when treating annual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. Rates of BANVEL 2S Herbicide in excess of 4 pounds per acre (dicamba acid equivalent) may cause temporary injury to sensitive grass species. For waiting period between treatment and grazing or harvest of treated grass, see IMPORTANT section.

Continued on page 61

“”

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11 60611

51

Continued on page 43

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN

See Back Panel
for Additional Cautions

EPA REG NO 876-255-AA
EPA EST NO

VELSICOL

Banvel 2S

HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO 11, ILL. 60611

Continued from page 60/61

**PASTURE AND RANGELAND
GRASSES AND NONCROPLAND
FOR USE ONLY IN THE STATES OF
TEXAS, OKLAHOMA AND NEW MEXICO**

Use	Weeds
Pasture and Rangeland Grasses and Noncropland areas such as: fence row, roadways, wasteland and similar area.	blueweed buckbrush wild carrot cottonwood (seedlings) creosotebush evergreen primrose grasses spotted knapweed rose western whorled milkweed climbing milkweed stringing nettle silverleaf nightshade pepperweed (leaf whitening) dingus dandelion bur ragwort lance ragwort (mature stages) redvine legume smartweed, perennia shepherd woodsorrel musk thistle trumpet creeper yellow yellow

Continued on page 64

**IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39**

Dosage/Acre	Application Directions
1 tank mix 4 Quarts BANVEL 2S Herbicide (4 lbs Dicamba acid equivalent) with up to 4 maximum of 2 lbs/acre acid equivalent 2-4-0 or 2-4-5-7 amine or V ester	Do not use this mixture in dense, un- derbrush grass pastures and cover and leafy pastures Do not apply when grass is in fruit stage Do not apply to newly seeded areas until seedling begins For control of suppression of weeds apply BANVEL 2S when weeds are actively growing For ground equipment use 10 to 15 gallons of water per acre when treating annual broadleaf weeds and for top growth control of perennial broadleaf weeds for maximum control of perennial broadleaf weeds use 100 gallons of water per acre Rates of BANVEL 2S Herbicide less of 4 pounds per acre (dicamba acid equivalent) may cause temporary injury to sensitive grass species For waiting period between treatment and grazing or harvest of treated area see IMPORTANT section

Continued on page 65

FOLLOW THE DIRECTIONS CAREFULLY

Read the entire label carefully before using this product. Do not use if the label is torn, faded, or missing. Do not use if the product is not in its original container. Do not use if the product is not in its original container. Do not use if the product is not in its original container.

CONTAINER DISPOSAL

Wash the container with water. Do not reuse.

LIMIT OF WARRANTY AND LIABILITY

The product is warranted to be free from defects in material and workmanship under normal use and conditions. The manufacturer disclaims any liability for damages, including consequential damages, arising from the use of this product. The user assumes all responsibility for the proper use of this product.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
 See Back Panel
 for Additional Cautions

E.P.A. REG. NO. 876-255-AA
 E.P.A. EST. NO.

VELSICOL
Banvel 2S
 HERBICIDE

Net Contents: 1 Gallon
 VELSICOL CHEMICAL CORPORATION
 341 E. OHIO ST., CHICAGO, ILL. 60611

66
 Continuing
 from page
 65

**PASTURE AND RANGELAND
 GRASSES AND NONCROPLAND
 FOR USE ONLY IN THE STATES OF
 TEXAS, OKLAHOMA AND NEW MEXICO**

Use	Weed
For use on hedges and roadways and similar areas	mesquite (tree) hustache

Continued on page 68

67
**IMPORTANT
 OBSERVE ALL PRECAUTIONS ON
 PAGES 4, 5, 6, 38 AND 39**

Dosage/Acre	Application Directions
1 to 2 pints BANVEL 2S herbicide (4.30 to 8.60 pounds acid equivalent plus 4 to 8 oz. acid equiv herbicide 2,4-D ester)	Do not use this mixture on bent, sorgho hedge grass pastures and clover and alfalfa pastures. Do not apply when grass is in bud or flower. Do not apply to newly seeded areas until seedling begins. For control of mesquite, apply this mixture BANVEL 2S herbicide (4.30 to 8.60 pounds acid equivalent) and apply 40 to 60 days after the first green growth appears. The best time to apply is when the foliage is green. Treatment may be made using aerial or ground equipment. Apply as a spray-water emulsion in 3 gallons of finished spray per acre. The finished spray may be prepared as follows:

Continued on page 69

WITH THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

WARRANTY AND LIABILITY

Read the entire label carefully before using this product. It contains important information about safe use, including first aid, fire, and disposal. It also contains information about the hazards of the product and the precautions to be taken. It is the user's responsibility to read and understand the label and to use the product in accordance with the directions.

CAUTION
KEEP OUT OF REACH
OF CHILDREN

See Back Panel
for Additional Cautions

EPA REG NO. 875-255-AA
EPA EST. NO.

VELSICOL

Banvel 2S
HERBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORP.
341 E. OHIO ST. CHICAGO, ILL. 60601

FEATURE AND RANGE, IN
HARVEST AND NONCROPPED
LANDS, IN THE
STATE OF ILLINOIS

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6, 38 AND 39

Dosage/Acre	Application Directions
Take one 1 to 1 pint BANVEL 2S Herbicide 4 to 8 ounces dicamba acid equivalent with 4 pints of 4% amine or water.	Do not use this mixture on Bent grass, etc. turf grass, Pasture and clover, etc. turf pastures. Do not apply when grass is in seed stage. Do not apply to newly seeded areas until heading begins. For established perennial grasses, make application between November 1st and April 1st, prior to seed stage. For new seedling, make application to forage spring after the seed crop has matured. Use sufficient water to give complete coverage. 1 to 4 gal. water per acre.

Continued on Page 2

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

Use of this product is restricted to the control of brush and trees on non-cropland. The user assumes all liability for any damage to crops, livestock, or other property resulting from the use of this product. The manufacturer's liability is limited to the replacement of the product if it is found to be defective. The user is responsible for the proper application of the product and for the safety of the application.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

EPA REG NO 876 255-AA
EPA EST NO

VELSICOL

Banvel 2S

HEPBICIDE

Net Contents 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST CHICAGO, IL 60601

8

FENCEROWS AND SIMILAR NONCROPLAND (NOT PASTURE AND RANGELAND)

USE	WEED
Fencerow Non-Cropland	To control mixed brush including both deciduous (hardwood) and evergreen species. Also the list of trees controlled by BANVEL is as follows: asp asp basswood cedar cherry chinquapin cucumber tree gum dogwood elm hackberry hornbeam box elder maples oaks persimmon pine poplar sassafras serviceberry sorbush sourwood sumac sycamore thornapple thornberry willow witchhazel yucca and others

Continued on page 84

5

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4 THRU 6
OBSERVE ALL ENVIRONMENTAL CAUTIONS
LABELING OF ALL PRODUCTS USED IN MIXTURE

DOSAGE PER ACRE	APPLICATION
1.0 quart of Banvel 2S per acre in 20 to 40 gallons of water.	Apply Banvel 2S to brush and trees. Apply the mixture to the stems and leaves. Use enough water to get the mixture on the leaves.
2.0 lbs. of Banvel 2S per acre in 20 to 40 gallons of water.	Apply Banvel 2S to brush and trees. Apply the mixture to the stems and leaves. Use enough water to get the mixture on the leaves.

Continued on page 84

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

LIMIT OF WARRANTY AND LIABILITY

CAUTION
KEEP OUT OF REACH
OF CHILDREN

See Back Panel
for Additional Cautions

EPA REG NO 876-255-AA
EPA EST NO

VELSICOL

Banvel 2S

HERBICIDE

Net Contents 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST. CHICAGO 1, ILL.

Continued from page 80 of

**FENCEROWS AND SIMILAR
NONCROPLAND
(NOT PASTURE AND RANGELAND)**

USE	WEED
Preventive	spiny aster slender aster balloonvine cover dwarf malow wild garlic goldenrod diffuse knapweed spotted knapweed wild onion
Control	povertyweed erect spinyweed wild tree rough spiny weed talbott sowthistle yellow toadflax star-thistle leaved wale
Spot Treatment	drumstick dark knot wild carrot wild radish seedling creosotebush evening primrose groundsel spotted knapweed wild mesquite western whorled wildflower limbing milkweed stinging nettle silverleaf nightshade
	drumstick dark knot wild carrot wild radish seedling creosotebush evening primrose groundsel spotted knapweed wild mesquite western whorled wildflower limbing milkweed stinging nettle silverleaf nightshade

Continued on page 90

IMPORTANT
OBSERVE ALL PRECAUTIONS ON
PAGES 4, 5, 6 AND 83

DOSAGE PER ACRE	APPLICATION
2 quarts (1 pound of active acid equivalent) for spot treatment mix 2.4 lbs. Banvel 2S herbicide with 1 gallon water to treat 1 square foot (277 square feet)	For control or suppression weeds apply BANVEL 2S when weeds are actively growing
4 quarts (2 pounds of active acid equivalent) for spot treatment mix 4.8 lbs. Banvel 2S herbicide with 1 gallon water to treat 1 square foot (277 square feet)	

Continued on page 91

FOLLOW THE DIRECTIONS CAREFULLY

CONTAINER DISPOSAL

Do not burn or bury. Wash thoroughly with water.

LIMIT OF WARRANTY AND LIABILITY

This product is warranted to be free from defects in material and workmanship under normal use and handling. The warranty is limited to the replacement of the product or a refund of the purchase price paid by the user. The warranty is void if the product is used in a manner not recommended by the manufacturer. The manufacturer is not liable for any damages, including consequential damages, arising from the use of this product. The user assumes all liability for the use of this product.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Back Panel
for Additional Cautions

EPA REG NO 876-255-AA
EPA EST NO

VELSICOL

Banvel 2S

HEPBICIDE

Net Contents: 1 Gallon
VELSICOL CHEMICAL CORPORATION
341 E OHIO ST., CHICAGO, IL

90

Continued from page 88 89

FENCEROWS AND SIMILAR NONCROPLAND (NOT PASTURE AND RANGELAND)

USE	WEED
Fencerows Non-cropland	redstem field bindweed blackberry bluebell bracken fern pricklypear locust hog clover dewberry crab Carolina petalium wild honeysuckle horsemint horsemint hairsch Russian knapweed kudzu
	bull nettle pawpaw running vine plantain stachys pokeweed teasty root sage Canada thistle sp. knot D. cottonwood velvet white lupine wild plum water hemlock willow yucca
	bracken fern

Continued on page 92

IMPORTANT OBSERVE ALL PRECAUTIONS ON PAGES 4, 5, 6 AND 83

DOSAGE PER ACRE	APPLICATION
1 to 4 gallons (4 to 8 pounds dicamba acid equivalent) For spot treatment mix 1 to 6 tablespoons BANVEL 2S Herbicide with 1 gallon water for treatment of 1 to 2 square feet.	For control or suppression of weeds apply BANVEL 2S when weeds are actively growing.
2 to 4 gallons (4 to 8 pounds dicamba acid equivalent)	Apply as a preemergence application before weed emergence.

Continued on page 92

AND SIMILAR
AND
(RANGELAND)

REC

perennial ryegrass
small leaf brome
rough brome
sorghum
sorghum
yellow foxtail
stachys
water hyacinth

perennial ryegrass