RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

VELSICOL

BARRICADE

EMULSIFIABLE CONCENTRATE INSECTICIDE

STOP—READ LABEL BEFORE USING!

ACTIVE INGREDIENTS

O, O-Dimethyl O-p-nitrophenyl phosphorothioate* 29.83%

Related compounds 1.57%

O-Ethyl O-(p-nitrophenyl) phenylthiophosphonothioate* 31.40%

Xylene Range Aromatic Solvent 26.14%

INERT INGREDIENTS 11.06%

DANGER - POISON - PRECAUTIONS



POISONOUS IF SWALLOWED. [illegible]

POISONOUS BY SKIN CONTACT—Poisonous if touched by hands or spread on skin or clothing. Liquid goes through clothing.



POISONOUS IF BREATHED—Breathing vapors [illegible]

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

POISON

[illegible] IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

POISON SIGNS (Symptoms)

WARNING SYMPTOMS include weakness, headache, tightness in the chest, [illegible] blurred vision, nausea, abdominal cramps, diarrhea, sweating, [illegible] salivation, and nonreactive pinpoint pupils.

[illegible] swallowed, inhaled or absorbed through skin. Report to your [illegible] eyes.

[illegible] the eyes or skin [illegible] Do not [illegible] contact may [illegible] symptoms [illegible] burning.

[illegible] material must be removed immediately.

[illegible] seek prompt medical attention [illegible]

INFORMATION FOR PHYSICIAN

[illegible]

PRECAUTIONARY STATEMENTS
(Hazards To Humans and Domestic Animals Continued)

PROTECTIVE CLOTHING AND EQUIPMENT REQUIRED

Wear protective clothing, goggles, and respirator as defined by the American National Standards Institute when applying or handling, or when reentering treated areas within 48 hours of treatment. The following protective clothing must be worn: lightweight, unlined rubber gloves at least wristlength; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a nonpermeable material covering the body from ankles to wrists (clothing not required, natural rubber boots at least ankle length; full-face respirators are recommended; half-face respirators and goggles are required. Aerial applicators in positive pressure cockpits and other applicators in comparable grade equipment with appropriate filters at all air intakes (see ... with these protective clothing requirements.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift. For aerial application, flagging must be by fully automated mechanical means, or, if by humans, by persons working in totally enclosed vehicles.

Do not rub eyes or mouth with hands. If you feel sick in any way STOP work and get help right away. Call a doctor (physician, clinic or hospital)—immediately. Explain that the victim has been exposed to a Methyl Parathion-EPN combination and describe his condition. After first aid is given (see Statement of Practical Treatment) and if a doctor cannot come take victim to clinic or hospital.

IMPORTANT: Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK: take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. First aid should be cleaned and after replacement according to instructions provided with respirator.

ENVIRONMENTAL HAZARDS
POISONOUS TO FISH AND WILDLIFE

[illegible] Birds and other wildlife [illegible] Do not apply or wash equipment [illegible] Do not contaminate water [illegible] Do not dump excess or waste [illegible] ARE [illegible]

[illegible]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[illegible] apply this product when weather [illegible] is favorable to the treated area.

[illegible] Apply to blooming cotton if bees are visiting the treatment area.

Do not apply to corn during the pollen-shed period if bees are visiting the treatment area.

Re-entry Statement

Workers entering treated fields within 48 hours of application should wear protective clothing.

Where a dosage range is given, use the lower rate on small, young plants or where insect infestation is light; use the higher rates on large or mature plants or where insect infestation is heavy.

BARRICADE may be applied by ground or aerial application. When making ground application, apply the recommended dosage in at least 3 gallons of water per acre, using sufficient water to give good coverage of the foliage. For aerial application, use 3 to 10 gallons of water per acre.

To make spray solutions, fill half of the spray tank with water, agitate, add the proper amount of BARRICADE, then fill the spray tank with water while still agitating. Continue to agitate while using.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting of treated areas.

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
Cotton	Aphids Bollworm Cotton Leafhopper Cotton Leafworm Fall Armyworms Fleahoppers Garden Webworms Grasshoppers Lygus Bugs (and other Mirids) Red Spider Mites Thrips	2/3 to 1-1/3 pints	Make application at 5 to 7 day intervals until control is obtained. Do not apply within 7 days of boll-picking or harvest. Do not feed treated forage or gin trash to livestock.
	Darkling Ground Beetle Mexican Bean Beetle Pink Bollworm Plant Bug (Rapid Terminals) Russet Mites Saddleback Caterpillar Southern Armyworms Sugarbeet Webworms Two Spotted Mites	2 1/2 pints	
	Beet Armyworms Bollworm Cabbage Looper Cotton Leaf Perforator Cutworms (Larvae only) Tobacco Budworm Yellow Striped Armyworm	[illegible]	

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
Corn	Chinkbug Complex Corn Leaf Aphids Corn Rootworms (adult) Fall Armyworms Sandburs	1/2 pint	Repeat applications as necessary. Do not apply within 14 days of harvest.
	Armyworms (up to third instar)	2/3 pint	Make first application when 25% of the plants show evidence of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From 1 to 3 treatments may be required. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.
	European Corn Borer (treated rows)	2/3 to 3/4 pints	
	[illegible]	[illegible]	Make application 10 days after appearance of leaf feeding in [illegible] Repeat 7 to 10 days after if new evidence of leaf feeding is observed. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.
	[illegible]	[illegible]	Apply when 25% of the crop [illegible]

CROP	PEST	RATE OF BARRICADE PER ACRE	APPLICATION CONDITIONS AND LIMITATIONS
[illegible]	Aphids [illegible] Bird mites Flea Beetles Leafhoppers Leaf Bugs Mites	[illegible] pints	Do not apply within 21 days of harvest
[illegible]	Dancing Ground Beetle Mexican Bean Beetle Russian Mite Serpentine Leafminer Sugarbeet Webworm Thrips Tomato Fruitworm Two Spotted Mite	[illegible] pints	
Soybeans	Combined Cutworm	[illegible] pint	[illegible]
	Stink bugs Velvetbean Caterpillar	1 pint	[illegible] during the growing season Do not harvest or graze or feed forage within [illegible] days after the last application
	Aphids Blister Beetle Cotton Bollworm Garden Webworm Mexican Bean Beetle Three Cornered Alfalfa Hopper Two spotted Spider Mite	[illegible]	
	Cabbage Looper Corn Earworm Green Clovenworm	[illegible]	
	Fall Armyworm Beet Armyworm	[illegible]	

[illegible]