



BARRICADE

CONCENTRATE INSECTICIDE

STOP. READ LABEL
BEFORE USING.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply if a product when weather conditions favor drift from treated area.

Do not apply to blowing cotton if bees are visiting the treatment area.

Do not apply to corn during the pollen-shed period if bees are visiting the treatment area.

Pre-entry Restrictions

Workers entering treated fields within 48 hours of application should wear protective clothing.

Where a dosage range is given, use the 1 rate on small young plants or where insect infestation is light; use the higher rate on large or mature plants or where insect infestation is heavy. BARRICADE may be applied by ground or aerial application. When making ground application, apply the recommended dosage in at least 3 gallons of water per acre, using sufficient water to give good coverage of the foliage. For aerial application, use 3 to 10 gallons of water per acre.

To make spray mixture, fill half of the spray tank with water, agitate, add the proper amount of BARRICADE, then fill the spray tank with water until agitating. Continue to agitate while using.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting of treated areas.

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
CORN	Aphids Belted Girdler Cotton Leafhopper Cotton Leafworm Fall Armyworm Fleahopper Grasshopper Green Weeviler Lepidopteran (and other larvae) Red Spotted Mite Thrips	2/3 to 1-1/2 pints	Make application at 5 to 7 day intervals until control is obtained. Do not apply within 7 days of hand-picking or harvest. Do not feed treated leaves or get them in livestock.
	Darkwing Ground Beetle Mexican Bean Beetle Pine Sawfly Plant Bug (Rhopal. Tortricid) Rust Mite Scorpionfly (Lepidopteran) Southern Armyworm Soybean Weevil Two Spotted Mite	2-1/2 pints	
	Beet Armyworm Cutworm Cotton Leaf Perforator Saltmarsh Caterpillar Tobacco Budworm Yellow Striped Armyworm	1-1/2 pints	

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
CORN	Climbing Cutworm Corn Leaf Aphid Corn Fleahopper (adult) Fall Armyworm Sawfly	1/2 pint	Repeat application as necessary. Do not apply within 14 days of harvest.
	Armyworm (up to third instar)	2/3 pint	Make first application when 25% of the plants show evidence of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From 1 to 3 treatments may be required.
	European Corn Borer (young stage)	2/3 to 1-1/4 pints	For control of second brood larvae apply when eggs begin to hatch. Do not apply within 14 days of harvest.
	European Corn Borer (old stage)	1 pint	Make application 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is observed. For control of second brood larvae apply when eggs begin to hatch. Do not apply within 14 days of harvest.

to be used in accordance with the label instructions.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Metal

Triple rinse immediately and thoroughly with water. Empty containers should be disposed of by procedures approved by state or local government officials.

Plastic

Triple rinse immediately and thoroughly with water. Empty containers should be disposed of by procedures approved by state or local government officials.

Small, unidentifiable fragments of plastic and metal are scattered across the bottom of the page, likely from a container or equipment used during the scanning process.