


FOR THE CONTROL OF PESTS  
USE ONLY BY CERTIFIED APPLICATORS  
OR UNDER THEIR DIRECT SUPERVISION AND ONLY ON  
THOSE USES COVERED BY THE CERTIFIED APPLICATOR REGISTRATION  
CATION



**VELSICOL**  
**BARRICADE**  
EMULSIFIABLE CONCENTRATE INSECTICIDE

**STOP READ LABEL  
BEFORE USING**

ACTIVE INGREDIENTS  
PERCENTAGE BY WEIGHT  
EPA REGISTRATION NO. 100-100000000-1  
EPA ESTABLISHED IN 1972

**DANGER-POISON-PRECAUTIONS**



**POISONOUS IF SWALLOWED**—The product can be fatal if swallowed. Do not swallow.

**POISONOUS BY SKIN CONTACT**—Poisonous if touched by hands or applied or spread on or over the face or clothing. Liquid goes through clothes.

**POISONOUS IF BREATHED**—Breathing vapors from the product may be fatal.

**PRECAUTIONARY STATEMENTS**  
**HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**



**POISON**

(SEE LIST OF SUSPECTED POISONING SYMPTOMS)

... symptoms of the product are: dizziness, drowsiness, headache, sweating, ...

... (SEE LIST OF SUSPECTED POISONING SYMPTOMS)

**PRECAUTIONARY STATEMENTS**  
**(Hazards To Humans and Domestic Animals Continued)**

**PROTECTIVE CLOTHING AND EQUIPMENT REQUIRED:**

Wear clean protective clothing, goggles, and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within 48 hours of treatment. The following protective clothing must be worn: lightweight, unlined natural rubber gloves at least midforearm in length; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a nonpermeable, non-pleated material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least mid-calf in length; full-face respirators are recommended; half-face respirators and goggles are required. Aerial applicators in positive pressure suits and other applicators in comparable ground equipment with appropriate filters at all air intakes need not comply with these protective clothing requirements.

**WORK SAFETY RULES**

Keep all unprotected persons and children away from treated area or where there is danger of drift. For aerial application, flagging must be by fully automated mechanical means, or, if by humans, by persons working in totally enclosed vehicles.

Do not rub eyes or mouth with hands. If you feel sick in any way STOP work and get help right away. Call a doctor (physician) clinic or hospital—immediately. Explain that the victim has been exposed to a Methyl Parathion-EPN combination and describe his condition. After first aid is given (see Statement of Practice: Treatment), and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT:** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK:** take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

**ENVIRONMENTAL HAZARDS**  
**POISONOUS TO FISH AND WILDLIFE**

This is toxic to fish and wildlife. Birds and other wildlife in treated areas may be poisoned by contact with any body of water. Do not apply when weather conditions favor runoff or a rain shower. Do not apply where runoff is likely to occur. Do not contaminate waterways with equipment or disposal of wastes. **APPENDIX THIS PRECAUTIONARY STATEMENT ON THIS LABEL.**

**PHYSICAL OR CHEMICAL HAZARDS**

AND OTHER INFORMATION CONCERNING THE HAZARDOUS NATURE OF THIS PRODUCT

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product when weather conditions favor drift from treated area.

Do not apply to blooming cotton if bees are visiting the treatment area.

Do not apply to corn during the pollen-shed period if bees are visiting the treatment area.

**No entry Statement**

Workers entering treated fields within 48 hours of application should wear protective clothing.

Where a dosage range is given, use the lower rate on small, young plants or where insect infestation is light; use the higher rate on large or mature plants or where insect infestation is heavy. BARRICADE may be applied by ground or aerial application. When making ground application, apply the recommended dosage in at least 3 gallons of water per acre, using sufficient water to give good coverage of the foliage. For aerial application, use 3 to 10 gallons of water per acre.

To make spray mixtures, fill half of the spray tank with water, agitate, add the proper amount of BARRICADE, then fill the spray tank with water while still agitating. Continue to agitate while using.

**POST TREATED AREA**

Consult your State Agricultural Extension Service or Experiment Station regarding posting of treated areas.

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
Cotton	Aphids Belt Weevil Cotton Leafhopper Cotton Leafworm Fall Armyworm Hobnobbler Garden Webworm Grasshopper Lemon Drop (and other species) Red Spined Miner Thrips	2/3 to 1-1/2 pints	Make application at 5 to 7 day intervals until control is obtained. Do not apply within 7 days of hand-picking or harvest. Do not treat treated crops or old trees in treatment.
	Dormant Ground Beetle Mole on Root Beetle Pink Bollworm Pine Sawfly (Royal, Tawny-bellied) Sawfly Miner Sawyerfly Larva Southern Armyworm Superbloom Webworm Two Spotted Miner	2-1/2 pints	
	Beet Armyworm Borer Cabbage Looper Cotton Leaf Perforator Sulphur Caterpillar Tobacco Borer Yellow Striped Armyworm	2-2/3 pints	

CROP	INSECT	RATE OF BARRICADE PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
Corn	Chickling Cutworm Corn Leaf Aphid Corn Rootworm (adult) Fall Armyworm Sandhopper	1/2 pint	Repeat application as necessary. Do not apply within 14 days of harvest.
	Armyworm (up to third instar)	2/3 pint	Make first application when 75% of the plants show evidence of leaf feeding and damage of 5 day intervals as long as leaf feeding is evident. Repeat 1 to 2 treatments only as needed.
	European Corn Borer (second instar)	2/3 to 3/4 pints	For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.
	European Corn Borer (third instar)	2/3 pint	Make application 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is observed. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.
	European Corn Borer (fourth instar)	1 pint	Apply when 10% of the plant population has eggs, masses. Repeat at 5 to 10 day intervals as needed. Do not apply within 14 days of harvest.

17-1111-1013

CROP	INSECT	RATE OF APPLICATION PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS
Beans (Dry green bean)	Aphids Armyworms (up to third instar) Flour Beetles Leafhoppers Lygus bugs Mites	1-1/2 pints	Do not apply within 21 days of harvest
Beans (Dry green)	Darkling Ground Sooty Mexican Bean Beetle Rust mite Soybean Leafminer Soybean Weeviler Thrips Tomato Fruitworm Two-Spotted Mite	1-1/2 pints	
Soybeans	Chewing Caterpillars	2/4 pint	Use restricted to the southeast Do not make more than 2 applications during the growing season Do not harvest or grade or feed forage from treated beans to livestock within 21 days after the last application
	Stinkbugs Vespaevine Caterpillars	1 pint	
	Aphids Bean Beetle Cotton Bollworm Garden Webworm Mexican Bean Beetle Three Corned Acker Hopper Two-spotted Spider Mite	1-1/2 pints	
	Cabbage Looper Caterpillar Cotton Bollworm	1 pint	
	Full Armyworm (up to third instar)	1 2/3 pints	
Tomatoes	Florida Aphid Soybean Leafminer Southern Armyworm Tomato Fruitworm	1-1/2 to 1 1/2 pints	Do not apply within 21 days of harvest
	Mites Phytophthora Rust mite Soybean Seed Ackerworm Thrips Two-spotted Mite Yellow Spider Armyworm	1 1/2 pints	
	1/2 Beetle Soybean Looper	1 1/2 pints	

AND MIXTURES FOR OTHER

... ..

... ..

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Metal

Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or (if allowed) state and local authorities or burning. If burned, stay out of smoke.

LIMITED WARRANTY AND LIABILITY

NOTICE: See this limited warranty on the back of the container. If the label is not placed on the container, the user should refer to the back of the container for the full text of this warranty. This warranty is made by the manufacturer of the product and is not a contract. It is subject to the terms and conditions of the purchase order and the terms and conditions of the product label. The manufacturer warrants that the product is free from defects in material and workmanship under normal use and handling for a period of one year from the date of purchase. The manufacturer does not warrant the product against damage caused by fire, theft, or other causes. The manufacturer does not warrant the product against damage caused by use of the product in a manner not intended in the label directions or from mixing this product before use with a substance not recommended by the product label. The manufacturer does not warrant the product against damage caused by use of the product in a manner not intended in the label directions or from mixing this product before use with a substance not recommended by the product label. The manufacturer does not warrant the product against damage caused by use of the product in a manner not intended in the label directions or from mixing this product before use with a substance not recommended by the product label. The manufacturer does not warrant the product against damage caused by use of the product in a manner not intended in the label directions or from mixing this product before use with a substance not recommended by the product label.