

Handwritten notes:
 New 16/1/74
 10/11/74
 10/11/74

**MILKING MACHINES
 AND UTENSILS
 CLEANING**

1. Immediately after use, rinse all equipment with cold or lukewarm water until all milk is flushed out. Do not use hot water.
2. Add 1 oz. BONCHEM BON-A-CIDE to each 5 gallons of lukewarm (90°F.) water. This provides 25 parts per million titratable iodine.
3. Dismantle milking machine and immerse parts, hoses, and inflations (but not pulsators) before brushing. Wash pails and utensils in same solution. Rinse with clear water and drain.

SANITIZING

1. Before using cleaned equipment sanitize with a solution consisting of 1 oz. BONCHEM BON-A-CIDE for each 5 gallons of lukewarm (90°F.) water, providing 25 parts per million titratable iodine. Immerse equipment in this solution for at least 2 minutes and drain.

CAUTION: Harmful if swallowed. Causes skin irritation. Avoid contact with skin and eyes. In case of contact flush skin or eyes with plenty of water for at least 15 minutes. For eyes get medical attention. Avoid contamination of foods.

DILUTION TABLE

| Ounces of BON-A-CIDE | Gallons of Water | Parts per million titratable iodine |
|----------------------|------------------|-------------------------------------|
| 1 | 5 | 25 |
| 2 | 5 | 50 |
| 4 | 5 | 100 |

When used at over 25 parts per million titratable iodine, potable water rinse is required before exposing equipment to food.

**BON-A-CIDE
 IODINE - DETERGENT - SANITIZER**

ACTIVE INGREDIENTS:

- a-(p-nonylphenyl)-omega-hydroxy poly (oxyethylene)-iodine complex, providing 1.75% titratable iodine 14%
- Phosphoric acid 15%

INERT INGREDIENTS:

- Water 71%
- Total 100%

**CAUTION: Keep out of reach of children. See side panels for further cautions.
 1 -- U. S. GALLON**

FARM BULK TANKS

1. Rinse tank thoroughly with lukewarm water until all visible milk is flushed out.
2. Brush tank with a solution of 1-oz. BONCHEM BON-A-CIDE for each 5 gallons of lukewarm (90°F.) water, providing 25 parts per million titratable iodine. Rinse and drain.
3. Before use, brush or spray (where code permits) the tank with a solution of 1/2-oz. BONCHEM BON-A-CIDE to a 10 quart pail of water, providing 25 parts per million titratable iodine. Allow 5 minutes contact and drain.

MILKSTONE REMOVAL

Mix equal parts of BONCHEM BON-A-CIDE and lukewarm (90°F.) water. Spread over surface to be cleaned, let stand a few minutes, then brush thoroughly and rinse.

UDDERS

For each cow use a fresh paper towel wet in a solution of 1/2-oz. BONCHEM BON-A-CIDE to 10 quarts of lukewarm water, providing 25 parts per million titratable iodine.

The treated equipment shall be so stored that all residual bactericide is drained before being used in contact with food products.

This solution must be discarded when its concentration reaches 12.5 ppm.

Rinse empty container thoroughly with water and discard it.

KEEP STOPPERED IN A COOL PLACE. Always follow Local Health Dept. regulations.

USDA Reg. No. 887-21
 U.S. Pat. Nos. 2,931,777 - 2,977,315
 3,028,299