

(Side Panel)

USE DIRECTIONS FOR UD-100

FOR SANITIZING

Prepare a solution of sufficient volume by adding 1 ounce of UD-100 to every 5 gallons of water used. Sanitize equipment by circulation, immersion, or spraying. Allow a 2 minute exposure period or contact time specified by governing sanitary code. Do not rinse unless required by local health authorities, and then only in accordance with established regulations.

FOR UDDER WASHING

Prepare a solution in a stainless steel or plastic bucket using 1 ounce of UD-100 with every 5 gallons of warm (115° - 120°F.) water. Use single service, disposable paper towels to bathe the areas to be cleaned and sanitized.

FOR RESTAURANTS AND INSTITUTIONS

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with Diversey cleaner of your choice.
3. Rinse with clean water.
4. Sanitize in a UD-100 solution containing 1 ounce per every 5 gallons of water. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.

NOTE: A clean potable water rinse following sanitation is not permitted under Section H 96.16 (2) (c) 2 of the Wisconsin Administrative Code.

CAUTION: This is a moderately acidic material. Do not take internally. If swallowed, give 1 or 2 glasses of water or milk. Call doctor. If product gets in eyes, or on skin, flush freely with water. Consult physician. Avoid contamination of foods.

## GENERAL DIRECTIONS

All surfaces to be treated should be thoroughly cleaned before using BONCHEM MIKRO-CIDE sanitizing solutions.

For all sanitizing applications, use 1 ounce of BONCHEM MIKRO-CIDE per gallon of water. BONCHEM MIKRO-CIDE is recommended for the sanitizing of all dairy and food plant equipment. Rinse equipment with potable water before use.

This product when used as directed fulfills the criteria of Appendix F of the Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the U. S. Public Health Service.

**NOTE:** Always follow State and local Health Department cleaning and sanitizing requirements.


# MIKRO-CIDE<sup>®</sup>

## Anionic Acid Sanitizer

<b>Active Ingredients:</b>	
Orthophosphoric Acid .....	22.5%
Glycolic Acid .....	14.0%
Decyl Benzene Sulfonic Acid, Sodium Salt .....	2.7%
<b>Inert Ingredients*</b> .....	<b>60.8%</b>
<b>*Includes Nonionic Detergent</b>	
	<b>100.0%</b>

**CAUTION:** Keep out of reach of children. See side panel for further cautions.

**Contents: One U. S. Gallon**



# MIKRO-CIDE<sup>®</sup>

## Anionic Acid Sanitizer

<b>Active Ingredients:</b>	
Orthophosphoric Acid .....	22.5%
Glycolic Acid .....	14.0%
Decyl Benzene Sulfonic Acid, Sodium Salt .....	2.7%
<b>Inert Ingredients*</b> .....	<b>60.8%</b>
<b>*Includes Nonionic Detergent</b>	
	<hr/>
	<b>100.0%</b>

**CAUTION:** Keep out of reach of children. See side panel for further cautions.

**Contents: One U. S. Gallon**

### SANITIZING DIRECTIONS

**Tank Sanitizing** (where code permits): Use one ounce of BONCHEM MIKRO-CIDE per gallon of water and a minimum contact time of 5 minutes. Rinse with potable water.

**Immersion or Flowing Methods:** Use one ounce of BONCHEM MIKRO-CIDE per gallon of water and a minimum contact time of 2 minutes. Rinse with potable water.

**CAUTION:** Harmful if swallowed. Causes skin and eye irritation. Avoid contact with skin and eyes. In case of contact, flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. Avoid contamination of foods.

**USDA Reg. No. 887-24**

ACCEPTED WITH COMMENTS

