

DIRECTIONS FOR USE

SANITIZATION OF BEVERAGE CONTAINERS

Follow local Health Department regulations, which usually suggest use solutions in the range of the lower ppm levels shown in the Dilution Table below.

Directions for Dilution

	Titratable Iodine
½ oz. BONCHEM NO RINS IODINE per 10 gal. water	6¼ ppm
1 oz. BONCHEM NO RINS IODINE per 10 gal. water	12½ ppm
2 oz. BONCHEM NO RINS IODINE per 10 gal. water	25 ppm
4 oz. BONCHEM NO RINS IODINE per 10 gal. water	50 ppm

CAUTION:

Do not get in contact with food.

Harmful if swallowed.

Avoid contact with skin and eyes. In case of contact with eyes, flush immediately with water; get medical attention.

U.S. Pat No. 3,028,299
EPA Reg. No. 887-33

NO RINS IODINE

Controls Bacteria, Mold, Yeast

Active Ingredients

Butoxypolypropoxypolyethoxyethanol

Iodine Complex (providing

1.6% titratable iodine) 12.54%

Inert Ingredients 87.46%

Total 100.00%

FOR INDUSTRIAL USE ONLY.

CAUTION:

Keep Out Of Reach Of Children.
See additional precautions on left panel.
One U.S. Gallon

DIRECTIONS FOR USE

EQUIPMENT SANITIZATION

Preclean equipment.

Fill with solution 2 oz. BONCHEM NO RINS IODINE to 10 gallon water. (Provides 25 ppm titratable iodine.)

When ready to use equipment, drain off BONCHEM NO RINS IODINE solution.

TANK AND VAT SANITIZATION

Preclean.

Add solution 2 oz. BONCHEM NO RINS IODINE to 10 gallon water. (Provides 25 ppm titratable iodine.)

After five minutes, drain off solution.

SANITIZATION OF WALLS, FLOORS, CONVEYORS, ETC.

Clean daily with solution 2 oz. BONCHEM NO RINS IODINE to 10 gallon water (25 ppm titratable iodine.)

Clean weekly or at batch run end with solution 4 oz. BONCHEM NO RINS IODINE to 10 gallon water (50 ppm titratable iodine) and rinse.

11-15-71
887-33