

ONS  
ould be  
re using  
D sanitiz-  
izing ap-  
-SAN  
r. After  
to allow  
zer solu-  
it shows  
er 2 gal-  
est Kit).  
directed  
endix F  
zed Milk  
ndations  
Service.  
ate and  
cleaning  
ts.  
roughly  
39

# BON-SAN 100

## Anionic Acid Sanitizer

|   |         |
|---|---------|
| <b>Active Ingredients:</b>  |         |
| Phosphoric Acid. . . . .  | 16.50%  |
| Dodecylbenzenesulfonic acid . . . . .   | 2.90%   |
| <b>Inert Ingredients:*</b> . . . . .  | 80.60%  |
| *Includes polyoxyethylene polyoxypropylene block copolymer and alpha-hydroxypropionic acid. |         |
|   | 100.00% |

**DANGER:** Keep Out Of Reach Of Children.  
See Side Panel For First Aid And Additional  
Precautions.

**CONTENTS: ONE U.S. GALLON**

### SANITIZING DIRECTIONS

For sanitizing, food equipment, dairy equipment and utensils.

**Spraying (where code permits):** Use one ounce of BONCHEM BON-SAN 100 to 1 gallon of water and a minimum contact time of 5 minutes. Allow equipment to drain before using.

**Immersion or Flowing Methods:** Use one ounce of BONCHEM BON-SAN 100 to 1 gallon of water and a minimum contact time of 2 minutes. Allow equipment to drain before using.

**DANGER:** Corrosive to skin. Causes eye damage. Do not get in eyes, on skin or clothing. Protect eyes and hands when handling. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**ACCEPTED**  
8 AUG 1975