



HERCULES

TOXAPHENE

ACTIVE INGREDIENTS:**TOXAPHENE (TECHNICAL CHLORINATED CAMPHENE CONTAINING 67-69% CHLORINE) 100%**

AN INSECTICIDAL TOXICANT

FOR MANUFACTURING USE ONLY



**POISON
GIFT
VELENO VENENO**

FIRST AID—ANTIDOTE

INTERNAL: Give sodium chloride 15 g. (one table-spoonful of table salt) in glass of warm water and repeat until vomit is clear.

EXTERNAL: Wash with soap and water.

CALL A PHYSICIAN IMMEDIATELY

PHYSICIAN'S NOTE: Toxaphene produces central nervous system excitation which responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons," which was prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

WHEN EMPTY, DESTROY THIS CONTAINER BEYOND ALL USE. BECAUSE OF THE POSSIBILITY OF POISONOUS MATERIAL ADHERING TO THIS CONTAINER, IT SHOULD NOT BE REUSED FOR ANY PURPOSE.

NET CONTENTS 250 POUNDS

USDA REG. NO. 891-1

HER. 19214

PRINTED IN U.S.A.



Hercules Incorporated makes no warranty, express or implied, concerning the use of this product other than that indicated on the label.

SYNTHETICS DEPARTMENT • HERCULES INCORPORATED

WILMINGTON, DELAWARE U.S.A.