

TOXAPHENE

20% DUST INSECTICIDE

ACTIVE INGREDIENT:	Percent by Weight
Toxaphene (technical chlorinated camphene containing 67-69% chlorine)	20.00
INERT INGREDIENTS:	80.00
TOTAL	100.00

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Read side panel for additional handling precautions.

EPA Registration No. 891-29

CAUTION

Harmful if swallowed. Avoid breathing of dust mist. Avoid contact with eyes, skin, and clothing. Wash thoroughly after using. Avoid contamination of feed or foodstuffs.

Keep away from children and domestic animals. Do not pasture dairy animals or animals being fattened for slaughter in treated fields. Do not use on cucurbits.

This product is toxic to fish, birds, and other wildlife.

Keep out of any body of water.

Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ANTIDOTE

Internal— Give sodium chloride 15 g. (one tablespoonful of table salt) in a glass of warm water and repeat until vomit is clear. Call a doctor immediately.

External— Wash with soap and water.

PHYSICIAN'S NOTE: Toxaphene produces central nervous system excitation which responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons," which was prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

DIRECTIONS FOR USE

Grain Sorghum: For control of climbing and surface-feeding cutworms, corn earworm, armyworms, and grasshoppers, apply at the rate of 7½ to 15 pounds per acre. Do not apply more than once after heads have begun to form.

Do not pasture dairy animals or animals being fattened for slaughter in treated fields. Do not ensile treated forage.

Do not apply within 4 weeks of harvest when using up to 10 pounds per acre.

Do not apply within 40 days of harvest when using 10 to 15 pounds per acre.

Collards and Kale: For control of climbing and surface feeding cutworms, thrips, serpentine leaf miner, armyworms, cabbage-worms, and green peach aphid, apply at the rate of 10 pounds per acre. For control of cabbage looper, apply at the rate of 10 to 20 pounds per acre. Do not apply within 14 days of harvest at the low rate. Do not apply within 21 days of harvest when using over 10 pounds up to 20 pounds per acre.

Spinach: For control of climbing and surface-feeding cutworms, thrips, serpentine leaf miner, armyworms, cabbage-worms, green peach aphid, and cabbage looper, apply at the rate of 10 pounds per acre. Do not apply more than once. Do not apply within 21 days of harvest.

ACCEPTED

JULY 24, 1971

891-29

Herco warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is reasonably fit for its intended purpose. Since timing, method of application, weather,