

**
WEED RHAP-20
Granular 2,4-D

WEED RHAP-20* Granular 2,4-D is formulated by impregnating selected mineral granules with low-volatile 2,4-D ester, to minimize drift. It is intended for use in the control of certain water weeds infesting lakes and ponds. It is also useful in the control of certain broad-leaved weeds infesting grasses and crops such as corn.

DIRECTIONS FOR USE

In Lakes and Ponds: To control Milfoil, Hornwort, Bladderwort, Fanwort, Pondweed, Cattail and Water Lillies infesting lakes and ponds, apply granules at the rate of 1 to 1-1/2 lbs. per 430 sq. ft. (10'x43') of water surface or 100 to 150 lbs. per acre. Distribute the granules evenly over the quiet water surface infested with weeds, using a dry fertilizer spreader attached to a boat. The granules settle to the water bottom around the roots and stems. Effects on waterweeds will be evidenced in 4 to 10 weeks and control may be expected for a year or longer. In running water, results are less satisfactory. DO NOT apply to more than 1/3 to 1/2 of a lake or pond in any one month. In tropical regions use 25 to 50 percent more granules.

In Yards and Parkways: To control broad-leaved weeds such as Wild Onion, Dandelion, Plantain, Ivy, Bitterweed and Dog Fennel, apply 10 to 15 lbs. of granules per acre in early spring when the shoots are young, using a dry fertilizer spreader. DO NOT use more than recommended; do not spread within 3 ft. of shrubs or flowers. Repeat application in 1 month to control skipped spots. Weeds over 6 inches tall are sometimes resistant and should be sprayed with R-H Low Volatile Weed Rhap.

In Corn: To aid control of annual grasses such as large and small Foxtail, Crab Grass, Barnyard Grass; broad-leaved weeds such as Smartweed, Sunflower, Wild Mustard, Cocklebur, Bindweed, Dandelion and Morning Glory; apply as follows: With a granule applicator designed to apply a band over the planted row; apply 3 to 3-1/4 lbs. per acre in a band 14 inches wide when rows are 42 inches apart. If the planting varies from this standard band width, or row separation, the granules should be applied proportionately. Be sure the granules are allowed to stay on top of the soil; preferably, they should be applied behind the packer wheel at planting time. Be sure the corn is planted 2-1/2 to 3 inches deep as damage may occur to shallow planted corn. DO NOT apply on sandy soil. (For Corn Belt Only)

In Strawberries: To control weeds such as Crab Grass, Foxtail, Barnyard or Winter Grass, Horse Weed, Bull Nettle, Carless Weed, Morning Glory, Dock, Smartweed, Jimson Weed and Dandelion: Apply at the rate of 12 lbs. per acre over the planted area. Application should be made broadcast using a spreader-seeder previously adjusted to deliver 12 lbs. per acre. Application should be made only in the dormant season (after first fall frost or before new leafing in spring).

On Cranberries: For control of annual broadleaf weeds in bearing cranberry marshes, apply at a rate of 10 to 20 lbs. per acre (2 to 4 lbs. 2,4-D equivalent per acre). Application should be made broadcast in the spring after removal of the winter flood from the dormant vines. Weeds controlled are Common Ragweed, Sticktites, Tearthumb, Smartweed, Northern Saint Johnswort, and common species of Annual Smartweed. Careful distribution of the granules is necessary if adequate control is to be obtained. Granules may also be distributed on ice over plants prior to spring thaw. (For Wisconsin area)

WARNING

Avoid application to susceptible plants because this product may injure cotton, soybeans, tomatoes, beans, peas, fruit trees, ornamentals, etc. Do not allow particles or dust to drift onto desirable crops or plants. Avoid contamination of water intended for irrigation or domestic use. Do not store near fertilizers, seeds, insecticides, or fungicides. It is best to use separate equipment for applying 2,4-D granules. Should the same equipment be used for applying other materials, however, clean it thoroughly before such use to remove all traces of 2,4-D particles or dust. Keep bag closed when it is not in use. Dispose of empty bag by burying or burning. Do not burn where smoke or fumes will drift to desirable crops or plants.

Avoid contact with skin, eyes, and clothing. In cases of skin contact, wash thoroughly with soap and warm water. Flush eyes with clear water and obtain prompt medical attention.

NON-WARRANTY: Because this product will be used under many different conditions over which the manufacturer has no control, we do not guarantee the suitability of this product in any particular situation. This product is sold without warranty, express or implied, and buyer assumes all responsibility for loss or damage arising from the handling and use of this product, whether in accordance with directions or not.

ACTIVE INGREDIENTS:

2-ethyl-hexylester* of 2,4-dichlorophenoxyacetic acid	30.12%
---	--------

INERT INGREDIENTS:

Clay Granules	69.88%
TOTAL	100.00%

* Equivalent to 20% 2,4-dichlorophenoxyacetic acid.

Net Contents - 50 lbs.

Contents licensed under U.S. Patent No. 2,792,295.

** Hercules Trademark

Made By

Agricultural Chemicals

Synthetics Department

HERCULES POWDER COMPANY

Incorporated

Wilmington, Delaware, U.S.A.

