


# TOXAPHENE

AN INSECTICIDE ESPECIALLY PREPARED FOR  
EUROPEAN CORN BORER CONTROL



ACTIVE INGREDIENT:	Percent by Weight
Toxaphene (technical chlorinated camphene containing 67.69% chlorine)	10
INERT INGREDIENTS:	90
TOTAL	100

**CAUTION:** KEEP OUT OF REACH OF CHILDREN.

Read side panel for additional handling precautions

MANUFACTURED BY  
SYNTHETICS DEPARTMENT • AGRICULTURAL CHEMICALS

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe dust. Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Empty bags should be buried or burned. (Stay away from market. If the bags are burned, the ashes should be buried in a safe place.)

**Internal** Give sodium chloride 15 g (1 tablespoonful of table salt in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

**External** Wash with soap and water.

**European Corn Borer Control** Apply 20 pounds per acre by either ground equipment or airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second brood here. Consult State Extension Service specialists for most effective application dates.

Do not feed milage to be treated with this product to dairy animals, including those being milked for 24 hours before milking. Do not feed to dairy animals or use on pastures or ranges for 24 hours after application. Do not use on pastures or ranges for 24 hours after application.

Do not use on pastures or ranges for 24 hours after application. Do not use on pastures or ranges for 24 hours after application. Do not use on pastures or ranges for 24 hours after application. Do not use on pastures or ranges for 24 hours after application.