

[871-150]

HERCULES TOXAPHENE
65% Emulsifiable Livestock Concentrate

An insecticide concentrate especially prepared for livestock use.

<u>ACTIVE INGREDIENTS:</u>	<u>Percent By Wt.</u>
Toxaphene (Technical chlorinated camplene containing 67-69% chlorine)	65.0
Refined Petroleum Distillate	22.5
<u>INERT INGREDIENTS:</u>	<u>12.5</u>
TOTAL	100.0%

WARNING: KEEP OUT OF REACH OF CHILDREN. Read Entire Label for Other Handling Precautions.

WARNING: May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist.

CAUTION: Use only as specified on the label. Avoid contamination of feed or foodstuffs. Do not use on dairy animals, in dairy barns or in milk rooms.

Do not use on poultry or in poultry houses.

Do not use on animals within 28 days of slaughter.

Do not use on calves less than 3 months of age.

Do not use on dairy goats.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply only as specified on the label. Rinse equipment and dispose of wastes by burying on non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying.

ANTIDOTE: Internal - Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

External - Wash with soap and water.

ACCEPTED

OCT 7 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
REGISTERED TOXIC POISON REGISTER
EPA NUMBER NC 711-50 SUBJECT
BY ATTACHED COMMENTS

DIRECTIONS FOR USE:

Cattle

Ticks - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. Thorough wetting of the animal is important for best results. Apply diluted material at the rate of 6 quarts per animal. Repeat application when necessary.

Horn Fly - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. Apply diluted material at the rate of two quarts per animal. Repeat application when necessary.

Lice - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. Apply diluted material at the rate of six quarts per animal. A single thorough application is usually all that is necessary to control lice.

Sheep and Goats

Lice - Sheep Tick (Keds) - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. A single thorough application is usually all that is required for control of lice and keds.

Ticks - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. Repeat application when necessary.

Hogs

Lice - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. A single thorough application is usually all that is required for control of lice.

Sarcoptic Mange - Mix Hercules toxaphene livestock concentrate at the rate of one gallon to 140 gallons of water. Repeat application when necessary.

NONWARRANTY:

Hercules warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant,

NONWARRANTY (contd.)

and soil conditions, mixtures with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Net Contents: _____

USDA Registration No. 891-26

MANUFACTURED BY:

SYNTHETICS DEPARTMENT · · AGRICULTURAL CHEMICALS

HERCULES INCORPORATED
WILMINGTON, DELAWARE, U.S.A.