

ACCEPTED

A Granular Formulation of Herbicide Amiben  
For Pre-emergence Weed Control in Soybeans

ACTIVE INGREDIENTS	% By Weight
Norco <sup>†</sup> (3-(benzylhydro-4,7-methoxy-1,2,4-oxadiazol-5-yl)-1,1-dimethylurea)	3.8
Reaction products related to Norco	0.2
*Ammonium salt of amiben (3-amino-2,5-dichlorobenzoic acid)	5.4
INERT INGREDIENTS:	90.6
TOTAL	100.0

†U. S. Patent No. 3,304,167

\*Contains 5% equivalent of  
3-amino-2,5-dichlorobenzoic acid.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

Read Side Panel For Use Directions And Additional Precautions.

Wash thoroughly after handling.

Harmful if swallowed.

Avoid inhalation of dust.

Do not contaminate food or foodstuffs.

Avoid contact with skin.

Do not get in eyes.

~~Herbicide~~ 30 granular KFF herbicide granules and broadcast of weeds in soybeans. Since this herbicide is applied before weed emergence and will be active only after rainfall applies, rain or irrigation is required for herbicide to be active. Do not apply to dry soil. Apply to soil before weeds emerge. Apply to soil before weeds emerge.

Weeds Controlled

~~Herbicide~~ 30 granular controls the following annual grasses and broadleaf weeds: foxtails (green, giant, yellow), crabgrass, barnyard grass, fall panic, summer panic, green goosegrass, pigweed, lambquarters, knotweed, purslane, chickweed, and ryegrass.

Large seeded weeds such as morning glory, cocklebur, Jimson weed and velvetleaf often germinate below the penetration level of the herbicide.

Populations of these weeds which are not immediately reduced under normal rainfall conditions.

DEVELOPS

Apply ~~Herbicide~~ 30 granular to the soil surface after planting and before soybeans and weeds emerge (pre-emergence).

Use 20 pounds of ~~Herbicide~~ 30 granular broadcast per acre on heavy and silt loam soils. On heavier soils such as clay, clay loam, or silty clay with high organic content, apply 30 pounds of ~~Herbicide~~ 30 granular broadcast per acre. Use proper placement for best application. A broadcast application can be mounted on the planter or the applicator used in planting can be used during operation.

Plant soybeans at a minimum depth of 1-1/2 inches. Do not use ~~Herbicide~~ 30 granular on sandy soils, heavy sands or deep soils.

**CAUTION:** Apply to soybeans to be harvested for the feed market. Do not graze the field or use the field for feed purposes.

*Handwritten notes:*  
Apply to soil before weeds emerge.  
Do not apply to dry soil.