



# TORBIDAN®

## 6-1.5

ACTIVE INGREDIENTS:

Percent by Weight

Permethrin 25.0%  
Cyfluthrin 1.5%  
Inert Ingredients 73.5%

INERT INGREDIENTS:

Water 68.0%  
Xylene 1.0%  
Nonylphenol 2.0%  
Fatty Acid Soap 1.5%  
Surfactant 1.0%  
Polyethylene Glycol 1.0%  
Other Inerts 1.0%



**POISON**



ALL INFORMATION ON THIS PRODUCT  
OBTAINED FROM THE MANUFACTURER  
OR THE DISTRIBUTOR  
IS THE PROPERTY OF THE MANUFACTURER

**DANGER: KEEP OUT OF REACH OF CHILDREN**

**NET CONTENTS 55 GALLONS**

remove clothing. Wash thoroughly with soap and water. Flush with plenty of water. Do not eat or drink until flushed.

Wash hands thoroughly with soap and water. Do not eat, drink, or use tobacco until hands are washed.

**DO NOT** breathe dusts of a type passed by the U. S. Bureau of Mines as being a union protector.

**KEEP** this product away from children and away from or vicinity where there may be a general population. Contaminated clothes should not be reentered until the spray has dried. If clothes does have dissipated. If treated clothes are used during or after use, get out of contaminated area at once for medical attention.

## POISON

**ANTIDOTE—IF SWALLOWED:** Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately.

**IF ON SKIN:** In case of contact remove contaminated clothing and immediately wash skin with soap and water.

**PHYSICIAN'S NOTE:** Methyl parathion exerts muscarinelike effects—i.e., stimulation of the parasympathetic nervous system. Repeated use of a parasympathetic depressant, such as atropine, to the limit of patient's tolerance is indicated to relieve symptoms. 2-PAM is also antidotal and may be administered in conjunction with atropine. The administration of oxygen is helpful. Never give morphine to patients suspected of parathion poisoning. Toxaphene produces central nervous system excitation that ordinarily responds to the barbituric acid derivatives, but these drugs should be considered only if complete atropinization fails to relieve symptoms. Refer to "Clinical Handbook on Economic Poisons," prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

### GENERAL DIRECTIONS

Prepare a spray mixture of Torbidan 6-1-5 as follows.

1. Add about half the total amount of water to be used to the spray tank.
2. Start agitator or pump bypass and add proper amount of Torbidan 6-1-5.
3. Continue mixing and add remaining water.

### DIRECTIONS FOR USE ON COTTON

Torbidan 6-1-5 may be applied by either ground equipment or air-craft. For ground equipment dilute in sufficient water to give adequate coverage. For aerial applications, dilute in 3 to 4 gallons of water per acre.

**Early-Season Program:** For general early season control, including plant bugs and fleahoppers, use  $1\frac{1}{2}$  to  $2\frac{1}{2}$  pints Torbidan 6-1-5 per acre. Make two or three applications at 7 day intervals starting at the two leaf stage. For thrips control only use 1 to  $1\frac{1}{2}$  pints per acre, and for cutworms use 2 to  $2\frac{1}{2}$  pints per acre.

**Midseason Insects:** Overwintering boll weevils, plant bugs, aphids, and cotton leafworms are controlled with 2 to  $2\frac{1}{2}$  pints

Caution:

### DIRECTIONS FOR USE ON SOYBEANS

Apply  $2\frac{2}{3}$  pints per acre to control armyworms, cutworms, and green cloverworms.

Apply suitably diluted with water. Do not use in late stages of growth. Diluted use is not recommended for control of cutworms.

**Caution:** Avoid spray drift to other crops. Do not use during growing season. Do not apply less than 4 weeks before harvest. Do not feed treated plants to livestock. Do not feed treated plants to poultry, dairy animals, or swine. Do not use on plants used for human food. Do not feed soybean products to livestock.

### FOR ARMYWORM CONTROL IN SMALL GRAINS

Apply  $2\frac{2}{3}$  pints per acre to control armyworms, cutworms, barley, or rice. Apply suitably diluted with water. With normal application, apply in a volume of 3 to 4 gallons per acre.

Do not apply this product within 15 days of harvest. This product is toxic to shrimp and crab. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

### DIRECTIONS FOR USE ON TOMATOES

For the control of aphids, armyworms, fruitworms, hornworms, leaf miner, loopers, russet mite, and tomato pinworm, apply  $2\frac{2}{3}$  to  $5\frac{1}{3}$  pints of product per acre in 30 to 250 gallons of water by ground application or in 3 to 10 gallons of water by air application. For stink bugs, use  $5\frac{1}{3}$  pints of product per acre. Do not apply within 15 days of harvest.

### FISH, WILDLIFE, AND ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife.

Birds feeding on treated areas may be killed.

Keep out of lakes, streams, or ponds.

**DO NOT** apply where runoff is likely to occur.

**DO NOT** apply when weather conditions favor drift from areas treated.

**DO NOT** contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

### USAGE CAUTIONS

**DO NOT** feed treated forage to dairy animals or animals being finished for slaughter.

**DO NOT** contaminate feed or foodstuffs.

Keep containers closed when not in use.

**DO NOT** reuse containers for other purposes.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**