

KROKE

RESIDUAL INSECT SPRAY

CONTAINS BAYGON

FOR INDUSTRIAL AND INSTITUTIONAL USE

Residual spray is a highly effective, ready to use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and long residual activity.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Millipedes, Paperlice, Sand Spiders.

USE ONLY OUTDOORS AGAINST: Fleas, Flies, Gnats, Mosquitos, Punkies.

Exposed stages of Cadelles, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Floor Moths, Red Flour Beetles, Sawtoothed Grain Beetles and Weevils.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Cautions

ACTIVE INGREDIENTS:	
o-Isopropoxyphenyl methylcarbamate	1.00%
Petroleum distillate	84.00%
INERT INGREDIENTS:	
	15.00%
	100.00%

* U.S. Patent No. 3,111,539
This material used under license from Chemagro Corporation.

† Baygon is the registered trademark of Farbenfabriken Bayer A. G.
Chemagro Corporation licensee
USDA Reg. No. 980097

Net contents, 1 U.S. Gallon

DISTRIBUTED BY

CHEMICAL SERVICE

DIVISION OF IMOCO GATEWAY CORPORATION
BALTIMORE MARYLAND 21230

DIRECTIONS FOR USE

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent surfaces where insects may crawl when they come out of hiding. Repeat treatment when necessary.

COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers, and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

ANTS: Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES AND WEEVILS: (Exposed stages) Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

SILVERFISH, FIREBRATS AND PAPER-LICE: Apply to localized areas infested by these insects.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS AND SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For clover mites give special attention to window sills.

FLEAS, FLIES, GNATS, MOSQUITOS, PUNKIES AND SANDFLIES: (outdoor only) Spray infested areas thoroughly including outside surfaces of screen doors, window frames, foundation, patios and other areas where insects may enter.

ACCEPTED

Mar 12, 1970

FOR USE IN MEAT PACKING AND FOOD PROCESSING PLANTS: Do not use in edible products areas of food processing plants. When using the spray in the inedible products areas where no exposed food or processing equipment is present, take any needed precautions to avoid contamination of packaged or stored food commodities. Any exposed food products, equipment, or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. The spraying of unloading docks and outside areas where insects congregate will aid in preventing re-infestations. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

INSECT CONTROL IN RESEARCH FACILITIES: The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated.

Do not allow children to contact treated areas until surfaces are thoroughly dry.

Do not store near food or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces.

Do not apply to animals. Remove pets and cover fish bowl before spraying. Do not spray plants.

COMBUSTIBLE: Do not spray into or near open flame. Do not smoke while using. Keep container closed and away from stoves, radiators, or other hot surfaces.

Do not reuse empty container. Perforate container and bury or discard in a safe place.

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Should illness occur, get prompt medical aid.

Antidote - Atropine Sulfate

railways
BEETLES,
IN MEAL
FLOUR
ES, SAW-
ND WEE-
ghly treat
sted area
ices, cor-
tion and
portance.
products.
PAPER-
infested
sleeping
seboards,
ll cracks,
repeat as
be placed
ying. Do
ARWIGS,
WBUGS,
oroughly
perimeter
or clover
o window
SQUITOS,
(outdoor
oroughly
screed
ndatio
e inse

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Millipedes, Paperlice, Sand Spiders.

USE ONLY OUTDOORS AGAINST: Fleas, Flies, Gnats, Mosquitos, Punkies.

Exposed stages of Cadelles, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Floor Moths, Red Flour Beetles, Sawtoothed Grain Beetles and Weevils.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Cautions

ACTIVE INGREDIENTS:

o-Isopropoxyphenyl methylcarbamate**	1.00%
Petroleum distillate	84.00%

INERT INGREDIENTS:

15.00%
<u>100.00%</u>

**U.S. Patent No. 3,111,539
This material used under license from Chemagro Corporation.

*Baygon is the registered trademark of Farbenfabriken Bayer A. G.
Chemagro Corporation licensee.
USDA Reg. No. 900-93

Net contents: 1 U.S. Gallon

DISTRIBUTED BY

CHEMICAL SERVICE

DIVISION OF IMOCO-GATEWAY CORPORATION
BALTIMORE, MARYLAND 21230

