

DANGER

POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Rapidly absorbed through skin. Wear natural rubber gloves, protective clothing, and goggles.

Do not get in eyes, on skin, or on clothing. In case of contact remove clothing; wash thoroughly with soap and water. If eyes flash with plenty of water and get prompt medical attention.

Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

DO NOT breathe spray mist or vapors. Wear a mask or respirator of a type passed by U. S. Department of Agriculture for methyl parathion protection.

KEEP all unprotected persons and animals out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be reentered until drifting spray mist and volatile residues have dissipated. If headache, dizziness, or nausea develops during or after use, get out of contaminated area at once. Get medical attention.



ANTIDOTE—IF SWALLOWED: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately. **IF ON SKIN:** In case of contact remove contaminated clothing and immediately wash skin with soap and water.

PHYSICIAN'S NOTE: Methyl parathion exerts muscarinic-like effects—i. e., stimulation of the parasympathetic nervous system. Repeated use of a parasympathetic depressant, such as atropine, to the limit of patient's tolerance is indicated to relieve symptoms. 2-PAM is also an antidote and may be administered in conjunction with atropine. The administration of oxygen is helpful. Never give morphine to patient suspected of parathion poisoning. Toxaphene produces central nervous system excitation that ordinarily responds to the barbiturate and derivatives, but these drugs should be considered only if complete atropinization fails to relieve symptoms. Refer to "Clinical Handbook on Economic Poisons," prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

GENERAL DIRECTIONS

Prepare a spray mixture of Torbidan 18 as follows:

1. Add about 1/2 all the total amount of water to be used to the spray tank.
2. Start agitator or pump before adding the amount of Torbidan 18.
3. Continue mixing until thoroughly mixed.

DIRECTIONS FOR USE ON COTTON

Torbidan 18 may be applied by either manual equipment or aircraft. For ground equipment dilute in sufficient water to give adequate coverage. For aerial application, dilute in 3 to 4 gallons of water per acre.

Hercules warrants that this product conforms to the chemical description on the label. When used in accordance with the directions under normal conditions, this product is reasonably fit for its intended purpose. Factors such as method of application, weather, soil, and soil conditions, mixture with other chemicals, and other factors affect the use of this product and the user assumes all risks of any such use.

Early-Season Program: For general early season control, including plant bugs and fleahoppers, use 1.5 pints Torbidan 18 per acre. Make two or three applications at 7-day intervals, starting at the two leaf stage. For thrips control only, use 0.75 to 1 pint per acre and for cutworms use 1.5 to 2 pints per acre.

Midseason Insects: This will control both weevils, plant bugs, and cutworms. Dilute with water to give 3 pints per acre. For cutworm control, dilute with 1.5 to 2 pints per acre. Make two or three applications at 7-day intervals if conditions indicate they are necessary.

Late-Season Control: Use 3 to 5 pints per acre to control plant weevils, bollworms, and aphids. Use the higher rate for heavy bollworm outbreaks. Repeat applications at 5 to 7 day intervals as insect counts indicate they are necessary. Do not use more than 4 pints per acre after bolls open.

Caution:

Do not apply within 7 days of hand picking or harvest. (Workers entering fields within 24 hours after application should wear protective clothing.) Do not graze dairy animals or animals being finished for slaughter in treated fields.

DIRECTIONS FOR USE ON SOYBEANS

Apply 2 to 3 pints of Torbidan 18 per acre to control aphids, armyworms, bean leaf beetles, corn earworms, grasshoppers, green cloverworms, stink bugs, and velvetbean caterpillars. Apply suitably diluted with water. For aerial applications, apply diluted insecticide at 3 to 4 gallons per acre.

Caution: Avoid spray drift. Do not apply less than 3 weeks before harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean trash to livestock or poultry.

FOR ARMYWORM CONTROL IN SMALL GRAINS

Apply 2 to 3 pints per acre to control armyworms in wheat, oats, barley, and rye. Apply suitably diluted with water. Aerial applications apply a volume of 3 to 4 gallons per acre. Do not apply within 15 days of harvest.

USAGE CAUTIONS

- DO NOT feed treated forage to dairy animals or animals being finished for slaughter.
- DO NOT contaminate feed or foodstuffs.
- Keep containers closed when not in use.
- DO NOT reuse containers for other purposes.
- This product is highly toxic to bees and other beneficial insects. Do not apply to blooming alfalfa. Protective information may be obtained from the Cooperative Agricultural Entomology Service.
- DESTROY all containers when empty, including the label. Then and always with waste. Do not burn.
- DO NOT use in the following states: ALABAMA, ARIZONA, CALIFORNIA, FLORIDA, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MISSISSIPPI, MISSOURI, NEBRASKA, NEVADA, NEW YORK, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, SOUTH CAROLINA, TEXAS, VIRGINIA, WISCONSIN, and WYOMING.

ENVIRONMENTAL CAUTIONS

- Do not apply to water bodies or to the water table. Do not apply to water bodies or to the water table. Do not apply to water bodies or to the water table. Do not apply to water bodies or to the water table. Do not apply to water bodies or to the water table.
- Do not apply where birds are likely to roost.
- Do not apply where wild animals are likely to roost.
- Do not apply to water bodies or to the water table.
- Do not apply to water bodies or to the water table.

NONWARRANTY

Outdoor Use:

ANTS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, FLIES, HORNETS, MILLIPEDES, MOSQUITOES, SOWBUGS, WASPS, AND BROWN DOG TICKS: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios, or other places where insects may enter. For ants, spray hills and runways. For control of wasps and hornets apply after dark when all of the insects have returned to the nest. Repeat as necessary. Do not spray vegetation.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of area being treated. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly after using. Avoid contamination of food, utensils, and food preparation areas. Should illness occur, get prompt medical aid. Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

TO PHYSICIAN—Atropine sulfate is antidotal.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

WARNING:

FLAMMABLE: Contents under pressure: Do not spray into or near open flame. Do not smoke while using. Keep container away from stoves, radiators, or other hot surfaces. Do not place in direct sunlight. Temperatures exceeding 120 degrees F may cause the container to burst. Do not puncture or throw container into fire.

Do not apply to rugs, carpets, draperies, wallpaper, or similar materials that may be stained. Avoid extensive wetting of plastic, rubber, and asphalt surfaces such as tiles and floor coverings.

KROKE

**Residual Insecticide
Contains Baygon®**

Active Ingredients:

o-Isopropoxyphenyl methylcarbamate*	0.5%
Petroleum distillates	69.0%
Total Actives	69.5%
Inert Ingredients:	30.5%

*U.S. Patent No. 3,111,539
Canadian Patent No. 725,204

Baygon® is a registered trademark of Farben Fabriken Bayer AG, Chemagro Corp., licensee

KILLS ANTS, ROACHES, SPIDERS AND CERTAIN OTHER CRAWLING INSECTS

INSECT-KILLING POWER LASTS FOR DAYS

SPRAY ONLY ON SURFACES DO NOT USE AS A SPACE SPRAY

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See other precautions on side panel


Net Weight 1 lb. 1.7 oz. avcir.

EPA Reg. No. 900-99

**DISTRIBUTED BY
CHEMICAL SERVICE**

DIVISION OF MACCO-GATEWAY CORPORATION

BALTIMORE, MARYLAND 21230



DIRECTIONS FOR USE

KROKE RESIDUAL INSECTICIDE is a highly effective formulation for use in industrial, commercial and institutional areas. It provides rapid kill of crawling insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides:

INDOORS

Ants	Saw-toothed grain beetle (exposed stage)
Brown dog ticks	Scorpions
Crickets	Silverfish
Roaches	Spiders

OUTDOORS

Clover mites	Hornets
Earwigs	Millipedes
Fleas	Mosquitoes
Flies	Sowbugs
	Wasps

In addition to rapid initial kill, KROKE RESIDUAL INSECTICIDE stays on treated surfaces and kills insects many days after the spray deposit has dried. Spray infested surfaces until wet, moving the spray stream rapidly to prevent excessive wetting of surfaces, and waste of material. Point spray away from people, plants, and pets.

Indoor Use:

ANTS, COCKROACHES, CRICKETS, SILVERFISH, SPIDERS, SCORPIONS, SAW-TOOTHED GRAIN BEETLES (exposed stage), **WATERBUGS:** Spray into cracks, crevices, around baseboards, and other places where these insects hide. Apply behind and beneath cabinets, refrigerators, sinks, stoves, and in and around waste containers. For ants, spray trails and places where ants enter premises. Repeat as needed. Do not spray food-stuffs or house plants. Remove pets and cover fish bowls before spraying. Do not allow children to contact surface until surfaces are completely dry.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

LEB

Covers 5,000 or 10,000 Sq. Ft.

NET WT. 20 LBS.



Net Wt. 20 Lbs.
Covers 5,000 or 10,000 Sq. Ft.

LEBANON INSECT and GRUB CONTROL WITH DURSBAN*

SPREADER SETTINGS

	5,000	10,000
Spreader Rate (lb/1000)	2	4
Spreader Setting	1	2
Spreader Speed (mph)	1.5	3.0
Spreader Width (ft)	10	20

These settings are approximate. Always check your own spreader with proportionate amount on 100 square foot area first.

HOW TO USE

5,000 sq. ft. for soil grubs
For Japanese Beetle Larvae, and European Chafer Larvae, apply Lebanon Insect and Grub Control at the rate of 4 lbs. per 1000 sq. ft. immediately after application, water the turf to wash granules into the soil area of infestation.

10,000 sq. ft. for soil surface insects
For sod webworms, cutworms, chinch bugs, ants, mole crickets, brown dog ticks and earwigs, apply Lebanon Insect and Grub Control at the rate of 2 lbs. per 1000 sq. ft. Treat when insects first appear. Immediately after application, sprinkle the turf with water to wash granules into the soil area of infestation. Repeat as necessary.

Use a fertilizer spreader to make an application. Follow the suggested spreader settings.
Keep children and pets off treated areas until the material is worked into the soil and grass is dry.
Do not mow for 24 hours following treatment.

CAUTION

Keep out of reach of children. Keep off equipment.
Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water, and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse.
Keep away from food, feedstuffs and domestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not re-use empty container. Dispose of empty container and wastes by burning or by burying. If burned, keep out of smoke and fumes.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

ACCENTED
JAN 23 1976

5,000 SQ. FT. COVERAGE Kills Soil Feeding Insects: Japanese Beetle Larvae, European Chafer Larvae
10,000 SQ. FT. COVERAGE Kills Soil Surface Insects: Sod Webworms, Cutworms, Chinch Bugs, Ants, Mole Crickets, Brown Dog Ticks, Earwigs

CAUTION: KEEP OUT OF REACH OF CHILDREN. See additional caution on back.

INSECTICIDE INGREDIENT STATEMENT

ACTIVE INGREDIENT	PERCENTAGE
Chlorpyrifos (0,0 diethyl 0,3,5,6-tetrachloro-2-pyridyl) phosphorothioate	1.14%
INERT INGREDIENTS	98.86%
TOTAL	100.0%

EPA REG. NO. 961-326 U.S. Patent No. 3,444,566

LEBANON