


Formula 93

Hospital Disinfectant Cleaner



• GERMICIDAL • STAPHYLOCIDAL • TUBERCULOCIDAL • SALMONELLACIDAL • PSEUDOMONACIDAL • FUNGICIDAL • VIRUCIDAL*

Effectively used by hospitals, schools, nursing homes, cafeterias, restaurants, industries, colleges, universities, medical clinics, public buildings and institutions.

Phenol Coefficient:
8 A. O. A. C. (*S. typhosa*)
10 A. O. A. C. (*S. aureus*)

ACTIVE INGREDIENTS23.25%
Potassium Laurate and
Myristate Soaps10.00%
Potassium
o-benzyl-p-chlorophenate 5.28%
Isopropanol 5.00%
Potassium
ortho-phenylphenate 1.83%
Tetrasodium
ethylenediamine
tetraacetate 1.14%

INERT INGREDIENTS:76.75%
(includes 0.25% sodium nitrite)

E.P.A. Reg. No. 900-105
E.P.A. Est. No. 900-CA-1, IL-1, MD-1

NET U.S. GALS.

Use dilutions range from a maximum of 2½ oz. down to as little as 1¼ oz. for a broad range of effective activity. This is a powerful germicidal cleaner-detergent suitable for use on walls, floors, and similar hard, non-porous surfaces in a hospital or institution. Highly concentrated, it swiftly destroys a broad range of micro-organisms including herpes simplex virus. It provides environmental sanitation by its thorough action while removing dirt and deodorizing the soiled area during the cleaning operation. It is an important contribution to the housekeeping procedure and aids in the reduction of nosocomial acquired infection from inanimate surfaces. Regular damp-mopping and regular scrubbing of floors and sponge-cleaning of walls is recommended to provide conditions that minimize potential cross infection.

DANGER: **KEEP OUT OF REACH OF CHILDREN.**

Do not use, pour, spill or store near heat or open flame.

Causes eye damage and skin irritation. Do not get in eyes, or on skin or clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food. Rinse empty container thoroughly with water and discard it.

FIRST AID — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Application of this material should be restricted to trained personnel. The exercise of basic industrial safety precautions and the wearing of industrial safety garments is recommended where appropriate.

D I R E C T I O N S **DISINFECTING — CLEANING —** **DEODORIZING.**

For **disinfecting** while cleaning, keep surfaces wetted with solution as recommended for at least 10 minutes. For **general cleaning and disinfecting**, use 1¼ ounces per gallon of water (100 to 1 dilution). Remove gross filth and heavy soil deposits from surface prior to application of the recommended solution. In hospitals and where **Pseudomonas** organisms are suspected, use 2½ ounces per gallon of water. **Tuberculocidal** against *Mycobacterium Tuberculosis* Var. Bovis at a Dilution of 1¼ ounces per gallon of water. *Virucidal against herpes simplex virus on environmental surfaces at a dilution of 2½ ounces per gallon of water. Effective against pathogenic fungi at a dilution of 2½ ounces per gallon of water.

When increased detergency is required in heavily soiled areas, the amount may be doubled or tripled. When used as directed it is harmless to flooring, tile, terrazzo, painted or other finished surfaces.



CHEMICAL SERVICE DIV. OF AMERICAN CHEMICAL CORPORATION
HOWARD & WENTZ STREETS / BALTIMORE, MARYLAND 21230 / (301) 752-2610

AD