

sinks, drain boards, stoves, and other areas frequented by insects. Repeat as necessary. Do not contaminate food or utensils. Do not use in edible product areas of food processing plants. Do not use where milk is stored.

ANT MOUNDS—Spray hills thoroughly, using 1½ teaspoon per gallon of water. Mix moist earth in the top of the hill. For other small ants in flower beds and lawns, spray lightly in the infested area. Repeat in 10-15 days if necessary.

MILLIPEDES, SPRINGTAILS & SOWBUGS in and around greenhouses and around homes—Mix 1 teaspoon in 1 gallon of water and apply at the rate of 1 gallon per 150 sq ft (10 ft. x 15 ft.) of soil surface where insects congregate. Repeat in 7-10 days if necessary. Do not apply to soil where food crops are grown.

WASPS & HORNETS—Mix 4 ozs. in 1 gallon of water. Apply to nests in the late evening as a coarse wet spray.


CAUTION—HARMFUL IF INHALED, SWALLOWED OR ABSORBED THROUGH THE SKIN. Malathion is a cholinesterase inhibitor. Note To Physicians: Atropine is antidotal. 2 PAM is also antidotal and may be administered in conjunction with atropine. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Store away from children and domestic animals. Avoid contact with feed and foodstuffs. In case of contact, wash with soap and water; for eyes, get medical attention. **KEEP CONTAINER CLOSED AND AWAY FROM HEAT AND OPEN FLAME.**

This product is toxic to fish. Keep out of any body of water. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Wash thoroughly with water and detergent. Wrap in newspaper and place in trash can for disposal.

Reg. T.M. American Cyanamid Co. E.P.A. Reg. No. 904-43

B. G. PRATT DIVISION

GABRIEL CHEMICALS LTD. PATERSON, N.J. 07509



PRATT

SINCE 1904

**MALATHION-50
SPRAY**

**CONTAINS
CYTHION
PREMIUM GRADE
MALATHION**

ACTIVE INGREDIENTS

Malathion*	By Weight
Xylene	13.3
INERT INGREDIENT	6.7
	Total 100.0

*O,O dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional CAUTIONS

NET CONTENTS, ONE QUART

PRATT

SINCE 1904

**MALATHION-50
SPRAY**

**CONTAINS
CYTHION
PREMIUM GRADE
MALATHION**

ACTIVE INGREDIENTS

Malathion*
Xylene
INERT INGREDIENT

By Weight


13.3
6.7
Total 100.0

*O,O dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional CAUTIONS

NET CONTENTS, ONE QUART



DIRECTIONS

Mix 1½ teaspoonfuls per gallon of water (1½ pints in 100 gallons). Make thorough, full coverage applications for control of aphids, leaf hoppers, thrips, Japanese beetles, Mexican bean beetles and mites, such as European red mite, two spotted mite and clover mite on apples, pears, beans, peas, tomatoes, cabbages and on flowers and ornamentals.

For control of tent caterpillars, canker worms, certain leaf miners and certain scale crawlers on ornamentals use 2 teaspoonfuls per gallon of water (2 pints in 100 gallons). Spray when insects appear and repeat when necessary. Do not apply to food crops later than 7 days before harvest. Injury may occur on certain ferns and some species of Crassula. May injure Bosc pears in Northeast. May injure McIntosh and Cortland varieties of apples if applied within four weeks of harvest.

FLEA CONTROL—To control fleas in kennels, yards, lawns, and under houses, dilute 5 ounces in one gallon of water and apply at the rate of one gallon of diluted spray per 1000 square feet. Clean area before treatment. Fresh bedding should be placed in animals quarters following treatment. Repeat as necessary. **DO NOT APPLY TO DOGS.**

MOSQUITO CONTROL—For the control of adult mosquitoes, flies and small flying insects in outdoor areas, dilute 1 to 28 with water (4½ fl. oz. per gal.) Apply as fine mist or fog. Three fl. oz. of concentrate will treat 1 acre. Repeat application as necessary. May cause spotting on automobile paint finish. Cars should not be sprayed directly. If accidental exposure occurs, the car should be washed immediately.

ROACH CONTROL—For control of roaches, household insects such as roaches, ants, spiders and silverfish and about homes, institutes, and industrial plants, dilute 4 ounces per gallon of water. Apply by means of a coarse spray or paint brush to walls, base boards, under

14-2293-10-73