

FOOD HANDLING ESTABLISHMENT - Places other than private residences in which exposed food is held, processed, prepared or served.

APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY
In areas for reception, storage, packing, canning, bottling, preparation, edible waste storage and enclosed systems, filters, wires, particle filters, syringes. Service areas where food is exposed and facility is in operation also considered a food area.

Apply small amounts directly into cracks and crevices using applicator. Avoid delivering a fine stream in joints between elements of construction, between equipment and floors, between piping to voids and hollow spaces in walls, equipment housings, conduits, motor housings, sections of switch boxes, where coaches, ants, and interbugs live.

Apply to application de-energize electrical equipment. After treatment, allow time for oil to evaporate before starting equipment placing switches to "on" position.

APPLICATION OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

Apply in facilities where prepared foods are served, such as restaurants, but excluding areas where foods may be prepared.

Apply only treatment to selective surfaces such as baseboards, elements of construction into cracks and crevices. Avoid treating surfaces likely to be contacted by food. (Do not apply if facility is in operation or foods are exposed.)

Apply to areas - includes garbage rooms, lavatories, floor drains, restrooms, lockers and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after cleaning or bottling).

Apply to baseboard areas, around water pipes, surfaces behind equipment, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

100-1000
100-1000
100-1000

PRATT'S VAPONA[®] FOGGING SPRAY

AN OIL BASE SPRAY READY-TO-USE IN FOG OR MIST GENERATORS

For use in control of flies, mosquitoes, wasps, gnats, fruit flies and other small flying insects in dairy and horse barns, restaurants and food processing plants. Also for direct application to Dairy and Beef Cattle.

ACTIVE INGREDIENTS

2,2-Dichlorovinyl Dimethyl Phosphate	0.92%
Related Compounds	0.08
Petroleum Hydrocarbons	99.00
	<hr/>
	100.00%

®Shell Trademark for DDVP

WARNING: May be fatal if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling, and before eating or smoking. Do not contaminate feed, water, or food-stuffs. Keep out of reach of children. If swallowed induce vomiting immediately and get medical attention. Atropine is antidotal.

DECONTAMINATE AND DESTROY USED CONTAINERS. Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

WARNING

KEEP OUT OF REACH OF CHILDREN

U.S.D.A. REG. #904-102

**B. G. PRATT COMPANY
PATERSON, NEW JERSEY**

DIRECTIONS

DAIRY & BEEF CATTLE SPRAY: For use in the control of certain insects attacking livestock (houseflies, horn-flies, stableflies, mosquitoes). Apply 1 to 2 fluid ounces per animal as a mist spray daily with hand or automatic sprayer. Apply spray so as to wet the hair. Do not apply in excess of 2 fluid ounces per animal.

FOGGING SPRAY FOR ANIMAL BUILDINGS: For use in control of flies, gnats and mosquitoes in Dairy and Horse Barns, Shelter Sheds, Milk Sheds and Outdoor Areas apply 1 pint of solution as a fog to each 8,000 cubic feet of enclosure. Before applying reduce air movement as much as possible by closing doors, windows and other openings.

DO NOT USE THIS SOLUTION FOR FOGGING PURPOSES IN AREAS WHERE ANIMALS HAVE RECEIVED DIRECT APPLICATION WITHIN 8 HOURS.

INDUSTRIAL SPRAY: For FLIES, MOSQUITOES, WASPS, GNATS, FRUIT FLIES and OTHER SMALL FLYING INSECTS for application only by pest control operators in ~~RESTAURANTS, BAKERIES and FOOD PLANTS processing vegetables, fruits, nuts, meat, poultry, fish and dairy products~~ apply as a space spray with a fog or mist generator at the rate of 1 gallon to 64000 cu. ft. Apply when plants are not in operation. Occupants should vacate before indoor treatment and should not reoccupy the premises until it has been ventilated. After application open several windows and doors to create air current to insure reduction of the VAPONA insecticide vapors. Food should be removed and food handling equipment covered with wrapping paper during application or washed after application is completed. During fogging or mist spraying indoors use a mask or respirator of the type passed by the U. S. Department of Agriculture for DDVP protection.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.