




FRUIT TREE SPRAY

Contains MALATHION, METHOXYCHLOR, CAPTAN
INSECT & DISEASE CONTROL ON
APPLES, PEARS, PEACHES, PLUMS



ACTIVE INGREDIENTS:	BY WEIGHT
Malathion*	6.00%
Methoxychlor Technical**	10.00%
Captan***	5.67%
Related Derivatives	0.33%
Sulfur****	25.00%
INERT INGREDIENTS:	53.00%
TOTAL	100.00%

*O,O dimethyl dithiophosphate of diethyl mercaptosuccinate.

**Equivalent to 8.8% 2,2 bis (p-methoxyphenyl) 1, 1, 1-trichloroethane and 1.2% of other isomers and related compounds.

***N-(trichloromethyl)-4-cyano-1, 2-dichlorobenzene.

****Not less than 88% passing 325 mesh.

EPA Reg. No. 904-128

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado completamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Possible mucosal damage may contraindicate the use of gastric lavage.

IF INHALED: Remove victim to fresh air.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Launder clothing before reusing.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

NET WEIGHT: ONE POUND

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER—Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Keep children and pets out of treated areas until spray has dried or dust has settled. Do not contaminate food or feed. Wash hands thoroughly after handling. Food utensils, such as tablespoons, should not be used for food after use with pesticides.

PRECAUTIONS—HOMEOWNER USES—Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal Law. The following endangered species and the general location of their habitats have been identified by county:

Blackwater Darter	Alabama	(Lauderdale, Limestone and Madison)
	Tennessee	(Lawrence and Wayne)
Pecos Gambusia	New Mexico	(Chaves and Eddy)
	Texas	(Reeves, Jeff Davis and Pecos)
Freshwater Mussels	Tennessee	(Hancock Claiborne, Hawkins and Sullivan)
	Virginia	(Smyth, Scott, Lee, Washington and Grayson)
Comanche Springs Pupfish	Texas	(Reeves and Jeff Davis)

Prior to use of this product in the above counties, you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

PHYSICAL OR CHEMICAL HAZARDS

Avoid contact with open flame

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in tightly closed original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

LIST OF INSECTS

ON CHERRIES:

Aphid
Thrips

ON PLUMS:

Curculio
Plant Bugs

ON STRAWBERRIES:

Spittlebug
Strawberry Weevil

ON APPLES, PEARS, PEACHES.

Aphids
Cankerworm
Codling Moth
Red Spider Mite
Leaf Hopper
Leaf Roller
Oriental Fruit Moth
Tent Caterpillar

LIST OF DISEASES

SCAB AND POWDERY MILDEW

On Apples and Pears

BROWN ROT

On Cherries, Peaches, Plums

BOTRYTIS ROT

On Strawberries

DO NOT ALLOW PERSONS TO ENTER TREATED AREAS WITHIN 4 DAYS FOLLOWING APPLICATION UNLESS PROTECTIVE CLOTHING IS WORN. CONSPICUOUSLY POST RE-ENTRY INFORMATION AT SITE OF APPLICATION.

***DO NOT ROTATE A FOLIAR TREATED CROP WITH CROPS OTHER THAN THOSE WITH REGISTERED CAPTAN USES.**

Mix 5 tablespoonsful per gallon, 1 cupful for 3 gallons or 1 pound to 12 gallons of water. For hose-end sprayer, follow manufacturer's directions. Mix thoroughly before using and occasionally during uses to assure uniformity of spray mix. Do not store diluted spray.

For best results follow a regular spray schedule, timing is important for both disease and insect control. Spray thoroughly to cover all fruit and leaf surfaces. A 6 to 8 foot tree requires about 3 gallons of spray; larger trees need 6 or more gallons.

APPLES AND PEARS: Spray when new growth appears and repeat at weekly intervals up to bloom. Then spray after petals fall at 10 day intervals. Do not spray within 7 days of harvest.

PEACHES: Make about seven spray applications at 10 day intervals, starting in the pink stage. Do not apply within 21 days of harvest. For Peach Tree Borer, spray or paint trunk from crotches to ground at monthly intervals, starting about July 1st, using ½ pound per gallon of water. Do not apply this concentration to fruit.

PLUMS: Make first application just before blooms open. After petals fall make about seven sprayings at 10 day intervals. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7 to 14 day intervals—not later than 14 days of harvest.

CHERRIES: Apply at pre-bloom, shuck fall, two weeks after petal fall, and then 2 weeks before harvest, but not later than 7 days before harvest.

DO NOT USE THIS MATERIAL WITHIN TWO WEEKS AFTER THE APPLICATION OF AN OIL SPRAY. AVOID SPRAYING WHEN TEMPERATURE IS IN EXCESS OF 85°F.


LIMITED WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

This material is packed at full net weight as listed. It is subject to settling and may appear less than full.



FRUIT TREE SPRAY

Contains MALATHION, METHOXYCHLOR, CAPTAN
INSECT & DISEASE CONTROL ON
APPLES, PEARS, PEACHES, PLUMS



ONLY FOR SALE TO,
STORAGE AND USE BY SERVICE PERSON OR FOR AGRICULTURAL USES.

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion*	6.0 %
Methoxychlor Technical**	10.0 %
Captan***	5.67%
Related Derivatives	0.33%
Sulfur****	25.0 %
INERT INGREDIENTS:	53.0 %
Total	100.00%

*0,0 dimethyl dithiophosphate of diethyl mercaptosuccinate.

**Equivalent to 8.8% 2,2 bis (p-methoxyphenyl) 1, 1, 1-trichloroethane and 1.2% of other isomers and related compounds.

***N-(trichloromethylthio)-4-cyclohexene-1, 2 dicarboximide.

****Not less than 98% passing 325 mesh.

EPA Reg. No. 904-128

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

PRECAUCION AL USUARIO: SI usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

IF INHALED: Remove victim to fresh air.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Launder clothing before reusing.

Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331
NET WEIGHT: 25 POUNDS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER—Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Keep children and pets out of treated areas until spray has dried or dust has settled. Do not contaminate food or feed. Wash hands thoroughly after handling. Food utensils, such as tablespoons, should not be used for food after use with pesticides.

PRECAUTIONS—AGRICULTURAL USES—Mixers, loaders, and applicators, when mixing, loading, and applying, must wear mid-forearm to elbow-length natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan; boots or overshoes; one-piece overalls which have long sleeves and long pants; face shield or goggles and a hat or other appropriate head covering.

Fieldworkers and harvesters must wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan residues. Leather or fabric gloves are not acceptable.

A dust mask and a chemical-resistant apron must be worn when mixing and loading this product.

Clothing worn while loading, mixing, and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

Applicator's protection may also be obtained by use of an enclosed tractor cab with a properly filtered air supply.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal Law. The following endangered species and the general location of their habitats have been identified by county:

Blackwater Cooter	Alabama	(Lauderdale, Limestone and Madison)
	Tennessee	(Lawrence and Wayne)
Peach Gambusia	New Mexico	(Chaves and Eddy)
	Texas	(Pecos, Jeff Davis and Pecos)
Freshwater Mussels	Tennessee	(Marion, Claiborne, Hawkins and Sullivan)
	Virginia	(Bryce, Scott, Lee, Washington and Wayson)
Comanche Springs Pupfish	Texas	(Pecos and Jeff Davis)

Prior to use of this product in the above counties, you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist or State Wildlife Agency) if there is any doubt as to the areas where the outdoor use of captan is prohibited.

PHYSICAL OR CHEMICAL HAZARDS

Avoid contact with open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in tightly closed, original container in a locked area away from children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIST OF INSECTS

ON CHERRIES:	ON APPLES, PEARS, PEACHES.
Aphid	Aphids
Thrips	Cankerworm
	Codling Moth
ON PLUMS:	Red Spider Mite
Curculio	Leaf Hopper
Plant Bugs	Leaf Roller
	Oriental Fruit Moth
ON STRAWBERRIES:	Tent Caterpillar
Spittlebug	
Strawberry Weevil	

LIST OF DISEASES

SCAB AND POWDERY MILDEW
On Apples and Pears
BROWN ROT
On Cherries, Peaches, Plums
BOTRYTIS ROT
On Strawberries

RE-ENTRY AND ADDITIONAL APPLICATION RESTRICTIONS FOR AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not allow persons to enter treated areas within 4 days following application unless protective clothing is worn. Conspicuously post re-entry information at site of application.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Sign or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Instruct workers of the area or fields that must be vacated and not re-entered until the sprays have dried. Acquaint workers with the information appearing on this label under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment". When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: 1) The name of this product; 2) Date of application; 3) The statement: "Do not enter without appropriate protective clothing until sprays have dried."; 4) The information given under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment" that is appropriate in case of accidental exposure.

DIRECTIONS FOR USE ON AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS

***DO NOT ROTATE A FOLIAR TREATED CROP WITH CROPS OTHER THAN THOSE WITH REGISTERED CAPTAN USES.**

Mix 5 tablespoonsful per gallon, 1 cupful for 3 gallons or 1 pound to 12 gallons of water. For hose-end sprayer, follow manufacturer's directions. Mix thoroughly before using, and occasionally during uses to assure uniformity of spray mix. Do not store diluted spray.

For best results follow a regular spray schedule, timing is important for both disease and insect control. Spray thoroughly to cover all fruit and leaf surfaces. A 6 to 8 foot tree requires about 3 gallons of spray; larger trees need 6 or more gallons.

APPLES AND PEARS: Spray when new growth appears and repeat at weekly intervals up to bloom. Then spray after petals fall at 10 day intervals. Do not spray within 7 days of harvest.

PEACHES: Make about seven spray applications at 10 day intervals, starting in the pink stage. Do not apply within 21 days of harvest. For Peach Tree Borer, spray on main trunk from crotch to ground at monthly intervals, starting about July 1st, using 1/4 pound per gallon of water. Do not apply this concentration to fruit.

PLUMS: Make first application just before blooms open. After petals fall, make about seven sprayings at 10 day intervals. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7 to 14 day intervals—not later than 14 days of harvest.

CHERRIES: Apply at pre-bloom, shuck fall, two weeks after petals fall, and then 2 weeks before harvest, but not later than 7 days before harvest.

DO NOT USE THIS MATERIAL WITHIN TWO WEEKS AFTER THE APPLICATION OF AN OIL SPRAY. AVOID SPRAYING WHEN TEMPERATURE IS IN EXCESS OF 85°F

LIMITED WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

This material is packed at full net weight as listed. It is subject to settling and may appear less than full.

50M MP

Prod. No. 01520