



FRUIT TREE SPRAY

Contains MALATHION, METHOXYCHLOR, CAPTAN
INSECT & DISEASE CONTROL ON
APPLES, PEARS, PEACHES, PLUMS

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion*	6.00%
Methoxychlor Technical**	10.00%
Captan***	5.87%
Related Derivatives	0.13%
Sulfur****	25.00%
INERT INGREDIENTS:	53.00%
TOTAL	100.00%

*O,O Dimethyl dithiophosphate of diethyl mercaptosuccinate.

**Equivalent to 8.8% 2,2 bis (p-methoxyphenyl) 1, 1, 1-trichloroethane and 1.2% of other isomers and related compounds.

***N-(trichloromethylthio)-4-cyclohexene-1, 2 dicarboximide.

****Not less than 98% passing 325 mesh.

EPA Reg. No. 904-128

**KEEP OUT OF REACH OF CHILDREN
DANGER**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Launder clothing before reusing.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For
Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

NET WEIGHT: ONE POUND

904-128

PM 12-

log 4

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER—Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Keep children and pets out of treated areas until spray has dried or dust has settled. Do not contaminate food or feed. Wash hands thoroughly after handling. Food utensils, such as tablespoons, should not be used for food after use of this pesticide.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Avoid contact with open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in tightly closed original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

LIST OF INSECTS

ON CHERRIES:
Aphid
Thrips

ON PLUMS:

Curculio
Plant Bugs

ON STRAWBERRIES:

Spittlebug
Strawberry Weevil

**ON APPLES, PEARS,
PEACHES.**

Aphids
Cankerworm
Codling Moth
Red Spider Mite
Leaf Hopper
Leaf Roller
Oriental Fruit Moth
Tent Caterpillar

LIST OF DISEASES

SCAB AND POWDERY MILDEW

On Apples and Pears

BROWN ROT

On Cherries, Peaches, Plums

BOTRYTIS ROT

On Strawberries

Mix 5 tablespoonsful per gallon, 1 cupful for 3 gallons or 1 pound to 12 gallons of water. For hose-end sprayer, follow manufacturer's directions. Mix thoroughly before using and occasionally during uses to assure uniformity of spray mix. Do not store diluted spray.

For best results follow a regular spray schedule, timing is important for both disease and insect control. Spray thoroughly to cover all fruit and leaf surfaces. A 8 to 8 foot tree requires about 3 gallons of spray; larger trees need 6 or more gallons.

APPLES AND PEARS: Spray when new growth appears and repeat at weekly intervals up to bloom. Then spray after petals fall at 10 day intervals. Do not spray within 7 days of harvest.

PEACHES: Make about seven spray applications at 10 day intervals, starting in the pink stage. Do not spray within 21 days of harvest. For Peach Tree Borer, spray or paint trunk from crotch to ground at monthly intervals, starting about July 1st, using 1/2 pound per gallon of water. Do not apply this concentration to fruit.

PLUMS: Make first application just before blooms open. After petals fall make about seven sprayings at 10 day intervals. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7 to 14 day intervals—not later than 14 days of harvest.

CHERRIES: Apply at pre-bloom, shuck fall, two weeks after petal fall, and then 2 weeks before harvest, but not later than 7 days before harvest.

DO NOT USE THIS MATERIAL WITHIN TWO WEEKS AFTER THE APPLICATION OF AN OIL SPRAY. AVOID SPRAYING WHEN TEMPERATURE IS IN EXCESS OF 85°F.

LIMITED WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

This material is packed at full net weight as listed. It is subject to settling and may appear less than full.

77M MP

Prod. No. 01518

© Copyright - 1983 - Miller Chemical & Fertilizer Corp.

BEST AVAILABLE COPY

SPRAY

Contains MALATHION, METHOXYCHLOR, CAPTAN
INSECT & DISEASE CONTROL ON
APPLES, PEARS, PEACHES, PLUMS



ACTIVE INGREDIENTS:	BY WEIGHT
Malathion ¹	8.0 %
Methoxychlor Technical ²	10.0 %
Captan ³	8.5 %
Residual Disinfectant	25.0 %
Sulfur ⁴	33.0 %
INERT INGREDIENTS:	33.0 %
Total	100.0%

¹0.5 dimethyl disulphosphorothioic acid dimethyl mercaptosuccinate.
²Equivalent to 8.5% 2,2 bis (p-methoxyph.), 1,1-trichloroethane and 1.2% of other isomers and related compounds.
³N-(2-chlorophenyl)-4-cyano-2-methyl-5, 2 dicarboximide.
⁴Not less than 90% passing 32.5 mesh.
 EPA Reg. No. 904-130

EPA Est. No. 440-1

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido traducida completamente.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For
Fruit Grafting Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331
NET WEIGHT: 25 POUNDS

BEST AVAILABLE COPY

3064
D.P.A.

8.28.87

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Launder clothing before reusing.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER—Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Keep children and pets out of treated areas until spray has dried or dust has settled. Do not contaminate food or feed. Wash hands thoroughly after handling. Food utensils, such as tablespoons, should not be used for food after use with pesticides.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Avoid contact with open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in the tightly closed, original container in a locked area away from children and pets. Do not reuse empty container. For disposal, wrap tightly closed original container in several layers of newspaper and discard in trash or at an approved waste disposal facility.

BEST AVAILABLE COPY

DAIRY

000000

LIST OF INSECTS

- | | |
|--|---|
| ON CHERRIES:
Aphid
Thrips

ON PLUMS:
Curculio
Plant Bugs

ON STRAWBERRIES:
Spittlebug
Strawberry Weevil | ON APPLES, PEARS, PEACHES:
Aphids
Cankerworm
Codling Moth
Red Spider Mite
Leaf Hopper
Leaf Roller
Oriental Fruit Moth
Tent Caterpillar |
|--|---|

- LIST OF DISEASES**
SCAB AND POWDERY MILDEW
On Apples and Pears
BROWN ROT
On Cherries, Peaches, Plums
BOTRYTIS ROT
On Strawberries

Mix 5 tablespoonsful per gallon, 1 cupful for 3 gallons or 1 pound to 12 gallons of water. For hose-end sprayer, follow manufacturer's directions. Mix thoroughly before using and occasionally during uses to assure uniformity of spray mix. Do not store diluted spray.

For best results follow a regular spray schedule, timing is important for both disease and insect control. Spray thoroughly to cover all fruit and leaf surfaces. A 6 to 8 foot tree requires about 3 gallons of spray; larger trees need 6 or more gallons.

APPLES AND PEARS: Spray when new growth appears and repeat at weekly intervals up to bloom. Then spray after petals fall at 10 day intervals. Do not spray within 7 days of harvest.

PEACHES: Make about seven spray applications at 10 day intervals, starting in the pink stage. Do not apply within 21 days of harvest. For Peach Tree Borer, spray or paint trunk from crotches to ground at monthly intervals, starting about July 1st, using 1/2 pound per gallon of water. Do not apply this concentration to fruit.

PLUMS: Make first application just before blooms open. After petals fall, make about seven sprayings at 10 day intervals. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7 to 14 day intervals—not later than 14 days of harvest.

CHERRIES: Apply at pre-bloom, shuck fall, two weeks after petals fall, and then 2 weeks before harvest, but not later than 7 days before harvest.

**DO NOT USE THIS MATERIAL WITHIN TWO WEEKS AFTER THE APPLICATION OF AN OIL SPRAY.
AVOID SPRAYING WHEN TEMPERATURE IS IN EXCESS OF 85°F.**

LIMITED WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

This material is packed at full net weight as listed. It is subject to settling and may appear less than full.

77M .MP

©COPYRIGHT - 1983 - MILLER CHEMICAL & FERTILIZER CORP.

Prod. No. 01526

BEST AVAILABLE COPY