

B.G. PRATT
SINCE 1904
**ROSE &
FLOWER
DUST OR SPRAY**



**KILLS
COMMON INSECTS
CONTROLS
DISEASES
in the
GARDEN**

B.G. PRATT
**ROSE & FLOWER
DUST OR SPRAY**



**CONTROLS DISEASES
KILLS COMMON INSECTS
on
ROSES, FLOWERS,
ORNAMENTALS**

CONTAINS SEVIN*, MALATHION, KELTHANE*

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for complete "CAUTION" statement

NET WEIGHT ONE POUND

B.G. PRATT
**ROSE &
FLOWER
DUST OR SPRAY**



**KILLS
COMMON INSECTS
CONTROLS
DISEASES
in the
GARDEN**

DIRECTIONS

Please Read Entire Label Before Using

Control common insects and diseases in the garden before the insects become numerous or the diseases apparent. Keep all plant surfaces protected throughout the growing season. Apply regularly at 7 to 14 intervals. During humid periods and new growth, apply every 4 to 5 days.

DUSTING: Material is ready for use and can be dusted from the sifter can or transferred to a dust gun. The best time to apply is early morning or late afternoon.

SPRAYING: Prepare a thin, smooth paste by mixing a little water with 8 tablespoonfuls of material. Add paste to water in sprayer to make one gallon. Mix thoroughly. Apply with sprayer designed for garden use. 1 lb. makes 10 gallons of spray. **AGITATE SPRAY WHILE USING.** Do not store mixed spray.

INSECTS CONTROLLED ON ROSES AND FLOWERS
Aphids, Leaf Pollers, Rose Slugs, Rose Chafers, Japanese Beetles, Red Spider Mites, Rose Leafhoppers

INSECTS CONTROLLED ON ORNAMENTAL SHRUBS
Aphids, Bagworms, Japanese Beetles, Boxwood Leaf Miners, Euonymus Scale Crawlers, Lacebugs, Spider Mites

DISEASES CONTROLLED ON FLOWERS
Black Spot and Powdery Mildew on ROSES, Septoria Leaf Spot on CHRYSANTHEMUMS, Didymellina Leaf Spot on IRIS, Rust and Alternaria Leaf Spot on CARNATIONS, Alternaria Leaf Spot, Powdery Mildew, and Anthracnose on MARIGOLDS, ASTERS, PHLOX, and SNAPDRAGONS.

Do not use on certain ferns including Boston, Maidenhair, and Pteris, as well as on some species of Crassula. Do not use on Boston Ivy. Do not use on Japanese Maple.

CAUTION.

HARMFUL IF INHALED OR SWALLOWED.

May cause irritation of nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Wash thoroughly after using. Keep out of reach of children. Store away from domestic animals. Avoid contamination of food and feedstuffs.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACTIVE INGREDIENTS:	By Weight
Carbaryl (1-naphthyl-N-methylcarbamate*)	3.0%
Malathion**	4.0%
Folpet (N-Trichloromethylthiophthalimide)	5.0%
1, 1-bis (Chlorophenyl) 2, 2, 2-trichloroethanol**	1.5%
INERT INGREDIENTS	86.5%
TOTAL	100.0%

*O,O dimethyl dithiophosphate of diethyl mercaptosuccinate
U.S.D.A. Reg. No. 904-130

SEVIN® Union Carbide Corp. *KELTHANE® Rohm & Haas Co

This material is packed at full net weight as listed.
It is subject to settling and may appear less than full.

**PACKAGED BY
B. G. PRATT COMPANY, PATERSON, N. J. 07059**
1969 B. G. P. Co

DIRECTIONS
LIFT HINGED CAP. SQUEEZE TO DUST.

To be most effective, the dust should be applied before the insects become numerous and diseases are apparent. Apply dust evenly to all plant surfaces at 7 to 14 day intervals. During periods of new growth and humid spells, apply at 4 to 7 day intervals.

INSECTS CONTROLLED ON FLOWERS

Aphids, Red Spider Mites, Japanese Beetles, Rose Chafers, Rose Leafhoppers, Leaf Rollers, Rose Slugs on GLADIOLI, DAHLIAS, ASTERS, CHRYSANTHEMUMS, SNAPDRAGONS.

INSECTS CONTROLLED ON ORNAMENTALS

Aphids, Japanese Beetles, Leafhoppers, Spider Mites, European Sawfly, Sawflies, Boxwood Leaf Miners, Bagworms on RHODOLARIX, BURNING BUSH, ALABASTER, JUNIPERS, EUNYMIUS, BOXWOOD and LILAC.

DISEASES CONTROLLED

Rosette, Powdery Mildew, Rust, Septoria Leaf Spot, Anthracnose, CHRYSANTHEMUMS, Dideronia Leaf Spot on IRIS, Rust on Asters, Anthracnose on APARTHEUS, Alternaria Leaf Spot, Powdery Mildew on CHRYSANTHEMUMS, MARIGOLD, ASTERS, SNAPDRAGONS.

Control of the following diseases on the Rose: Mildew, Powdery Mildew, Rust, Black Spot, Leaf Spot, and other diseases. Do not use on Boxwood.

CAUTION

Do not breathe dust. Wash face and hands after use. Do not use in enclosed spaces. Do not use near food. Do not use near children. Do not use near pets. Do not use near electrical equipment. Do not use near open flames. Do not use near heat. Do not use near machinery. Do not use near moving vehicles. Do not use near power lines. Do not use near water. Do not use near swimming pools. Do not use near hot tubs. Do not use near saunas. Do not use near spas. Do not use near hot tubs. Do not use near saunas. Do not use near spas.

NOTICE

Do not use on roses. Do not use on flowers. Do not use on ornamentals. Do not use on lawns. Do not use on trees. Do not use on shrubs. Do not use on vines. Do not use on ground covers. Do not use on perennials. Do not use on annuals. Do not use on vegetables. Do not use on fruits. Do not use on nuts. Do not use on seeds. Do not use on plants. Do not use on animals. Do not use on humans. Do not use on children. Do not use on pregnant women. Do not use on nursing mothers. Do not use on infants. Do not use on the elderly. Do not use on the infirm. Do not use on the disabled. Do not use on the sick. Do not use on the dying. Do not use on the dead.

THIS SQUEEZE DUSTER IS REFILLABLE
REMOVE BOTTOM CLOSURE TO REFILL

This container is not to be refilled. For test dust action, it contains the amount of dust which will fill the container.

Manufactured for
B. G. PRATT COMPANY, PATERSON, N. J. 07649
 © 1969 B. G. P. Co.



KILLS COMMON INSECTS
CONTROLS DISEASES
 on
ROSES, FLOWERS,
ORNAMENTALS

CONTAINS SEVIN®, MALATHION, KELTHANE®
CAUTION: KEEP OUT OF REACH OF CHILDREN
 See back label for complete caution statement

NET WEIGHT 10 OZ AV