

WOODY PLANTS and WEEDS in Industrial Yards, Fence Rows and Right-of-Ways for Poison Ivy, Poison Oak, Honeysuckle, Sumac, Brambles and general noxious weed control use 1/2 cup (4 fl. oz.) per gallon of water. Spray to thoroughly drench all plants. Application may be repeated in a month.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. Wash hands after using. Keep containers closed. Do not store near heat or open flame, nor seeds, fertilizers, insecticides or fungicides. Do not contaminate water used for domestic or irrigation purposes. Do not use same spray equipment for purposes other than application of lawn chemicals and herbicides. Vapors from this product may injure susceptible plants in immediate vicinity. Destroy container when empty. Do not reuse. Keep out of reach of children.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTICE

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

B. G. PRATT CO., PATERSON, N.J. 07509

© 1971 B.G.P. Co.

PRATT IMPROVED
**LAWN WEED
KILLER**

**KILLS CHICKWEED
AND OTHER LAWN WEEDS**

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS

For most effective control apply when weeds are growing vigorously after a rain in the Spring or Fall. Poor results will be attained during periods of drought and turf injury may occur. Do not apply during heat of the day. If rain is expected within 24 hours delay application. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Apply as coarse spray to prevent spray drift to desirable plants.

LAWN WEEDS: (Clover, Chickweed, Chickory, Dandelion, Herbit, Plantain, Spotted Spurge, Wild Onion and certain other weeds) Mix 4 tablespoonfuls (2 fl. oz.) in 3 gallons of water and apply to 600 sq. ft. of lawn. If lawn begins to brown after application water thoroughly. Do not use on Dichondra, Lippia, Clover, St. Augustine or Bent Grass lawns. Wild Onions may require repeated applications in late Fall and early Spring. Do not apply to new lawns until after at least second cutting. Do not reseed for 3 to 4 week period after use of this material.

HARD-TO-KILL WEEDS: (Ragweed, Dock, Thistle, Morning Glory) in Lawns, Fence Rows and Right-of-Ways: Use 3 tablespoonfuls per gallon of water (1 to 100 dilution). Spray to thoroughly wet all plant surfaces. Repeat as necessary.

RECEIVED
JAN 24 1974

B
C | PRATT

LAWN WEED KILLER

Contains 2, 4-D & SILVEX

EPTED

19. 1974

ACTIVE INGREDIENTS:	BY WEIGHT
Equivalent of 2,4-Dichlorophenoxyacetic Acid	18.2%
Equivalent of 2,4-Dichlorophenoxypropionic Acid	7.1
INERT INGREDIENTS	74.7
Total	100.0%

NO. 904-136 SUBJECT
 Equivalent to 10% 2,4D Acid **Equivalent to 5% Silvex
 — Contains the equivalent of 0.8 pounds of 2,4D Acid and 0.6 pounds of Silvex per gallon

USDA Reg. No. 904-136

See other precautions on side panel.

NET CONTENTS: 1 PINT