

WARNING

May be fatal if swallowed.

Can be absorbed through skin.

Do not breathe vapors or spray mist.

Do not get on skin or in eyes or on clothing.

Avoid contamination of feed or foodstuffs.

Wash thoroughly after using.

ANTIDOTE

INTERNAL: Give sodium chloride 15 gm. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear.

EXTERNAL: Wash with soap and water.

CALL A PHYSICIAN IMMEDIATELY

WARNING

Not for use or storage in or around the home. Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. Do not apply to any area not specified on the label.

Do not graze treated areas to dairy animals or animals being finished for slaughter.

Do not store or handle near open flame.

Keep container tightly closed when not in use.

Use only as directed on label.

Do not reuse container. Destroy when empty.

THIS PRODUCT IS NOT EMULSIFIABLE. DO NOT DILUTE WITH WATER.

DIRECTIONS FOR USE:

Bananas:

For control of leaf-eating caterpillars (*Ceratomia* sp.) on bananas, make aerial application of 1 - 3 pints undiluted Toxaphene 50% Concentrate per acre. Repeat application as necessary.

M-111

ACCEPTED
 JAN 10 1970
 FEDERAL BUREAU OF INVESTIGATION
 U.S. DEPARTMENT OF JUSTICE
 COMMENTS
 904-137

tals per gallon of water (4 fl. oz. to 12 gals. or 1 pint to 50 gals.) when insects or damage appear. Repeat at weekly intervals or as necessary for such insects as apple aphid, bagworm, leaf miners, elm leaf beetle, gypsy moth, Japanese beetle, lacebug, rose slug, tent caterpillars, leaf rollers, scale insects and periodical Cicada.

VEGETABLE GARDEN SPRAY: Mix 1 fl. oz. in 3 gallons of water (1 quart in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with directions!

BEANS (green lima and snap) Japanese and Mexican Bean Beetle, Tarnished Plant Bug. **TOMATO, EGGPLANT & PEPPER** Leaf hopper, Lacebug, Tomato fruit worm. **PUMPKIN, SQUASH, CUCUMBER, MELON** Flea beetle, Melonworms, Cucumber beetle, Squash bug. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust or spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Avoid contamination of food and keep out of reach of children. Do not re-use container. Destroy when empty.

NOTICE: Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

EPA REG. NO. 961137
B.C. PRATT DIV., GABRIEL CHEMICALS LTD.
 Paterson, N. J. 07509

PRATT
FLOWABLE SEVIN®
CHINCH BUGS,
JAPANESE BEETLES
 and certain other
LAWN, GARDEN &
ORNAMENTAL INSECTS
 Contains 4 pounds
 of Sevin per gallon

ACTIVE INGREDIENT:	By Weight
Carbaryl (1-naphthyl-N-methylcarbamate)	41.8%
INERT INGREDIENTS	58.2
Total	100.0%

®Trade Mark of Union Carbide Corp. for the active ingredient

See side panel for additional cautions
NET CONTENTS: 1 LIQUID PINT

PROTECT FROM TEMPERATURES BELOW 32°F
 Keep cover on opened container to prevent drying.

DIRECTIONS: Flowable Sevin is an extremely fine dispersion of Sevin in water. **SHAKE WELL BEFORE USING** Pour recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.

PRECAUTIONS: Avoid spray drifting to adjoining food or forage crops or crops in bloom. For protection of honey bees, avoid unnecessary use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. For application to food plants other than those listed, get directions on rates and timing from local agricultural authority. Sevin is compatible with common insecticides, fungicides and miticides, but not with lime or other highly alkaline materials.

LAWN SPRAY: For Ants, Chinch Bugs, Earwigs, Fleas, Fall armyworm, Mosquitoes, Sod webworms (lawn moths) and Millipedes. For best results treat after cutting and when the ground is moist from rain or irrigation using 1/4 pint in 15 to 20 gallons of water to each 500 square feet of established lawn. Apply with hose end or pressure sprayer with ample water to insure turf penetration. Repeat in 2 to 3 weeks if necessary.

ORNAMENTAL PLANT SPRAY: For flowers, Shrubs and Shade Trees including roses, azaleas, mums, lilac, arborvitae, juniper, pine, birch, dogwood, oak and maple (Do not use on Boston Ivy). Apply spray mixture containing 2 teaspoon