


WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Rag Weed, Morning-Glory, Thistles and many other broad-leaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 1.5 fl. oz. in 1 gallon of water or 1 cupful in 5 gallons of water.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable tree or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides or fungicides. Do not use on centipede grass. St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2, 4-D combinations only to turf.


Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

NET CONTENTS: 1 QUART



CAUTION: KEEP OUT OF REACH OF CHILDREN.
See side panel for additional cautions.

B. G. PRATT COMPANY, Paterson, New Jersey



DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Henbit, Wild Onion and many other broadleaf weeds in the Spring . . . FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

SCHEDULE A	SCHEDULE B	WATER GALLONS	AREA SQ. FT.
1 Tablespoonful	3.5 Teaspoonfuls	1	200
1/2 fl. oz.	0.6 fl. oz.		
5 Tablespoonfuls	6 Tablespoonfuls	5	1000
2 1/2 fl. oz.	3 fl. oz.		
1 pint	10 fl. oz.	30	6500

ACTIVE INGREDIENTS:

Dimethylamine Salt of Dicamba (3, 6-dichloro-o-anisic acid)*	3.43%
Dimethylamine Salts of Related Acids	0.55
Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic acid**	17.22
INERT INGREDIENTS	78.80
	100.00%

* Equivalent to 2.84% 3, 6-dichloro-o-anisic acid (dicamba) or 0.26 pounds per gallon.
** Equivalent to 14.27% 2, 4-Dichlorophenoxyacetic acid or 1.25 pounds per gallon.

U.S.D.A. Reg. No. 904-146

10732-2-12-66

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Red Wheel Morning-Glory, Thistles and most other broad leaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 5.0 fl. oz. per gallon of water or 1.0 cupful in 5 gallons of water.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water. For eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray. These sprays are less likely to drift on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation water. Do not pour contents used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra on lawns or turf where desirable flowers or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for application of combinations only to turf.

Contamination of shallow fish ponds or other applicable water reservoirs may result in the death of fish.

NET CONTENTS: 1 QUART



with BANVEL D-3



CAUTION: KEEP OUT OF REACH OF CHILDREN.

See side panel for additional cautions.

B. G. PRATT COMPANY, Paterson, New Jersey

DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Henbit, Wild Onion and many other broadleaf weeds in the Spring. **FOLLOW-DILUTION IN SCHEDULE A.** For late summer and fall applications and for hard-to-kill weeds such as Sheep Squirrel, Oxalis, Yarrow and matric Knotweed, increase the dosages to Schedule B. If weeds persist repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

SCHEDULE A	SCHEDULE B	WATER GALLONS	AREA SQ. FT.
1.0 teaspoonful	3.5 teaspoonfuls	1	200
1.0 fl. oz.	0.6 fl. oz.	5	1000
5.0 teaspoonfuls	3.0 teaspoonfuls	10	1500
1.0 fl. oz.	1.0 fl. oz.	20	3000

ACTIVE INGREDIENTS:
 2-methyl-4-chlorophenoxyacetic acid 3.43%
 2,4-dichlorophenoxyacetic acid 0.55%
 2,4-dichlorophenoxyacetic acid 17.22%
 2,4-dichlorophenoxyacetic acid 76.80%
NET INGREDIENTS 100.00%
 Contains 2.0% 2,4-Dichlorophenoxyacetic acid and 0.5% 2,4-Dichlorophenoxyacetic acid.
 U.S.O.A. Reg. No. 504-146

10732-2-12-66

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Rag Weed, Morning-Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray as re-growth develops. Use 1.5 fl. oz. in 1 gallon water or 1 cupful in 5 gallons of water.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, lawns, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable tree or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, pesticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra on lawns or turf where desirable bent-grass clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never use. Use spray equipment for applying 2,4-D formulations only to turf.


Elimination of shallow fish-bearing water at application rates recommended on this label may kill fish.

NET CONTENTS: 1 QUART



CAUTION: KEEP OUT OF REACH OF CHILDREN.
See side panel for additional cautions.

B. G. PRATT COMPANY, Paterson, New Jersey



DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Henbit, Wild Onion and many other broadleaf weeds in the Spring . . . FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

SCHEDULE A	SCHEDULE B	WATER GALLONS	AREA SQ. FT.
1 Tablespoonful	3.5 Teaspoonfuls	1	300
2 1/2 fl. oz.	0.8 fl. oz.	5	1000
5 Tablespoonfuls	6 Tablespoonfuls	30	6500
2 1/2 fl. oz.	3 fl. oz.		
1 pint	10 fl. oz.		

ACTIVE INGREDIENTS:
Dimethylamine Salt of Dicamba 3.43%
3,6-dichloro-o-anisic acid** 0.55
Dimethylamine Salts of Related Acids 17.22
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid* 78.80
INERT INGREDIENTS 100.00%

* Equivalent to 2.84% 3,6-dichloro-o-anisic acid (dicamba) or 0.26 pounds per gallon.
** Equivalent to 14.27% 2,4-Dichlorophenoxyacetic acid or 1.25 pounds per gallon.
U.S.D.A. Reg. No. 994-146

10732-2-12-66