

B G PRATT COMPANY

2004 Periody First Avenue Potens in the 2 de 1, 37502

plums. Use 2 gallons per 100 gallons of water. Apply in dormant or late dormant period.

HARMFUL IF SWALLOWED. May be absorbed through skin. Avoid breathing vapor or spray mist. Avoid centact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

lighly toxic to fish and wildlife. Birds feeding on treated areas may be killed.

Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Do not reuse empty containers. Destroy by crushing or perforating and burying in a safe place.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk-white spray mixture results, provided the spray iniviture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitation running, is not milk-white, do not use the resulting spray or the remainder of this product. Do not mix more material than can be applied in one day.

Do not apply within 2 weeks of a sulfur spray or within A days of Captan.

DO NOT STORE AT TEMPERATURES BELOW 32 4

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR

NOTICE: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

B. G. PRATT COMPANY, PATERSON, N. J. 07509



See side panel for additional autions.

NET CONTENTS: 1 LIGHT SALLON

U.S.D.A. Reg. No. 904-157

DIRECTION

Fill the spray tank about one tator and pour in the requ SPRAY. If other spray mate mix the wettable powder or water thoroughly, and then a oil. 2 gallons per 100 gallo fi oz. (5 tablespoon uls) per 100 gallo fi oz.

DIRECTIC

Spray on warm days in late fore the buds separate interfreezing weather.

ORNAMENTALS: Taxus, Arb p'e Elm, Catalpa, Boxwood Willow, Dogwood, Euonymu canium, Magnolla, Tulip tre European Fruit. Hard or Arr San Jose, Pine Needle, Jul ornamentals, use 2 to 3 gal evergreen ornamentals, us water. Apply as domant ar nights has passed but be Spray on a clear mild day noon. Be cautious with oil jured plants such as sugar ory, walnut, and butternut buds and remove blue cold

APPLES & PEARS: For ove Mite, San Jose Scale, Oyst apples and Pear Psylla on lons of water. Make a sindormant period. Do not ap; period.

PEACHES, PLUMS & CHE European Red Mits, San J Aphid on cherries and O



Total 100.00%

O' --- Totroothyl S,S' --methylane bispho Pet. No. 2,873,228

ieum Oil Specifications. Iphonated Residue (Min.) 92%

Viscosity 60 sees

ins 0,15 pounds Ethion per gallen U.S.D.A. Reg. No. 904-157

> See side panel for additional sautions. NET CONTENTS: 1 LIGHT CALLON

DIRECTIONS FOR MIXING

Fill the spray tank about one third full of water, start the agitator and pour in the required amounts of PRATT 6E OIL SPRAY. If other spray materials are to be used with it, first mix the wettable powder or emulsifiable concentrate in the water thoroughly, and then as the tank is being filled add the oil. 2 gallons per 100 gallons of water is equivalent to 2½ fl. oz. (5 tablespoonfuls) per gallon.

DIRECTIONS FOR USE

Spray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather.

ORNAMENTALS: Taxus, Arborvitae, Magnolia, Tulip tree, Maple Elm, Catalpa, Boxwood, Privet, Oak, Ash, Poplar, Birch, Willow, Dogwood, Europymus, Pine, Juniper, Spit Scales: Lecanium, Magnolla, Tulip tree, Cottony Maple, European Elin, European Fruit. Hard or Armored Scales: Oystershell, Scurly, San Jose, Pine Needle, Juniper, Europymus—For deciduous ornamentals, use 2 to 3 gallons per 100 gallons of water. For evergreen ornamentals, use 2 gallons per 100 gallons of water. Apply as dominant spray after most danger of freezing nights has passed but before buds have begun to open. nights has passed but before buds have begun to open. Spray on a clear mild day and do NOT spray in late afternoon. Be cautious with oil sprays. Do NOT use on easily injured plants such as sugar and Japanese maple, beech, hickory, walnut, and butternut. Oil will injure Douglas fir flower buds and remove blue color from Colorado blue spruce.

APPLES & PEARS: For overwintering eggs of European Rad Mite, San Jose Scale, Oystershell Scale, also Apple Aphid on apples and Pear Psylla on pears—use 2 gallons per 100 gallons of water. Make a single application during the delayed dormant period. Do not apply later than the delayed dormant

PFACHES, PLUMS & CHERRIES: For overwintering eggs of European Red Mits, San Jose Scale, also for Black Cherries Aphid on cherries and Oystershell Scale on peaches and



mike Jaks

FUNCTOES

SPECIALTIES

B G PRATT COMPANY

204 to arts Fort Avenue Pater on New Yorks, 07509

1 1 1 4 4 4 1/2

POS R + 194

plums. Use 2 gallons per 100 gallons of water. Apply in dormant or late dormant period.

HARMFUL IF SWALLOWED. May be absorbed through skin Avoid breathing vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

fighly toxic to fish and wildlife. Birds feeding on treated areas may be killed

Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Do not reuse empty containers. Destroy by crushing or perforating and burying in a safe place.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if when this product is added to the water in the spray tank, a uniform milk white spray mixture results, provided the spray mixture is being agitated. If the fispray mixture, after adding the oil to the spray tank with the agitation running, is not milk white, do not the the resulting spray or the remainder of this product. Do not mix more miterial than can be applied in one day.

Do not apply within 2 weeks of a suitur spray or within A days of Capten


DO NOT STORE AT TEMPERATURES BELOW 32 4.

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CHOP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR

NOTICE: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith

B. G. PRATT COMPANY, PATERSON, N. J. 07509

-1973 B & P Co



See side panel for additional coutions.

NET CONTENTS: 1 LIQUID GALLON

Trootices Mountaining Some 1901 -

Fill the tator SPRA) mix th water oil. 2 fl oz.

> Spray fore th

> freezin

ORNAI
ple Eli
Willow,
canium
Europe
San Jo
ornam
evergre
water,
nights
Spray
noon,
jured ;
ory, wi
buds a

apples lons o dorma period PEACI Europ Aphid

APPLE Mite,

T COMPANY

Paterson New Jersey 07509

PO Box . .



DIRECTIONS OR MIXING
Fill the spray tank about one-third full of water, start the agretator and pour in the required amounts of PRATT 6E OIL SPRAY. If other spray materials are to be used with it, first mix the wettable powder or emulsifiable concentrate in the water thoroughly, and then as the tank is being filled add the oil. 2 gallons per 100 gallons of water is equivalent to 2½ fl oz (5 tablespoonfuls) per gallon.

DIRECTIONS FOR USE

Spray on warm days in late Winter or early_Spring, and before the buds separate into the clusters. Do not spray in freezing weather.

ORNAMENTALS: Taxus, Arborvitae, Magnolia, Tulip tree, Mape Elm, Catalpa, Boxwood, Privet, Oak, Ash, Poplar. Birch, Willow, Dogwood, Euonymus, Pine, Jumper. Sptt Scales: Lecanium, Magnolla, Tulip tree, Cottony Maple. European Elm, European Fruit. Hard or Armored Scales: Oystershell, Scurfy, San Jose, Pine Needle, Jumper, Euonymus—For deciduous ornamentals, use 2 to 3 gallons per 100 gallons of water. For evergreen ornamentals, use 2 gallons per 100 gallons of water. Apply as dommant apray after most danger of freezing nights has passed but before buds have begun to open. nights has passed but before buds have begun to open. Spray on a clear mild day and do NOT spray in late afternion. Be cautious with oil sprays. Do NOT use on easily injured plants such as sugar and Japanese maple, beech, hickcry, walnut, and butternut. Oil will injure Douglas fir flower buds and remove blue color from Colorado blue spruce.

APPLES & PEARS: For overwintering eggs of European Red Mite, San Jose Scale, Oystershell Scale, also Apple Aphid on apples and Pear Psylla on pears—use 2 gallons per 100 gallons of water. Make a single application during the delayed dormant period. Do not apply later than the d. ayed dormant

PEACHES, PLUMS & CHERRIES: For overwintering eggs of European Red Mits, San Jose Scale, also for Black Cherries Aphid on chernes and Oystershell Scale on peaches and

HARMFUL IF SWALLOWED. May be absorbed through skin Avoid threathing vapor or spray mist. Avoid intact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or for products.

I'ghly toke to fin and weldlife. Birds feer no on treated areas may be killed.

Do not containing the any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Proposed on the containers of post reuse empty containers. Destroy by crushing or perforating and burying in a safe place.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk-white spray mixture is being agitated. If the stray mixture, after adding the oil to the spray tank with the instation running, is not milk-white, do not use the resulting agree or the remainder of this product. Do not mix more many or the applied in one day.

I must apply within 2 weeks of a senter spray or within ${m Z}>{\it distance}$ Captan.

NOT STORE AT TEMPERATURES BELOW 32 F.

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESTRABLE RESULTS MAY OCCUR

NOTICE. Buyer a some care of the following control of the simple care of the simple car

IS C PRATTICE EMPANY PATERSON OF 1 0/509

B

W.61414.

plums. Use 2 gallons per 100 gallons of water. Apply in dormant or late dormant period.

HARMFUL IF SWALLOWED. May be absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

lighly toxic to fish and wildlife. Birds feeding on treated areas may be killed.

Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Do not reuse empty containers. Destroy by crushing or perforating and burying in a safe place.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk-white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitation running, is not milk-white, do not use the resulting spray or the remainder of this product. Do not mix more material than can be applied in on day.

Do not apply within 2 weeks of a sulfur spray or within days of Captan.


DO NOT STORE AT TEMPERATURES BELOW 32 1

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

NOTICE: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

B. G. PRATT COMPANY, PATERSON, N. J. 07509

1970 B G P Co



See side panel for additional cautions.

NET CONTENTS: 1 LIQUID GALLON

DIRECTI

Fill the spray tank about tator and pour in the r SPRAY. If other spray memory the wettable powder water thoroughly, and thoughly, agailous per 100 graph oz. (5 tablespoonfuls

DIREC

Spray on warm days in fore the buds separate freezing weather.

ORNAMENTALS: Taxus, ple Elm, Catalpa, Boxwo Willow, Dogwood, Euony canium, Magnotta, Tulip European Fruit. Hard or San Jose, Pine Needle, ornamentals, use 2 to 3 evergreen ornamentals, water. Apply as dormand nights has passed but Spray on a clear mild on son. Be cautious with jured plants such as sugory, walnut, and buttern buds and remove blue of

APPLES & PEARS: For Mite, San Jose Scale, O apples and Pear Psylla clons of water. Make a dormant period. Do not period.

PEACHES, PLUMS & C European Red Mits, Sa Aphid on cherries and



ACTIVE INGREDIENTS:

*Ethion
Petroleum Oil (Superior type Oil)
INERT INGREDIENTS:

1.65
Total 100.00%

*O,O,O' --- Tetraethyl S,S' --- methylene bisphosphorod thioate *U. S. Pat. No. 2,873,228

Petroleum Oil Specifications.

Unsulphonated Residue (Min.) 92% Viscosity 60 seconds contains 0.15 pounds Ethion per gallon

U.S.D.A. Reg. No. 904-157

See side panel for additional cautions.

NET CONTENTS: 1 LIQUID GALLON

DIRECTIONS FOR MIXING

Fill the spray tank about one-third full of water, start the agitator and pour in the required amounts of PRATT 6E OIL SPRAY. If other spray materials are to be used with it, first mix the wettable powder or emulsifiable concentrate in the water thoroughly, and then as the tank is being filled add the oil. 2 gallons per 100 gallons of water is equivalent to $2\frac{1}{2}$ fl. oz. (5 tablespoonfuls) per gallon.

DIRECTIONS FOR USE

Spray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather.

ORNAMENTALS: Taxus, Arborvitae, Magnolia, Tulip tree, Mape Elm, Catalpa, Boxwood, Privet, Oak, Ash, Poplar, Birch, Willow, Dogwood, Euonymus, Pine, Juniper. Soft Scates: Lecanium, Magnolla, Tulip tree, Cottony Maple, European Elm, European Fruit. Hard or Armored Scales: Oystershell, Scurfy, San Jose, Pine Needle, Juniper, Euonymus—For deciduous ornamentals, use 2 to 3 gallons per 100 gallons of water. For evergreen ornamentals, use 2 gallons per 100 gallons of water. Apply as domnant spray after most danger of freezing nights has passed but before buds have begun to open. Spray on a clear mild day and do NOT spray in late afternoon. Be cautious with oil sprays. Do NOT use on easily injured plants such as sugar and Japanese maple, beech, hickory, walnut, and butternut. Oil will injure Douglas fir flower buds and remove blue color from Colorado blue spruce

APPLES & PEARS: For overwintering eggs of European Red Mite, San Jose Scale, Oystershell Scale, also Apple Aphid on appies and Pear Psylla on pears—use 2 gallons per 100 gallons of water. Make a single application during the delayed dormant period. Do not apply later than the delayed dormant period.

PEACHES, PLUMS & CHERRIES: For overwintering eggs of European Red Mits, San Jose Scale, also for Black Cherries Aphid on cherries and Oystershell Scale on peaches and