PYRETHRUM FOGGING SPRAY

904-162

ONLY FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS (Statement For Sizes Over One Gallon)

Containing

PYRETHRUM WITH ACTIVATOR FOR HOME, FARM AND INDUSTRY

A contact spray for control of listed flying insects in home, restaurants, hotels, food storage warehouses, dairies, meat packing plants and similar establishments with electric mist spraying or fogging equipment. Also outdoor adult mosquito control.

ACTIVE INGREDIENTS:	BY WEIGHT
Pyrethrins	0.10%
Piperonyl Butoxide, Technical*	0.75%
Isoparaffinic Petroleum Hydrocarbons	99.15%
TOTAL	

*Consists of 0.60% (butylcarbityl)(6-propylpiperonyl) ether and 0.15% related compounds.

EPA Reg. No. 904-162

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poisen Control Center. Do not induce IF SWALLOWED:

vomiting. Vomiting may cause aspiration pneumonia. Do not induce vomiting or give anything to an unconscious or

convulsing person.

Flush with plenty of water. Get medical attention if IF IN EYES:

irritation persists.

Remove contaminated clothing. Immediately wash affected IF ON SKIN:

area with soap and water. Launder clothing before reusing.

Get medical attention if irritation persists.

Remove victim to fresh air. Apply espiration if indicated. IF INHALED:

Empty stomach if material was taken internally. NOTE TO PHYSICIAN:

Gastric lavage may be indicated. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be ad-

ministered in conjunction with atropine.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

AUG 1 2 1985

ACCEPTED

Under the Enderell se ticide

Pratt-Gabriel Division MILLER CHEMICAL & FERTILIZER CORPORATION Hanover, Pennsylvania 17331

> 1 GALLON LIQUID NET CONTENTS:

> > 5 GALLONS LIQUID

30 GALLONS LIQUID

55 GALLONS LIQUID

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION -- Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin and clothing. Avoid breathing vapors - spray mist. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Healt! under the provisions of 30 CFR Part 11. Wash thoroughly after handling. Wash hands, face and arms with soap and water before eating and smoking. Wear chemical resistant gloves and longsleeved work clothing. Change to clean clothing daily. Wash all contaminated clothing with soap and hot water before reuse. Do not remain in treated area. Ventilate the area thoroughly before re-entry. Do not apply directly on very young animals. Do not use in poultry houses. Remove pets, birds and cover fish aquariums before spraying. Do not apply in a food processing plant while plant is in operation. When applied in a food processing plant, remove or protect food during treatment. Facilities and equipment must be thoroughly washed with an acceptable detergent solution and then rinsed with potable water to remove all traces of contamination.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARD

Do not use or store near heat or open flame. Keep Container Closed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers. Do not reuse empty container. Plastic containers may be incinerated or burned if allowed by state and local authorities. If burned, stay out of smoke. Do not reuse empty container.

DIRECTIONS

OUTDOOR SPRAY

TO REDUCE THE ANNOYANCE OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS around patios, swimming pools, home yards, and outdoor areas in parks and playgrounds. When the air is still, spray as a fine mist or fog to penetrate hiding places such as grasses, shrubs and ornamentals around homes, picnic areas, stadiums and open air theaters. Treat as large an area as possible surrounding the patio, pool or grounds, where insect protection is desired. Repeated applications may be necessary when infestation is heavy. Apply at the rate of 1.5 gallons per acre (4.4 fl. oz. per 1,000 sq. ft.).

Do not spray flowers and ornamentals at close range. Allow only the fine mist or "dry" feg to contact desirable plants. Direct spray away from the swimming pool. Application to any body of water is prohibited.

INDOOR SPRAY

FOR CONTROL OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS in homes, food establishments, farm buildings and warehouses: Close all windows and stop all ventilators. Pilot lights should be extinguished. Remove pets, birds and cover fish aquariums before spraying. Allow only fine mist or "dry" fog to contact fabric or surfaces that might be affected by the oil base. Direct fine mist or fog upward toward the center of the room and around the sides until the area is filled with fog or mist particles. Keep area closed for 10 to 15 minutes. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Clean up and destroy fallen insects.

CONTACT SPRAY -- FOR USE WITH SPRAYERS THAT GIVE COARSE WETTING SPRAY PARTICLES TO KILL COCKROACHES AND ACCESSIBLE STAGES OF GRANARY INSECTS: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RED FLOUR BEETLES, INDIAN MEAL MOTH LARVAE AND ADULTS, CADELLES, GRAIN MITES, AND SAW-TOOTHED GRAIN BEETLES: Cover or remove exposed food. Spray outside walls of grain bins, floorings, about the base of milling machinery, walls, baseboards, areas behind cabinets, around pipes, and inside surfaces of boxcars and trucks using 1 gallon of spray per 1,000 sq. ft. of space. Apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream in points between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, conduits, motor housings, and junction or switch boxes. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in "on" position. Repeat applications as necessary. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

CONTACT AND SPACE SPRAY -- TO KIL. ROACHES, SILVERFISH, EARWIGS, ANTS, AND SPIDERS IN SHELTERED SITUATIONS: Adjust spraying device to deliver a coarse wet spray. Apply as a contact spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Then adjust sprayer to deliver aerosol size particles. Apply as a space spray at the rate of 3 to 6 fl. oz. per 1,000 cu. ft. of room space. Repeat dosage as necessary for control. Leave the room closed for 1 hour. Do not remain in treated area. Ventilate the area thoroughly before re-entry.

FOR WASPS AND SPIDERS: Spray insects directly and apply heavy spray to nests and webs. The most effective wasp control is obtained by spraying after dark when all insects have returned to the nest.

FOR CONTROL OF CARPET BEETLES: Clean all rugs, carpets and fabrics to be treated. Spray both surfaces of carpet and floor space beneath evenly with a fine mist spray. Particular attention should be paid to cracks and crevices around the base-boards.

ANIMAL SPRAY FOR HORSES AND DAIRY AND BEEF CATTLE: For protection against Stable-flies, Hornflies and Houseflies attacking animals, spray the animal against the grain of the hair, directing a fine mist around the base of the horns, on the legs and along the sides and belly and over the back. Do not spray udders. Apply just the enough to wet the ends of the hair but not enough to wer the hide. One or two ounces of spray is sufficient to treat an adult cow. Apply once daily or as necessary.

LIMITED WARRANTY: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.