

RATT

EC2 METHOXYCHLOR INSECT SPRAY

**USED BY ARBORISTS —
A SUBSTITUTE FOR DDT ON
SHADE & FRUIT TREES — ORNAMENTALS**

An Emulsifiable Concentrate
Contains — 2 Pounds of Technical Methoxychlor Per Gallon

Methoxychlor, Technical	25.0%
Emulsifier	72.0%
Other Ingredients	3.0%
TOTAL	100.0%

Methoxychlor, technical is equivalent to 22.2% 2,2-bis (p-Methoxy-1,1-Trichloroethane and 3% other isomers and reaction products.

USDA Reg. No. 904-165

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional CAUTIONS

DIRECTIONS

Please read entire label BEFORE using.

CONTROL FRUIT TREE INSECTS

Apply only to listed fruit crops. (Apples, pears, peaches, apricots, cherries and plums.) Apply 2 to 3 quarts of Concentrate per 100 gallons in a Hydraulic Tank to cover all surfaces of foliage and fruit. Begin at petal fall and repeat at 7 to 14 day intervals as needed to control apple maggot, cherry fruit worm, codling moth, leaf hoppers, plum curculio, tent caterpillar and Japanese beetle. Do not apply later than 7 days before harvest on apples, cherries, pears, plums or later than 21 days of harvest on peaches and apricots. DO NOT USE with materials containing sulfur or other substances incompatible with oil. DO NOT USE on crops within 14 days of application of any material containing sulfur.

AREA CONTROL OF ADULT MOSQUITOES

Dilute 5 gallons of concentrate in 20 gallons of water (5% actual Methoxychlor by weight.) Apply at the rate of ½ gallon of spray mix per acre. Adjust Mist Blower to deliver a fine mist and apply to penetrate the areas occupied by adult mosquitoes. For best results misting should be done during the cooler hours of the night or early morning when the wind is not in excess of 3 miles per hour. Timing should coincide with the greatest mosquito activity. Repeat as necessary. Special care should be taken when applying space treatment to large areas to avoid drift over streams, lakes and ponds. Do not apply this treatment where food crops are grown.

NOTE:

DO NOT store at temperatures below 20 degrees F. If exposed to prolonged cold, allow to warm to 50 degrees F, and agitate well before using. Observe the usual precautions concerning application of oil-containing products to crops during high temperatures, since many crops are then more susceptible to solvent injury.

CAUTION

Keep out of reach of children. DO NOT use on humans. DO NOT get on skin, may cause irritation. In case of contact, wash with soap and water. DO NOT breathe spray mist. DO NOT contaminate feed or foodstuffs. DO NOT store or use near fire or open flame.

This product is toxic to fish. Keep out of streams, lakes or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. Do not reuse empty drum. Return drum to reconditioner or destroy by perforating or crushing and burying in a safe place.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

NOTICE

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

B. G. PRATT COMPANY, PATERSON, N. J. 07509

©1969 B. G. P. Co.

122-6M-4/69-W4151