

February 5, 1969

PRATTtm
LOGO2 0 2 K
S H A D E T R E E
I N S E C T S P R A YAn Emulsifiable Foliar Spray
For Hydraulic Sprayers and Mist BlowersContains Lindane-Malathion-Kelthane^R
2.5 pounds Malathion and 0.83 pounds Lindane
per gallon.

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride from Lindane	9.6 %
Malathion*	29.00%
1,1-Bis(p-Chlorophenyl)-2,2,2-trichloroethanol	3.00%
Xylene	55.40%

INERT INGREDIENT

Total	3.00%
	<u>100.00%</u>

U.S.D.A. Reg. No. 904-x

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate
^RKelthane Reg. trademark Rohm and Haas.


CAUTION: KEEP OUT OF REACH OF CHILDREN

See right side panel for additional cautions.

B. G. PRATT COMPANY

PATERSON, NEW JERSEY 07509

NET CONTENTS: 55 GALLONS



DIRECTIONS

As a general foliar spray on shade trees and ornamentals, use 2.5 pints with 100 gallons water in a Hydraulic Sprayer or 1 to 1½ gallons per 50 gallons of spray mix in Mist Blower. Apply thoroughly to all plant surfaces when insects appear. Repeat as necessary.

List of Insects Controlled

<u>Aphids</u> <u>Mites</u>	<u>Most Shade Trees & Ornamentals</u>	When insects appear.
<u>Lacebug</u>	<u>Andromeda, Azalea, Hawthorn, Oak, Sycamore</u>	When small nymphs appear.
<u>Cankerworm</u> <u>Tent Caterpillar</u> <u>Gypsy Moth</u> <u>Elm Leaf Beetle</u> <u>Willow Leaf Beetle</u>	<u>Most Deciduous Shade Trees & Shrubs</u>	When small worms first appear.
<u>Japanese Beetle adult</u> <u>Certain Leaf-Choppers</u> <u>Mealy Bug</u>	<u>Many Deciduous Shade Trees & Shrubs</u>	When insects appear.
<u>Leaf Miners</u>	<u>Birches</u> <u>Holly</u> <u>Boxwood</u>	Early to mid-May Repeat in 10 days.
<u>Sawflies</u> <u>European Pine Shoot Moth</u>	<u>All Pines</u>	When first small worms appear.
<u>Oyster Shell</u> <u>Euonymus</u> <u>Pine Needle Sawfly</u> <u>Juniper Scales</u>	<u>Many Ornamentals & Shade Trees</u>	Apply when scale crawlers first appear.

AREA CONTROL OF ADULT MOSQUITOES ON NON-AGRICULTURAL LAND

Mix 1 gallon 202K SHADE TREE INSECT SPRAY with water to make 50 gallons of spray. Adjust mist blower to deliver a fine mist and apply to penetrate the areas occupied by adult mosquitoes. For best results, misting should be done during the cooler hours of the night or early morning, when the wind is not in excess of 3 miles per hour. Timing should coincide with the greatest mosquito activity. Apply at the rate of 6 to 8 gallons of dilute spray per acre. Repeat as necessary.

DIRECTIONS

As a general foliar spray on shade trees and ornamentals, use 2-3 pints with 100 gallons water in a Hydraulic Sprayer or 1 to 1½ gallons per 50 gallons of spray mix in Mist Blower. Apply thoroughly to all plant surfaces when insects appear. Repeat as necessary.

List of Insects Controlled

<u>Aphids</u> <u>Mites</u>	<u>Most Shade Trees & Ornamentals</u>	When insect appear.
<u>Lacebug</u>	<u>Andromeda, Azalea, Hawthorn, Oak, Sycamore</u>	When small nymphs appear.
<u>Cankerworm</u> <u>Tent Caterpillar</u> <u>Gypsy Moth</u> <u>Elm Leaf Beetle</u> <u>Willow Leaf Beetle</u>	<u>Most Deciduous Shade Trees & Shrubs</u>	When small worms first appear.
<u>Japanese Beetle adult</u> <u>Certain Leafhoppers</u> <u>Mealy Bug</u>	<u>Many Deciduous Shade Trees & Shrubs</u>	When insects appear.
<u>Leaf Miners</u>	<u>Birches Holly Boxwood</u>	Early to mid May Repeat in 10 days.
<u>Sawflies</u> <u>European Pine Shoot Moth</u>	<u>All Pines</u>	When first small worms appear.
<u>Oyster Shell</u> <u>Euonymus</u> <u>Pine Needle Saw</u> <u>Juniper Scales</u>	<u>Many Ornamentals & Shade Trees</u>	Apply when scale crawlers first appear.

AREA CONTROL OF ADULT MOSQUITOES ON NON-AGRICULTURAL LAND

Mix 1 gallon 202K SHADE TREE INSECT SPRAY with water to make 50 gallons of spray. Adjust mist blower to deliver a fine mist and apply to penetrate the areas occupied by adult mosquitoes. For best results, misting should be done during the cooler hours of the night or early morning, when the wind is not in excess of 3 miles per hour. Timing should coincide with the greatest mosquito activity. Apply at the rate of 6 to 8 gallons of dilute spray per acre. Repeat as necessary.

RIGHT SIDE PANEL

February 5, 1969

The dilutions given for Mist Blowers are based on 8 times the concentration used for Hydraulic Sprayers. Adjustments should be made in accordance with the manufacturer's directions for the capacity of each Mist Blower.

CAUTION: Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, wash with soap and water; for eyes, get medical attention. Store away from children and domestic animals. Avoid contamination of feed and food stuffs. Solvents are inflammable. **KEEP CONTAINER CLOSED. DO NOT STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish and wild life. Do not let spray get into bird baths or fish ponds. Birds feeding in treated areas may be harmed. **KEEP OUT OF ANY BODY OF WATER.**


This product is highly toxic to bees exposed to direct treatment or residues left on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

Do not apply in any manner not specified on the label. Wash thoroughly after using.

Injury may occur on certain ferns, some species of Crassula and Canaerti Junipers.

Do not apply during periods of high humid or excessive drought.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



February 5, 1969

The dilutions given for Mist Blowers are based on 8 times the concentration used for Hydraulic Sprayers. Adjustments should be made in accordance with the manufacturer's directions for the capacity of each Mist Blower.

CAUTION: Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, wash with soap and water; for eyes, get medical attention. Store away from children and domestic animals. Avoid contamination of feed and food stuffs. Solvents are inflammable. **KEEP CONTAINER CLOSED. DO NOT STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish and wild life. Do not let spray get into bird baths or fish ponds. Birds feeding in treated areas may be harmed. **KEEP OUT OF ANY BODY OF WATER.**

This product is highly toxic to bees exposed to direct treatment or residues left on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

Do not apply in any manner not specified on the label. Wash thoroughly after using.

Injury may occur on certain ferns, some species of Crassula and Canaerti Junipers.

Do not apply during periods of high humid or excessive drought.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

