



SHOO FLY™

Bug *SHOOT*
Fly Spray Concentrate

Mixes with water to make a contact spray for control of certain insects* in and around homes, industrial plants, hotels, warehouses, barns, dairies, and kennels. Also, for direct application to horses, dairy and beef cattle.

*As listed on back panel

ACTIVE INGREDIENTS	By Weight
Pyrethrins	1.00%
Technical Piperonyl Butoxide*	10.00%
Isoparaffinic petroleum hydrocarbons	76.50%
INERT INGREDIENTS	12.50%
TOTAL	100.00%

NET CONTENTS:

*Equivalent to 8.0% of (Butylcarbityl) (6-propylpiperonyl) ether and to 2.0% of related compounds.

E.P.A. Reg. No. 904-171-9803

CAUTION: Keep Out Of Reach Of Children
See back panel for complete "CAUTION" statement

FORMULATED FOR MUNICIPAL,
PROFESSIONAL & INDUSTRIAL USE EXCLUSIVELY

MuniChem Corporation

SPARKS, NEVADA • MILWAUKEE, WIS.

ACCEPTED
9/24/77

[904-171] 2092

~~SHOO-FLY~~ BUG SHOT
Fly Spray Concentrate

DIRECTIONS

FOR OUTDOOR USE: Mix 1 part concentrate with 9 parts of water and apply as a fine mist to reduce the annoyance of FLIES, MOSQUITOES, GNATS AND ADULT MOTHS around patios, swimming pools, home yards and outdoor areas in parks and playgrounds. Spray or fog to penetrate hiding places such as grasses, shrubs, and ornamentals around homes, picnic areas, stadiums and open air theaters. Treat as large an area as possible surrounding the patio, pool or grounds where insect protection is desired. Repeated application may be necessary when infestation is heavy. Do not apply more than 1.5 gallons of diluted spray per acre.

Do not spray flowers and ornamentals at close range. Allow only the fine mist to contact desirable plants. Direct spray away from the swimming pool.

Horseflies, Deerflies, Stableflies and Houseflies: To kill and repel these insects on HORSES, DOGS, DAIRY & BEEF CATTLE dilute 1 gallon of ~~SHOO-FLY~~ ^{BUG SHOT} with 9 gallons of water. Apply diluted spray at the rate of 2 ounces per large animal for daily treatment or as a wet spray to all portions of the animal, using up to 1-2 quarts per animal the size of an adult cow. This treatment will give 3-8 day protection from stableflies and houseflies and up to 3 days protection from horseflies and deerflies. Repeat as necessary. Horses or nervous animals may be sponged with a 1-9 solution.

For automatic treadle sprayers use without dilution. Each nozzle should be set to deliver 0.3-0.5 mls. Animals should make 2 or 3 round trips daily.

Sucking and Biting Lice: For control of sucking lice and biting lice on CATTLE, POULTRY, DOGS, HORSES, SHEEP, GOATS and HOGS use ~~SHOO-FLY~~ ^{BUG SHOT} at a 1-100 dilution as a dip. For POULTRY LICE treat birds by submerging in a barrel, tank or tub of the solution, holding the wings away from the body to insure thorough wetting of all infested areas. Avoid chilling of the fowl by dipping on a warm day or in a warm building.

For fly control in and around BARNs, DAIRIES, CREAMERIES, FOOD PLANTS, and KENNELS: Close all windows and doors and stop all ventilators. Set machine to deliver fine mist. Direct spray upward toward center of the room and around the sides until the treated area is filled with mist particles. Dilute 1 part of ~~SHOO-FLY~~ ^{BUG SHOT} with 9 parts of water. Use 1/2 to 1 ounce of spray for each 1000 cu. ft. of space. Keep area closed for 10 to 15 minutes. Clean up and destroy fallen insects.

For control of ROACHES, SILVERFISH, ANTS, dilute 1-9 with water, directing coarse spray into all cracks and crevices where insects hide and on areas frequented by them. Apply spray to surfaces to the point of runoff.

If fog space treatment for roach control is desired, set machine for fine mist, using 2 to 3 ounces per 1000 cu. ft. It is also necessary to apply surface treatment as directed above. Repeat treatment for crawling insects as necessary.

CAUTION: HARMFUL IF SWALLOWED. If swallowed do not induce vomiting. Keep undiluted material closed and away from open flame. Do not contaminate feed or food stuffs. Avoid spraying directly on foodstuffs. Do not apply directly on very young animals. Do not apply as a space spray in a food processing plant while plant is in operation. Remove or protect food during treatment. After treatment wash all equipment, benches, shelves, etc. with potable water. Do not apply as a residual spray in edible product areas of food processing plants. Do not remain in an enclosed area immediately after fogging. Ventilate when treatment is completed. Apply this product only as specified on this label.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

^{BUG SHOT}
* ~~SHOO-FLY~~ is the trademark of MUNICHEM CORP.
© COPYRIGHT 1973 BY MUNICHEM CORP.

Sold By

12/73 1M LL



MuniChem Corporation

SPARKS, NEVADA • MILWAUKEE, WIS.