

1% ROTENONE
DUST OR SPRAY

A BOTANICAL INSECTICIDE FOR CONTROL OF CERTAIN CHEWING INSECTS
ON • FLOWERS • VEGETABLES
AS LISTED ON BACK PANEL

ACTIVE INGREDIENTS:	BY WEIGHT
Rotenone.....	1.0%
Other Cube Resins.....	2.0%
INERT INGREDIENTS:.....	97.0%
TOTAL.....	100.0%

EPA Reg. No. 904-176

EPA Est. No. 4-NY-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.


STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Call a Physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF INHALED: Remove victim to fresh air.
- IF IN EYES: Flush with plenty of water. Get medical attention.
- IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For
Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

- NET WEIGHT: ONE POUND
- 4 POUNDS
- 10 POUNDS
- 25 POUNDS
- 50 POUNDS



ACCEPTED

JUL 19 1985

U.S. Environmental Protection Agency
Federal Insecticide, Fungicide, and Rodenticide Act
Registered under the pesticide
Reg. No. 904-176

INSECTS CONTROLLED ON VEGETABLES

<u>ASPARAGUS</u> Asparagus Beetle	<u>BEANS</u> Mexican Bean Beetle Bean Leaf Beetle	<u>BEETS (Garden), CARROTS, EGGPLANT & PEPPERS</u> Flea Beetle	<u>TOMATOES</u> Aphids Fruit Worm Flea Beetle
--------------------------------------	---------------------------------------------------------	---------------------------------------------------------------------------	--------------------------------------------------------

<u>CABBAGE, CAULIFLOWER; BROCCOLI & BRUSSELS SPROUTS</u> Imported Cabbage Worm Cabbage Looper Begin applications as soon as plants are established in the field.	<u>MELONS, SQUASH, & CUCUMBERS</u> Striped Cucumber Beetle Spotted Cucumber Beetle To reduce damage, keep plants thoroughly covered with dust or spray throughout the season.	<u>PEAS</u> Pea Aphids Pea Weevil
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------

DO NOT APPLY WITHIN 24 HOURS OF HARVEST OF VEGETABLES. REMOVE ANY SPRAY RESIDUE AT HARVEST BY WASHING EDIBLE PORTIONS.

This material is packed at the full net weight as listed. It is subject to settling and may appear less than full.

LIMITED WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

