




SULFUR

PRATT
WETTABLE
SULFUR
*Dust or
Spray*

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions



LF3632-3-2-73



FINENESS: Not less than 95% passing 325 mesh.
ACTIVE INGREDIENT:

Sulfur	Not less than	95.0%
INERT INGREDIENTS:	Not more than	5.0%
		100.0%

A SULFUR FUNGICIDE FOR USE IN THE CONTROL OF BLACK SPOT AND MILDEW ON ROSES, APPLE SCAB AND SIMILAR

**FUNGUS DISEASES ON
FRUITS and FLOWERS**

NET WEIGHT: 2 LBS.

914-177

DIRECTIONS

This material may be applied as a dust or spray for the control of such diseases as Black Spot on Roses; Powdery Mildew of Roses, Phlox, Zinnias, Sweet Peas and Lilacs; Rust on Snapdragons and Carnations; Brown Rot of Stone Fruits; Scab on Apples and Pears and to aid in keeping Red Spider Mites in check on evergreens.

Apply every week or ten days during the growing season, or more often during rainy weather. Cover all parts of the plant with a thin film of dust or spray. On stone fruits for Brown Rot apply just as trees are coming into bloom, and after the petals fall, and at intervals of two weeks during the season. During the last month before harvest dust lightly every week or ten days.

For scab on apples and pears make weekly applications from the time new growth starts until 30 days after petal fall. Use more frequently during rainy periods. For red spider mite delay application on evergreens until the new growth has hardened.

DUSTING — Material is ready for use and is applied with a duster. Dust the plants lightly from two sides when the air is quiet.

SPRAYING — Add the required amount of material to water and stir thoroughly. Pour the mixture into the sprayer and keep agitated while spraying. In small sprayers shake frequently for agitation.

DILUTIONS — Apples and pears 6 to 8 pounds in 100 gallons of water — 4 level tablespoonfuls in each gallon of water.

Peaches, plums and prunes — also for red spider mite — 4 to 6 pounds in 100 gallons of water — 3 level tablespoonfuls in each gallon of water.

CAUTION: Do not use sulfur on apple foliage within two weeks of an oil spray.

Avoid spraying when temperature is warmer than 85°F.

Do not reuse empty container. Wrap in newspaper and place in trashcan for safe disposal.

NOTICE: User assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

E.P.A. Reg. No. 904-177

PRATT
WETTABLE
SULFUR

*Dust or
Spray*

PRATT