

Begin when plants are small. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Do not use on beans.

ABBAGE: Alternaria Leaf Spot, Downy Mildew. Begin 7 to 10 days after planting. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest.

CURRY: Sweet. Hemitelespium Blight. Begin when plants are 6 inches high. Repeat at 7 to 10 day intervals until harvest. Do not use within 10 days of harvest.

CUMBERS: Alternaria. Mosaic. Downy Mildew. Leaf Spot. Anthracnose. Leaf Blight. Downy Mildew. Fruit Rot. Begin when diseases threaten. Do not use within 10 days of harvest. Repeat at 7 to 10 day intervals until severe disease conditions. Do not use within 5 days of harvest.

CUCUMBER: Anthracnose. Early Blight. Alternaria. Phomopsis. Blight of Fruit. Botry. Begin when fruit starts to form. Repeat at 7 to 10 day intervals.

ENDIVE: LITTLE. Downy Mildew. Begin when disease appears. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove diseased leaves and stems by stripping and burning. Do not use water culture and avoid spraying with other effective means.

PEPPERS: Anthracnose. Cercospora Leaf Spot. Phytophthora Blight. Botry. Begin when disease threatens. Repeat at 7 to 10 day intervals.

PUMPKIN: Summer and Winter. Anthracnose. Downy Mildew. Begin when plant starts to form. Repeat at 7 to 10 day intervals. Do not use within 5 days of harvest.

TOMATOES: Greenhouse and Field. Cadusporium Leaf Mold. Early and Late Blights. Gray Leaf Spot. Stemphylium. Septoria Leaf Spot. Begin at first fruit cluster and repeat at 7 to 10 day intervals. Do not use within 5 days of harvest. Apply 1 to 2 quarts of diluted spray per acre.

CAUTION: May cause irritation to eyes, nose, throat and skin. Avoid breathing dusts. Do not be absorbed through skin. Do not spray on clothes, skin or clothing. Do not use on food plants. Do not use on plants that get medicinal benefit. Keep away from fire and sparks.

This product is for use only on plants. Do not apply to any body of water. Apply this product as specified on label. Do not apply where prohibited. Do not use on plants when weather conditions favor drift. Do not use on plants that are damaged by frost. Do not use on plants that are damaged by insects. Do not use on plants that are damaged by other diseases. Do not use on plants that are damaged by other pests.

B. G. PRATT DIVISION
LABORATORY AND SALES DEPARTMENT

B. G. PRATT
SINCE 1904

Liquid MANEB

PLANT FUNGICIDE

CONTROLS COMMON DISEASES ON TURF, ORNAMENTALS AND VEGETABLES

(As listed on back panel)

55-5

ACTIVE INGREDIENT	By Weight
Maneb (Manganese Ethylenebisdithiocarbamate)	37%
(Total manganese as metallic—7.6%)	
INERT INGREDIENTS:	63%
	Total.....100%

E.P.A. Reg. No. 904-190

See side panel for additional CAUTIONS
NET CONTENTS 1 PINT

SHAKE WELL BEFORE USING

GENERAL DIRECTIONS
Mix with water according to directions. Apply thoroughly to all plant surfaces.

FLOWER AND ORNAMENTAL USES
GENERAL DILUTION: 1.500 (2 Teaspoons per gallon) 1 pint per 5 gallons.

CARNATIONS, DAHLIAS: Alternaria Leaf Spot, Anthracnose, Botry. Begin when new growth starts. Repeat weekly through late summer.

GLADIOLI: Anthracnose. Begin when bud opens. Repeat weekly through late summer.

ADONIS: Botry. Blight, Curculionid and Stemphylium Leaf Spot. Begin when flower spikes are developing. Repeat weekly at weekly intervals.

GLADIOLI: Botry. Begin with new growth. Repeat weekly.

PANSIES: Anthracnose. Begin with new growth. Repeat weekly.

ROSES: Downy Mildew, Botry. Blight, Phyllosticta Blight. Apply in early spring and early fall.

WILDS: Botry. Blight, Curculionid Leaf Spot, Rust, Gall. Begin when first leaves unfold. Repeat at 7 to 10 day intervals.

WATER LILIES: Botry. Begin with new growth. Repeat weekly.

TURF AND GRASS USES
Apply 1 to 2 quarts of diluted spray per acre. Mix prescribed volume of liquid with sufficient water to cover 1,000 square feet. Use 5 to 10 gallons of water. Do not use on lawns where children or pets play. Do not use on lawns where children or pets play. Do not use on lawns where children or pets play.

VEGETABLE USES
Apply 1 to 2 quarts of diluted spray per acre. Mix prescribed volume of liquid with sufficient water to cover 1,000 square feet. Use 5 to 10 gallons of water. Do not use on lawns where children or pets play. Do not use on lawns where children or pets play.

1077