

PRATT

SINCE 1904
TM

5% SEVIN* DUST

CONTROLS MANY
INSECTS† ON

- VEGETABLES
- SMALL FRUITS
- FLOWERS
- TREES • SHRUBS
- PETS

ACTIVE INGREDIENT	By Weight
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS	95.0%
TOTAL 100.0%	

**Registered trademark of Union Carbide Corporation for carbaryl insecticide (1-naphthyl N-methylcarbamate). U. S. Pat. Nos. 2,903,478; 3,009,855.

†As listed in "DIRECTIONS."

E.P.A. Reg. No. 904-191

CAUTION: KEEP OUT OF REACH OF CHILDREN

See back panel for additional cautions.

NET WEIGHT — ONE POUND

Read entire label thoroughly before use. Use strictly according to label directions.
Observe CAUTION and LIMITATION statements carefully.

DIRECTIONS FOR USE

Apply this dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Repeat as necessary, usually at 7 day intervals. Do not apply in excess. Observe preharvest limitations.

HOME GARDENS. Apply with dust gun, or from shaker-top can.

ACREAGE APPLICATIONS. Apply 40 lbs. per acre to all crops listed on this label except as follows: Cucumbers, melons, pumpkins, squash (20 lbs. per acre); Peas (30 lbs. per acre)

VEGETABLE INSECTS

ASPARAGUS. Controls asparagus beetle, Apache cicada. Apply during cutting season. Do not repeat application within 3 days.

BEANS. (Green, lima, and snap beans; cowpeas, black-eyed peas). Controls Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs, tarnished plant bugs.

SWEET CORN. Controls corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetles, sap beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Application of this product to the tassel region of corn during the pollen shed season will seriously destroy bee populations.

CUCUMBER, MELONS, PUMPKIN, SQUASH. Controls pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during prolonged high humidity. SEVIN injures watermelons in Florida.

TOMATO, POTATO, EGGPLANT, PEAS, PEPPERS. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA. For corn earworm, stink bugs.

CARROTS, CHINESE CABBAGE, COLLARDS, DANDELIONS, ENDIVE (ESCAROLE), GARDEN BEETS, HORSERADISH, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY, SPINACH, SWISS CHARD, TURNIPS. For flea beetles, harlequin bug, leafhoppers, armyworm, imported cabbage worm, corn earworm, spittlebugs, tarnished plant bugs, stink bugs. To control small cabbage loopers, use on a 5 to 7 day schedule.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI. For flea beetles, harlequin bug, armyworms, imported cabbage worm, corn earworm. To control small cabbage loopers, use on a 5 to 7 day schedule.

SMALL FRUIT INSECTS

GRAPES. For grape berry moth, Japanese beetle leafroller, grape leafroller, grape leafhopper, grape vine sawfly. Apply just before first brood leafroller larvae emerges from eggs.

STRAWBERRIES. For meadow spittlebug, strawberry root weevil.

BLUEBERRIES. For blueberry maggot, cherry and blueberry sawfly.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, RASPBERRIES. For Japanese beetles, leaf rolling sawfly, aphids.

GRASSHOPPERS ON ABOVE CROPS. Apply as directed to control grasshoppers.

FLOWERS - SHRUBS - TREES

This dust can safely be used on a wide variety of trees. **ROSES, CARNATIONS, GLADIOLUS, CHRYSALEPIDAE, ARBORVITAE, JUNIPER, HYDRANGEA, AZALEA, DOGWOOD, BIRCH, PINES.** Do not use on **BOSQUET**.

For apple aphid, bagworm, birch leaf miner, boxwood leaf miner, elm leaf aphid, elm leaf miner, Japanese beetle, June beetles, lace bugs, leaf beetles, mimosa webworm, oak leaf miners, orange plant bugs, puss caterpillars, psyllids, rose aphid, sawflies, caterpillars, thorn bugs, thrips (exposed), willow sawfly.

PET USE

ON DOGS, CATS. For ticks and fleas dust liberally on the animal, avoiding the eyes, and rub thoroughly into the fur a few hours after treatment. Repeat at weekly intervals. For treatment of kittens and puppies under 4 weeks of age, use **IN and AROUND DOG and CAT QUARTERS** liberally in and around sleeping quarters, kennel, and under the cover of a shaker can, hand or dust gun.

LIMITATIONS OF USE BEFORE HARVEST

Allow the designated time limit shown between the crops listed below:

1 day: Asparagus, strawberries.

3 days: Broccoli, Brussels sprouts, cabbage, kohlrabi, garden beets (roots), horseradish, parsley (roots), turnip (roots).

7 days: Blackberries, boysenberries, dewberries, raspberries.

14 days: Chinese cabbage, collards, dandelion beets (tops), kale, leaf lettuce, mustard green, spinach, Swiss chard, turnips (tops).

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid contact with skin. Wash hands after use. Avoid contamination of food.

This product is highly toxic to bees exposed to on crops. Protective information may be obtained from Agricultural Extension Service.

Injury to tender foliage may occur if application is made wet or in the presence of high humidity of severe weather. Do not reuse container. Destroy when empty and place in trash can for disposal.

Note for Physician: Carbaryl is a moderate sedative. Atropine is antidotal.

NOTICE. Seller makes no warranty of any kind concerning effects of use of this product other than that stated on label. Buyer, or User, accepts all responsibility for use of this product.

B. G. PRATT DIVISION GABRIEL CHEMICALS LTD., PATERSON, N.J. 07509

Read entire label thoroughly before use. Use strictly according to label directions.
Observe CAUTION and LIMITATION statements carefully.

DIRECTIONS FOR USE

Apply this dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Repeat as necessary, usually at 7 day intervals. Do not apply in excess. Observe preharvest limitations.

HOME GARDENS. Apply with dust gun, or from shaker-top can.

ACREAGE APPLICATIONS. Apply 40 lbs. per acre to all crops listed on this label except as follows: Cucumbers, melons, pumpkins, squash (20 lbs. per acre); Peas (30 lbs. per acre)

VEGETABLE INSECTS

ASPARAGUS. Controls asparagus beetle, Apache cicada. Apply during cutting season. Do not repeat application within 3 days.

BEANS. (Green, lima, and snap beans; cowpeas, black-eyed peas). Controls Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs, tarnished plant bugs.

SWEET CORN. Controls corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetles, sap beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Application of this product to the tassel region of corn during the pollen shed season will seriously destroy bee populations.

CUCUMBER, MELONS, PUMPKIN, SQUASH. Controls pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during prolonged high humidity. SEVIN injures watermelons in Florida.

TOMATO, POTATO, EGGPLANT, PEAS, PEPPERS. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA. For corn earworm, stink bugs.

CARROTS, CHINESE CABBAGE, COLLARDS, DANDELIONS, ENDIVE (ESCAROLE), GARDEN BEETS, HORSERADISH, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY, SPINACH, SWISS CHARD, TURNIPS. For flea beetles, harlequin bug, leafhoppers, armyworm, imported cabbage worm, corn earworm, spittlebugs, tarnished plant bugs, stink bugs. To control small cabbage loopers, use on a 5 to 7 day schedule.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI. For flea beetles, harlequin bug, armyworms, imported cabbage worm, corn earworm. To control small cabbage loopers, use on a 5 to 7 day schedule.

SMALL FRUIT INSECTS

GRAPES. For grape berry moth, Japanese beetles, June beetles, red-banded leafroller, grape leafroller, grape leafhopper, grape leaf skeletonizer. Apply just before first brood leafroller larvae emerges from rolls and as needed for leafhoppers.

STRAWBERRIES. For meadow spittlebug, strawberry leafroller, strawberry weevil.

BLUEBERRIES. For blueberry maggot, cherry and cranberry fruitworm.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES,

RASPBERRIES. For Japanese beetles, leaf rollers, leafhoppers, raspberry aphids.

GRASSHOPPERS ON ABOVE CROPS. Apply as often as necessary on these crops to control grasshoppers.

FLOWERS - SHRUBS - TREES

This dust can safely be used on a wide variety of flowers, shrubs, and small trees. **ROSES, CARNATIONS, GLADIOLUS, CHRYSANTHEMUM, LILAC, ARBORVITAE, JUNIPER, HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH, PINES.** Do not use on **BOSTON IVY.**

For apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf roller, mealy bugs, mimosa webworm, oak leaf miners, orange tartrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

PET USE

ON DOGS, CATS. For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens and puppies under 4 weeks of age.

IN and AROUND DOG and CAT QUARTERS. For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

LIMITATIONS OF USE BEFORE HARVEST

Allow the designated time limit shown between application and harvest of the crops listed below:

1 day: Asparagus, strawberries.

3 days: Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, kohlrabi, garden beets (roots), horseradish, parsnips, radishes, rutabagas, salsify (roots), turnip (roots).

7 days: Blackberries, boysenberries, dewberries, loganberries, raspberries.

14 days: Chinese cabbage, collards, dandelion, endive (escarole), garden beets (tops), kale, leaf lettuce, mustard greens, parsley, salsify (tops), spinach, Swiss chard, turnips (tops).

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of food.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment. Do not reuse container. Destroy when empty. Wrap in newspaper and place in trash can for disposal.

Note for Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE. Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or User, accepts all responsibility for results due to misuse of this product.