

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals—CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes with plenty of water, wash skin with soap and water. If irritation develops, get medical attention. Avoid contamination of feed or food stuffs.
Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate domestic or irrigation water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with these directions.

STORAGE AND DISPOSAL

Keep from freezing. Do not store near food or foodstuffs. Do not store near fertilizers, insecticides or fungicides. **DO NOT REUSE EMPTY CONTAINER. WRAP AND PUT IN TRASH COLLECTION.**

OTHER USE PRECAUTIONS: Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solution near these plants. Whether using a hose-end sprayer or pressure-sprayer, do not use sprays that are fine because they can drift and cause damage to non-target plants. Coarse, drop-type sprays are less likely to drift. Use only lawn-type sprayers. Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful with drift on trees and other ornamental species. This product may be injurious to St. Augustine and Centipede grasses. Since injury can be expected, spot-spray weeds only with a pressure sprayer.

May be used on Bahia, Bermuda, Bluegrass, Fescue, Ryegrass, Zoysia lawns, vacant lots, playgrounds and other similar areas where broad leaf weeds are present. This product is especially effective on annual-broad leaf weeds such as Poinsettia, Poison Oak, Sheep Sorrel and Scurf. In addition, this product controls most other broadleaf weeds including sensitive weeds such as Clover, Dandelion, Knotweed and Plantain. Spray at any time during the growing season when weeds are actively growing. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, spot-treatment may be made at any time with a small pressure-sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or sods down in bare spots. For best results, make repeat sprays if applied during drought or 1 before rain. For best results, make before treating. Do not water lawn within 24 hours after treatment.

ACCEPTED
DEC 13 1988
Federal Insecticide, Fungicide, and Rodenticide Act
under
No. 201-112

CHICKWEED & CLOVER KILLER

Contains *TRIMEC* Herbicide:
2, 4-D, MCPP, Dicamba
1 Pint Covers 4,000 Sq. Ft.

Controls: BEDSTRAW, BLACK MEDIC, BUCKHORN, CHICORY, CHICKWEED, CLOVER, DANDELION, DOCK, GROUND IVY, HEAL-ALL, HENBIT, KNOTWEED, LAMBSQUARTER, LESPEDEZA, MALLOW, MORNING GLORY, PEPPERGRASS, PIGWEED, PLANTAINS, POISON IVY, POISON OAK, PURSLANE, RAGWEED, SHEPHERD'S PURSE, SPEEDWELL, SPURGE, WILD CARROT, WILD LETTUCE, WILD ONION, YARROW and many others.

ACTIVE INGREDIENTS

***Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	7.62%
***Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	3.66%
***Dimethylamine Salt of Dicamba (3,6-dichloro-2-anisic acid)	0.64%
INERT INGREDIENTS	87.81%
TOTAL	100.00%

THE PRODUCT CONTAINS:
***0.65 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 6.34%
***.26 lb. 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 3.02%
***0.09 lb. 3,6-dichloro-2-anisic acid per gallon or 0.64%
*Other Symbols: AOAC Method 6, D01-5
*Trade Name is a registered trademark of PBI GORDON CORP.

EPA Reg. No. 904-193 EPA Est. No. 72-PA-1
KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Drink 2 to 4 glasses of water immediately by sipping slowly. Do not induce vomiting. Give 1 glass every 15 minutes.
IF IN EYES: Flush with water for 15 minutes.
IF INHALED: Move to fresh air.
IF ON SKIN: Wash with soap and water.

See back panel for additional precautionary statements.
NET CONTENTS: 8 Fl. Oz. 1 Pint 1 Quart 1 Gallon



HOW MUCH TO USE (ENTIRE LAWN) HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of product (Column 1) in sprayer and add water to indicated level (Column 2) and spray on area shown in Column 3. For example, to spray 2,500 sq. ft. measure 10 ozs. (20 tbsps) in jar, add water to the 10 gallon level and mix. Turn on hose and spray as evenly as possible over 2,500 sq. ft.

NOTE: If you have a CHEMLAWN LIQUID APPLICATOR, use pint of Chickweed & Clover Killer in 2 1/2 gallons of spray and cover 4,000 sq. ft.

Column 1	Column 2	Column 3
Amt. Chickweed & Clover Killer to use (Tbsp.)	Fill Bottle To (Gallons)	Area to Treat (Sq. Ft.)
10	5	1,250
20	10	2,500
30	15	3,750
40	20 (Full)	5,000

PRESSURE-SPRAYERS: Measure the amount of Chickweed & Clover Killer in tank and dilute with water according to the following chart. Use a spray pattern that gives small spray droplets without any fine mist and cover the area indicated.

Amt. Chickweed & Clover Killer to use (Tbsp.)	Amt. of water in sprayer (Fl. Oz.)	Area to be Sprayed (Sq. Ft.)
8	3	750
32	16	3,000
64	32	6,000

SPOT-SPRAYING: To spray small areas or individual weeds, do not use a hose-end sprayer. Mix 1 oz. (2 tbsps) in a half-gallon of water in a pressure sprayer and spray, as required, being certain to wet each weed.

LIMITED WARRANTY: BUYER ASSUMES ALL RISKS OF USE, STORAGE AND HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

Manufactured By
PRATT-GABRIEL DIVISION
 MILLER CHEMICAL & FERTILIZER CORPORATION
 Hanover, Pennsylvania 17331 An Alco Standard Company

