

PRESSURE SPRAYERS Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated.

Amt. Turf Herbicide 6000 to use (Tbsp.)	(Fl. Oz.)	Amt. of water in sprayer: (Gallons)	Area to be Sprayed: (Sq. Feet)
2	1	1	190
4	2	2	380
6	3	3	760

HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required being certain to "wet" each weed.

CAUTION: Harmful if swallowed. Avoid skin, eye and clothing contact. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

CAUTION: Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

ACTIVE INGREDIENTS:	By Weight
*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	2.77%
**Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	6.10%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.63%
Dimethylamine salts of related compounds	0.08%
INERT INGREDIENTS	90.42%
Total	100.00%

*2-(2-methyl-4-chlorophenoxy)propionic acid equivalent to 0.19 pounds per gallon, equivalent to 2.22%.

**2,4-dichlorophenoxyacetic acid equivalent to 0.43 pounds per gallon, equivalent to 5.01%.

***3,6-dichloro-o-anisic acid equivalent to 0.045 pounds per gallon, equivalent to 0.52%.

KEEP FROM FREEZING
E.P.A. Reg. No. 904-194

B. G. PRATT DIVISION
GABRIEL CHEMICALS, LTD., PATERSON, N.J. 07509

PRATT

SINCE 1904

TURF HERBICIDE 6000

CONTAINS TRIMEC** HERBICIDE, 2,4-D,
MCPP AND DICAMBA

1 PINT COVERS
3000 Square Feet

- KILLS**
- CHICKWEED
 - DANDELION
 - KNOTWEED
 - POISON IVY
 - GROUND IVY
 - WILD ONION
 - CLOVER
 - SPURGE
 - PLANTAINS
 - HENBIT
 - RAGWEED
 - AND MOST OTHER BROADLEAF WEEDS

*TRIMEC is a registered trademark of P.B.I.—GORDON Corporation
U.S. Patent No. 3,284,186

E.P.A. REG. NO. 904-194

CAUTION:
KEEP OUT OF REACH OF CHILDREN

See back panel for complete caution statement.

NET CONTENTS: ONE PINT

OTHER WEEDS CONTROLLED: BESSIBAW, BLACK MEDIC, BUCKHORN, CHICORY, CHICKWEED, CLOVER, DANDELION, DOCK, GROUND IVY, HEAL-ALL, HENBIT, KNOTWEED, LAMBSQUARTER, LESPEDEZA, MALLOW, MORNING GLORY, PEPPERGRASS, PIGWEED, PLANTAINS, POISON IVY, POISON OAK, PURSLANE, RAGWEED, SHEPHERD'S PURSE, SPEEDWELL, SPURGE, WILD CARROT, WILD LETTUCE, WILD ONION, YARROW

DIRECTIONS FOR USE

WHERE TO USE: Turf Herbicide 6000 may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, Turf Herbicide 6000 Lawn Weed Killer controls most other broadleaved lawn weeds, including Dandelion and Plantain, and a partial list is given above. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed. Repeat if necessary for hard to kill weeds.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 2 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE—ENTIRE LAWN

HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example, to spray 2,500 sq. feet, measure 10 ozs. (20 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,900 sq. feet.

Col 1		Col 2	Col 3
Amt. Turf Herbicide 6000 to use (Tbsp.)	(Fl. Oz.)	Fill Bottle To: Gallons.	Area to Treat: (Sq. Feet)
10	5	5	950
20	10	10	1900
30	15	15	2850
40	20	20 full	3800

NOTE: Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

June 6, 1974
904-194

FRISSE SPRAYERS Measure indicated amount of chemical to be used and dilute with water according to the following chart. Use a spray pattern which gives spray droplets about 1/2 inch in diameter and cover the area indicated.


Amount of Herbicide 6000 to Use (lb/gal)	Amount of water in sprayer (Gallons)	Area to be Sprayed (Sq. Feet)
2	1	100
4	2	200
6	3	300

HOW MUCH TO USE - SPOT SPRAY
To spray small areas of individual weeds, do not use hose and sprayer. Mix 1 lb. of 6000 in 10 gal. water in pressure sprayer and spray as required being certain to "beat" each weed.

CAUTION: Hazardous if swallowed. Avoid skin and clothing contact in case of contact, flush with water. Do not contaminate food or foodstuffs. Do not get in eyes. Avoid drift of spray mist onto vegetables, ornamentals, plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or herbicides. Do not spray on chickens, ducks, geese, turkeys or other domestic birds. Do not spray on clovers or alfalfa. Do not spray on lawns or lawns where desirable plants are growing. Do not spray on lawns where desirable plants are growing. Do not spray on lawns where desirable plants are growing.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material in strict accordance with directions given hereon.
ACTIVE INGREDIENTS:
 - 2,2-Dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid 2.77%
 - Dimethylamine salt of 2,2-dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid 8.30%
 - Dimethylamine salt of 2,2-dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid 8.83%
 - Dimethylamine salt of 2,2-dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid 0.58%
 - Inert ingredients 89.52%
 Total 100.00%
 2,2-Dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid and its dimethylamine salt are equivalent to 2.27% active ingredient.
 2,2-Dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid and its dimethylamine salt are equivalent to 2.27% active ingredient.
 2,2-Dichloro-1,1-dimethyl-3-(4-chlorophenyl)propane acid and its dimethylamine salt are equivalent to 2.27% active ingredient.

KEEP FROM FREEZING
EPA Reg. No. 504-104
B. G. PRATT DIVISION
GABRIEL CHEMICALS, LTD. PATERSON, N.J. 07650



PRATT
SINCE 1904
TURF HERBICIDE 6000
CONTAINS TRIMEC * HERBICIDE 74 D,
N'OPP AND DICAMBA

**1 PINT COVERS
3000 Square Feet**

KILLS

- CHICKWEED
- DANDELION
- KNOTWEED
- POISON IVY
- GROUND IVY
- WILD ONION
- CLOVER
- SPURGE
- PLANTAINS
- HENBIT
- RAGWEED
- AND MOST OTHER BROADLEAF WEEDS

NET CONTENTS ONE GAL.

OTHER WEEDS CONTROLLED: BLACK MEDIC, BUCKHORN, CHICORY, LAMBQUARTER, DANDELION, DOG, GROUND IVY, HAZEL, HENBIT, LAMBQUARTER, LAVENDER, LINDSEY, MORNINGS GLORY, PEPERomia, PIGWEED, PLANTAINS, POISON IVY, POISON OAK, PUNNETT, RAGWEED, SHEPHERD'S PURSE, SPEEDWELL, SPURGE, WILD CARROT, WILD LUCI, WILD ONION, YARROW.

DIRECTIONS FOR USE:
WHERE TO USE: Turf Hericide 6000 may be used on lawns, Bermuda or Zoysia lawns, golf courses, and other areas where hard-turf grasses are desired. This product is especially effective on all types of broadleaf weeds such as HENBIT, LAMBQUARTER, SPOTTED SPURGE, DANDELION, BLACK MEDIC, and SHEPHERD'S PURSE. In addition to these, Turf Hericide 6000 kills many other weeds. A partial list is given above. Do not allow spray to get on vegetables or ornamental plants. Spot spray may result in some injury to be expected when used on these plants or on the grass lawn. Spot spray weeds only. Clover present will be damaged if sprayed. Retreat if necessary for hard to kill weeds.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat lawns, HENBIT, Wild Onion and Garlic in the fall in the spring, as needed. Do not reseed until the following year. Do not spray on lawns until the grass is 2 inches tall. Make a straight spray treatment. Spot spray weeds. Retreatment may be made at any time with a hand sprayer. Spray on the weeds at ground level. Wait 3 weeks before cutting lawns, mowing or stirring them in the bare spots. Do not use on lawns until spray is applied during drought or just before rain. Do not water lawn before treating. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE - ENTIRE LAWN:
MIXING AND SPRAYING: Use a hose end sprayer designed for lawns. Put recommended amount of chemical (see chart) in bottle and add water to indicated level. Spray evenly and spray on area indicated. Spray 2,500 sq. ft. with 10 oz. of 10-10-10 fertilizer and add water to the 10 oz. of herbicide. Mix with the fertilizer and spray as evenly as possible over the lawn.

Rate	Per 1000 Sq. Ft.	Per 5000 Sq. Ft.	Per 10000 Sq. Ft.
1/2 oz.	1/2	1	2
1 oz.	1	2	4
2 oz.	2	4	8
4 oz.	4	8	16
8 oz.	8	16	32

NOTE: Do not use on lawns where desirable plants are growing. Do not spray on lawns where desirable plants are growing. Do not spray on lawns where desirable plants are growing.

June 6, 1974
504-194