

**PRATT** CO.

# TURF HERBICIDE 10,000

ACCEPTED

THE FEDERAL BUREAU OF INVESTIGATION  
DEPARTMENT OF JUSTICE  
WASHINGTON, D. C. 20535  
RECEIVED 1/24/1955

## CONTROLS

- BEDSTRAW
- BLACK MEDIC
- BUCKHORN
- CHICORY
- CHICKWEED
- CLOVER
- DANDELION
- DOCK
- GROUND IVY
- HEAL-ALL
- HENBIT
- KNOTWEED
- LAMBSQUARTER
- LESPEDEZA
- MALLOW
- MORNING GLORY
- PEPPERGRASS
- PIGWEED
- PLANTAINS
- POISON IVY
- POISON OAK
- PURSLANE
- RAGWEED
- SHEPHERD'S PURSE
- SPEEDWELL
- SPURGE
- WILD CARROT
- WILD LETTUCE
- WILD ONION
- YARROW

## DIRECTIONS FOR USE

**HOW TO USE:** Turf Herbicide 10,000 may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, it controls most other broadleaf lawn weeds, including Dandelion and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

**WHEN TO USE:** Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

## HOW MUCH TO USE - ENTIRE LAWN

**HOSE-END SPRAYER:** Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example— to spray 1,700 sq. feet, measure 5 ozs. (10 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

**PRESSURE SPRAYERS:** Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated:

Col 1	Col 2	Col 3
AMT. Turf Herbicide 10,000 to use (Tbsp.) (Fl. Oz.)	Fill Bottle To (Gallons)	Area to Treat (Sq. Ft.)
1 1/2	1	170
2 1/2	2	340
5 2 1/2	5	850
10 5	10	1700
15 7 1/2	15	2550
20 10	20	3400
32 16	32	5440

AMT. Turf Herbicide 10,000 to Use (Tbsp.) (Fl. Oz.)	AMT. of water in sprayer (Gallons)	Area to be Sprayed (Sq. Ft.)
2 1/2	1	335
4 2	1	670
8 4	2	1340
12 6	3	2010

**NOTE:** Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

## HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required being certain to "wet" each weed.

**CAUTION:** Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

**CAUTION:** Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

**NOTICE:** Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

**KEEP OUT OF REACH OF CHILDREN**

**ACTIVE INGREDIENTS:**  
 \*\*\*\*  
 \*\*\*\*  
 \*\*\*\*  
**INERT INGREDIENTS** 78.94%

**Equipment:**  
 • 1.0% 2.5-D A-40 or 0.5% 2.5-D A-40  
 • 4.0% WDFP A-40 or 0.5% 2.5-D A-40  
 • 1.1% D-40 A-40 or 0.5% 2.5-D A-40

**CAUTION: Keep Out of Reach of Children**

See Back Panel for Additional Cautions

\*Trimes is a registered trademark of P.M.I. - Gordon Corporation. U. S. Patent No. 3,284,185

**NET CONTENTS GALLONS**  
**EPA REG. NO. 994-185**

**E. G. PRATT DIVISION GARNER CHEMICALS LTD.**  
**PATTERSON, N. J. 07650**

**PRATT**

**HERBICIDE**

**ACCEPTED**

JANUARY 30, 1955

THE FEDERAL INSECTICIDE AND RODENTICIDE ACT AND ECONOMIC POISON REGISTERED UNDER NO. 904-195

**CONTROLS**

- BEDSTRAW**
- BLACK MEDIC**
- BUCKHORN**
- CHICORY**
- CHICKWEED**
- CLOVER**
- DANDELION**
- DOCK**
- GROUND IVY**
- HEAL-ALL**
- HENBIT**
- KNOTWEED**
- LAMBSQUARTER**
- LESPEDEZA**
- MALLOW**
- MORNING-GLORY**
- PEPPERGRASS**
- PIGWEEED**
- PLANTAINS**
- POISON IVY**
- POISON OAK**
- PURSLANE**
- RAGWEED**
- SHEPHERD'S PURSE**
- SPEEDWELL**
- SPURGE**
- WILD CARROT**
- WILD LETTUCE**
- WILD ONION**
- YARROW**

**DIRECTIONS FOR USE**

**WHERE TO USE:** Turf Herbicide 10,000 may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, it controls most other broadleaf lawn weeds, including Dandelion and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover control will be damaged if sprayed.

**WHEN TO USE:** Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

**HOW MUCH TO USE - ENTIRE LAWN**

**HOSE-END SPRAYER:** Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example— to spray 1,700 sq. feet, measure 5 ozs. (10 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

**PRESSURE SPRAYERS:** Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated:

Col. 1	Col. 2	Col. 3
AMT. Turf Herbicide 10,000 to use: (Tbsp.) (Fl. Oz.)	Fl. Bottle To: (Gallons)	Area to Treat: (Sq. Ft.)
1 1/2	1	170
2 1	2	340
5 2 1/2	5	850
10 5	10	1700
15 7 1/2	15	2550
20 10	20	3400
32 16	32	5440

AMT. Turf Herbicide 10,000 to Use: (Tbsp.) (Fl. Oz.)	AMT. of water in sprayer: (Gallons)	Area to be Sprayed (Sq. Ft.)
2 1	1/2	335
4 2	1	670
8 4	2	1340
12 6	3	2010

**NOTE:** Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

**HOW MUCH TO USE: SPOT SPRAY**

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required being certain to "wet" each weed.

**CAUTION:** Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

**CAUTION:** Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

**NOTICE:** Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.