

7

SECRET

Diazinon^(K) Reg. trademark of CIBA-GEIGY Corp., Ardsley, New York.
CONTAINS 2 LBS. OF DIAZINON^(K) PER GALLON.

FILE NO. 72-PA-1

RECEIVED
JAN 10 1962
U.S. DEPT. OF JUSTICE
FEDERAL BUREAU OF INVESTIGATION
WASHINGTON, D.C.

1. *Chlorophyll a* (Chl *a*)

21.1.1

21.1.1

DO NOT ALLOWED: Call a Physician or Poison Control Center immediately.
DO NOT INDUCE VOMITING: Vomiting may cause a aspiration pneumonia.
NOT A PHYSICIAN: Do not use to aspiration hazard. Gastric lavage is indicated if material was taken orally. If 10% sodium chloride solution inhibition is present, intravenous injection of atropine sulfate is indicated. 2-4M is also antidote for atropine if administered in conjunction with atropine. If NaOH is used, dilute to fresh water. If aspiration is indicated, dilute NaOH to fresh water. Do not use material if it is fresh after contact with soap and water. Wash skin before removal. Wash with 10% NaOH. Wash eyes with plenty of water. If swallowed, do not induce vomiting.

THE UNIVERSITY OF CHICAGO PRESS

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

5 GALLON	14.99
10 GALLON	14.99
55 GALLON	14.99
1 GALLON	14.99

PRATT DIAZINON^(R) 25E INSECT SPRAY
EPA Reg. No. 904-201

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS- WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist or vapors. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep children and pets off treated areas until spray has dried. Do not use on humans or household pets. Utensils such as teaspoons and cups used to measure pesticides should not be used for food products. ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur or when weather conditions favor drift from treated areas. Cover bird baths and fish ponds. This product is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Do not apply when honey bees are actively visiting the crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next interval period of bee activity. PHYSICAL OR CHEMICAL HAZARDS: Do not use, pour or store near heat or open flame.

AN INSECTICIDE CONCENTRATE THAT MIXES WITH WATER TO MAKE A SPRAY THAT WILL CONTROL THE INSECTS AS LISTED. APPLY WHEN INSECTS APPEAR AND REPEAT AS NOTED IN DIRECTIONS, OR AS NECESSARY. FOR LAWNS AND VEGETABLE GARDENS, MEASURE AREA TO BE TREATED AND APPLY AT COVERAGE RATE.

1 FL. OZ. IN 3 GALS. OF WATER IS EQUIVALENT TO A 1 TO 400 DILUTION.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its label.

STORAGE AND DISPOSAL (8 Oz., 1 Pt., 1 Qt., 1 Gal.)

Store this product in its closed, original container in a locked area away from children and domestic animals. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. For Disposal, securely wrap original container in several layers of newspaper and discard in trash, or at an approved waste disposal facility.

STORAGE AND DISPOSAL (5 Gals., 30 Gals., 55 Gals.)

Do not contaminate water, food or feed by cleaning of equipment or disposal of waste. Store in the closed original container in a locked area away from children and domestic animals. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL (Metal): Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. FOR PLASTIC CONTAINERS: Follow above procedure or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. DO NOT REUSE CONTAINER.

PRAFT DIAZINON^(R) 25L INSECT SPRAY
EPA Reg. No. 904-201

DILUTION AND APPLICATION DIRECTIONS

LAWN SPRAY: To control Chinch Bugs, Sod Webworms, Ants, Brown Dog Ticks, and Earwigs, mix 1 fl. oz. of Diazinon^(R) 25E in 3 gals. of water to treat 125 sq. ft., or 8 fl. oz. in 24 gals. to cover 1,000 sq. ft. For Millipedes, use 1 pt. in 24 gals. to cover 1,000 sq. ft. Apply a few hours after a heavy rain or thorough watering for control of all listed lawn insects. For White Grubs (Japanese Beetle Larvae and European Chater) and Billbugs when insects or damage first appear. For Hyperodes Weevil (New York State) apply 6 fl. oz. in 10-30 gals. of water for 1,000 sq. ft. First treatment in mid-April, repeat in mid-May. For Digger Wasps, make application at dusk when insects are less active.

FLORAL, SHRUB AND SHADE TREE SPRAY: Mix 1 fl. oz. of Diazinon^(R) 25E in 3 gals. of water (8 fl. oz. in 24 gals.) to control Aphids, Mites, Leafhoppers, Thrips, Leafminers, and Scale Crawlers on Roses, Azaleas, Chrysanthemums, Gladioli, Carnations, Oaks, Maple, Locust, Holly, Boxwood, Pine, Spruce, Willow. Penetrate dense foliage to thoroughly cover all plant surfaces.

Do Not Use On Certain Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, Since Injury to The Plants May Occur.

DO NOT TREAT CROPS within the number of days of harvest shown in parentheses after each crop.

FRUIT SPRAY: Mix 8 fl. oz. of Diazinon^(R) 25E in 24 gals. of water to control Codling Moths, Mites, Aphids, Leafrollers, Scale Crawlers on Apples (14), Peaches (21), Plums (10), Cherries (10), and Strawberries (5), Grapes (7). If sticker is used on Grapes (18), Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Apply as a full cover spray.

VEGETABLE SPRAY: Mix 2 tablespoonful or 1 fl. oz. of Diazinon^(R) 25E in 3 gals. of water (8 fl. oz. in 24 gals.). Three gals. of diluted spray will cover 1,000 sq. ft. Make full coverage applications at 7 to 10 day intervals if needed starting when insects first appear. Used to control Bean Beetles, Aphids, Leafminers, Imported Cabbageworms and Cucumber Beetles. For use on Tomatoes (1), Carrots (10), Potatoes (35), Beans (7), Cucumbers (7), Onions (10), Cantaloupes (3), Peppers (5), and Cabbage (7). For cutworms (Surface and Subterranean), Wireworms, Garden Symphylans and Root Maggots, apply 15 fl. oz. in sufficient water to cover 1,000 sq. ft. Make applications to soil surface as a broadcast spray prior to planting Cabbage, Carrots, Lettuce, Peas, Radishes, Snapbeans or Tomatoes. Thoroughly mix spray into the top 4-8 inches of soil.

OUTDOOR AREA NUISANCE PESTS: Mix 8 fl. oz. of Diazinon^(R) 25E in 1.0 gals. of water. Thoroughly spray window frames, doors, sills, screens, porches, patio walls, foundations, walks, garbage cans and into cracks and other places where insects may hide to control Flies, Ants, Cockroaches, Boxelder Bugs, Clover Mites, Spiders, Brown Dog Ticks, Chiggers and Earwigs. Do not treat plants with this dosage rate.

LIMITED WARRANTY: Buyer Agrees All Uses of this Material are in Accordance with Directions Given Herein.