

For Cutworms (Surface and Subterranean), Wireworms, Garden Symphians and Root Maggots, apply 15 fl. oz. in sufficient water to cover 500 sq. ft. Make application to soil surface as a broadcast spray prior to planting Cabbage, Carrots, Lettuce, Peas, Radishes, Snapbeans or Tomatoes. Thoroughly mix spray into the top 4-8 inches of soil.

OUTDOOR AREA NUISANCE PESTS: Mix 8 fl. oz. of Diazinon 12.5E in 1.5 gals. of water (1 pt. to 3 gals.) Thoroughly spray window frames, doors, sills, screens, porches, patio walls, foundations, walks, garbage cans and into cracks and other places where insects may hide to control Flies, Ants, Cockroaches, Boxelder Bugs, Clover Mites, Spiders, Ticks, Chiggers and Earwigs. Do not treat plants with this dosage rate.

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go on grass or other treated areas until spray has completely dried. Keep out of reach of domestic animals. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to the plants may occur.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS HEREWITH.

B. G. PRATT DIVISION
GABRIEL CHEMICALS LTD., PATERSON, N.J. 07509



CONTAINS 1 POUND DIAZINON® PER GALLON

FOR LAWNS, FLOWERS, SHRUBS, SHADE TREES AND CERTAIN FRUITS AND VEGETABLES as listed on back panel.

ALSO CONTROLS CLOVER MITES, EARWIGS, MILLIPEDES, ANTS

ACTIVE INGREDIENTS:	By Weight
O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	12.5%
Aromatic Petroleum Derivative Solvent	73.0%
INERT INGREDIENTS	14.5%
E.P.A. Reg. No. 904-205	TOTAL.....100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional warnings.

AN INSECTICIDE CONCENTRATE THAT MIXES WITH WATER TO MAKE A SPRAY THAT WILL CONTROL THE INSECTS AS LISTED. APPLY WHEN INSECTS APPEAR AND REPEAT AS NOTED IN DIRECTIONS, OR AS NECESSARY. FOR LAWNS AND VEGETABLE GARDENS, MEASURE AREA TO BE TREATED AND APPLY AT COVERAGE RATE.

DIRECTIONS FOR USE

LAWN SPRAY: To control Chinch Bugs, Sod Webworms, Ants, Ticks and Earwigs, mix 2 fl. oz. of Diazinon 12.5E in 3 gals. of water to treat 125 sq. ft. or 1 pt. in 24 gals. to cover 1000 sq. ft. For Millipedes use 1 qt. in 24 gals. for 1000 sq. ft. Apply a few hours after a heavy rain or thorough watering for control of all listed lawn insects. For White Grubs (Japanese Beetle Larvae and European Chafer) and Billbugs, apply 16 fl. oz. mixed in 5-15 gals. of water for 1000 sq. ft. For Grubs apply between late July and early October. Water thoroughly after application. Apply for Billbugs when insects or damage first appear. For Hyperodes Weevil (New York State) apply 12 fl. oz. in 5-15 gals. of water for 500 sq. ft. First treatment in mid-April, repeat in mid-May.

FLORAL, SHRUB AND SHADE TREE SPRAY: Mix 2 fl. oz. of Diazinon 12.5E in 3 gals. of water (1 pt. to 25 gals.) to control Aphids, Mites, Leafhoppers, Thrips, Leafminers and Scale Crawlers on Roses, Azaleas, Chrysanthemums, Gladioli, Carnations, Oaks, Maple, Locust, Holly, Boxwood, Pine, Spruce, Willow. Penetrate dense foliage to thoroughly cover all plant surfaces.

FRUIT SPRAY: Mix 1 pt. of Diazinon 12.5E in 24 gals. of water to control Codling Moths, Mites, Aphids, Leafrollers and Scale Crawlers on Apples, Peaches, Plums, Grapes, Cherries and Strawberries. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of picking.

VEGETABLE SPRAY: Mix 4 tablespoons or 2 fl. oz. of Diazinon 12.5E in 3 gals. of water (1 pt. in 25 gals.) Three gallons of diluted spray will cover 1000 sq. ft. of Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cantaloupes, Peppers and Cabbage to control Bean Beetles, Aphids, Leafminers, Imported Cabbageworms and Cucumber Beetles. Make applications at 7-10 day intervals when insects appear. Do not spray within 14 days of picking. Tomatoes may be sprayed 1 day before picking. Wash home garden vegetables before using.

ACCEPTED

APR 23 1973

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 709-265