

LAWN INSECT CONTROL

To control insects listed below, apply recommended rates. For insects such as Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs and Collembola (Springtails), which are commonly found near the house and may enter the house, spray a 5 ft. band of soil around the house, as well as the house foundation wall to a height of 2-3 ft. Repeat application if necessary.

Insects	Rate/1,000 sq. ft.		Remarks
	Diazinon AG4E	Water	
Lawn Chinch Bugs	2-3 fl. oz. or 3-6 fl. oz.	25 gals.	Use higher rate for longer residual control and in lawns of dense growth, such as St. Augustine lawns.
Ants, Armyworms, Clover Mites, Collembola (Spring-tails), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Sod Webworms (Lawn Moths), Sowbugs	4 fl. oz.	3 gals.	Spot spray ant hills and wasp ground nest openings. For lawn billbugs, irrigate the lawn thoroughly immediately after treatment. For frit flies, mow grass and water well before treatment but delay application until grass is dry. Do not water again until necessary.
Brown Dog Ticks, Bermuda Mites, Chiggers, Fleas, Leafhoppers, Ticks	1 1/4 fl. oz.	3 gals.	Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Millipedes	8 fl. oz.	3 gals.	
Rhodes-Grass Scales	5 fl. oz.	25 gals.	Apply when crawlers first emerge.

Billbugs and White Grubs (Japanese Beetle Larvae, European Chafer and Southern Chafer): Apply 4 fl. oz. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Hyperodes Weevil (New York only): Apply 3 fl. oz. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Boxelder Bugs: Spray bugs outside the home with 1 1/4-2 1/2 fl. oz. (2 1/2-5 tablespoons) in 1 gal. of water. At the same time, thoroughly spray around foundation, windows, doors, tree trunks and into cracks and other places where the insects hide.

FOREST INSECTS

PINE (SEED ORCHARDS, PLANTATIONS, FORESTS): Arkansas Sawfly (Loblolly Sawfly): Apply 1 pt. per acre over the tree canopy in sufficient water to assure thorough coverage. Application should be made at or near the complete egg hatch. Aerial applications should be made with a minimum of 2 gals. of spray per acre, in swaths not more than 90 ft. wide. Repeat application as needed.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

B. G. PRATT DIVISION

GABRIEL CHEMICALS LTD., PATERSON, N.J. 07503



DIAZINON AG4E
FOR TURF AND TREE

FOR CONTROL OF INSECTS ON SHADE TREES, ORNAMENTALS AND TURF
as listed on the label

DIAZINON AG4E CONTAINS 4 LBS. DIAZINON PER GALLON.

ACTIVE INGREDIENTS:	By Weight
0, 0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	48%
Xylene	36%
INERT INGREDIENTS	16%
TOTAL	100%

DIAZINON®—Trademark of CIBA GEIGY CORP., Ardsley, N.Y.

E.P.A. Reg. No. 904-206

WARNING: KEEP OUT OF REACH OF CHILDREN

See side panel for complete caution statement.

NET CONTENTS

ACCEPTED WITH COMMENTS

ACCEPTED
APR 4, 1972

PRATT DIAZINON AG4E IS AN INSECTICIDE TO BE MIXED WITH WATER ACCORDING TO DIRECTIONS FOR CONTROL OF INSECT PESTS ON

DIRECTOR

ORNAME

To control certain insects on Boxwood, Camellia, Carnation, Hawthorn, Holly, Juniper, Lilac, Magnolia, Rose, Spruce and Yew.

Insect

Aphids, Bagworms, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, European Red Mites, Holly Bud Moths, Leafhoppers, Omnivorous Leaf Hoppers, Privet Mites, Cottony-cushion, Lecanium, Pinescale, Soft Scale, Inrips, Two-spotted Whiteflies

Apple-and-Thorn Skeletonizers, Fall Webworms, Hemlock Cherry Borer, Oblique-banded Leaf Roller, Pines Caterpillars

1 fl. oz. = 2 tablespoons

Do not use on certain Ferns, since injury to the plants may result. For Mimosid Webworms: Mix 1/2 fl. oz. water, and apply as a thorough second and third application.

WARNING: KEEP

May be fatal if swallowed. May irritate eyes. Do not get in eyes. Avoid contact with skin and clothing. Avoid contamination of animals. Do not use on human children or pets to go on to

This product is toxic to fish. Do not apply to lakes or streams. See label for complete directions on this label. Do not apply to trees.

This product is highly toxic to shade trees, ornamentals and shrubs. Do not apply to trees from your Cooperative Agricultural

Do not reuse empty container. Discard in a safe place.

MINIMUM FLASH POINT: 82°F OR OPEN FLAME



DIAZINON AG4E FOR TURF AND TREE

FOR CONTROL OF INSECTS ON SHADE TREES, ORNAMENTALS AND TURF

as listed on the label

DIAZINON AG4E CONTAINS 4 LBS. DIAZINON PER GALLON.

ACTIVE INGREDIENTS:	By Weight
O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	48%
Xylene	36%
INERT INGREDIENTS	16%
TOTAL	100%

DIAZINON®—Trademark of CIBA-GEIGY CORP., Ardsley, N. Y.

E.P.A. Reg. No. 904-206

WARNING: KEEP OUT OF REACH OF CHILDREN

See side panel for complete caution statement.

NET CONTENTS

PRATT DIAZINON AG4E IS AN EMULSIFIABLE SOLUTION WHICH, WHEN DILUTED WITH WATER ACCORDING TO THE DIRECTIONS BELOW, IS INTENDED FOR THE CONTROL OF INSECT PESTS ON SHADE TREES, ORNAMENTALS AND TURF.

DIRECTIONS FOR USE

ORNAMENTAL INSECT CONTROL

To control certain insects on ornamentals such as Arbervitae, Azalea, Birch, Camellia, Gardenia, Chrysanthemum, Douglas Fir, Elm, Gladiolus, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Plum, Poplar, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated below.

Insects	Amount Diazinon AG4E to Use	
	per 3 gals. water	per 100 gals. water
<u>Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leaf Miners, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Mites, Leafhoppers, Obscure Root Weevils, Omnivorous Leaf Eaters, Privet Mites, Scale Crawlers, Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft Scale, Tetranychus, Two-spotted Mites, Webworms, Whiteflies</u>	1/2 fl. oz.	1 pt.
<u>Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Chermes, Oak Looper, Orange-banded Leaf Rollers, Pear Slugs, Tent Caterpillars</u>	1 1/2 fl. oz.	3 pts.

1 fl. oz. = 2 tablespoons 16 fl. oz. = 1 pint

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to the plants may occur.

Minors Webworms: Mix 1/2 fl. oz. in 3 gals. of water for 1 pt. in 100 gals. of water, and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

WARNING: KEEP OUT OF REACH OF CHILDREN.

May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not permit children or pets to go on to sprayed grass until spray has completely dried.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be poisoned. Keep out of lakes, streams or ponds. Apply this product only as directed on this label. Do not apply when weather conditions favor drift from treated area. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on shade trees, ornamentals and turf. Protective information may be obtained from our Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

MINIMUM FLASH POINT: 92 F. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

insects such as
bees, Millipedes,
found near the
and the house, as
that application if

marks

for longer
and in lawns
th, such as
lawns.

t hills and wasp
penings.

ugs, irrigate the
ly immediately
nt.

ow grass and
ore treatment
lication until;
Do not water
cessary.

to animals.

ticks, spray
er shrubbery,
ear house.

awlers first

an Chafer and
ater grass thorny
time between
activity is first
ng damage from

sq. ft. of grass.
in mid-April and

oz. (2 1/2-5 table-
/ around founda-
places where the

ly (Loblolly Saw-
water to assure
he complete egg
2 gals. of spray
ation as needed.

THIS MATERIAL
ITH.

ACCEPTED

APR 4 1973

ACCEPTED