

904-208

07/08/1973

1/2

**VEGETABLE GARDEN SPRAY:** Mix 1 fl. oz. in 3 gals. of water to 650 sq. ft. (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7-10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with directions.

**BEANS (Green Lima and Snap):** Japanese and Mexican Bean Beetle, Tarnished Plant Bug; **TOMATO, EGGPLANT and PEPPER:** Leafhopper, Lace Bug, Tomato Fruit Worm; **PUMPKIN, SQUASH, CUCUMBER, MELON:** Flea Beetle, Melonworms, Cucumber Beetle, Squash Bug. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

**CAUTION: HARMFUL IF SWALLOWED OR INHALED.** Do not breathe spray mist or take internally. Avoid contact with skin, wear gloves and long-sleeved work clothing. Change to clean clothing daily. Wash hands and wash face before eating or smoking. Wash thoroughly after handling. Do not contaminate food or feed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. For protection of bees, avoid use during periods when honey bees are visiting crops. When necessary to use during such periods, warn bee keepers to locate hives at a safe distance until one week after application. Further protective information may be obtained from your Cooperative Agricultural Extension Service. Do not allow drifting to adjoining food or forage crops. Do not use in any manner not described on this label.

Do not reuse container. Wrap in newspaper and place in trash can for safe disposal or bury in a safe place away from domestic water supplies.

**BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.**

**PRATT**

**LIQUID 2 SEVIN®**  
Plant Insecticide

**Kills CHINCH-BUGS;  
JAPANESE BEETLES;  
MOSQUITOES**  
and certain other  
**LAWN, GARDEN &  
ORNAMENTAL INSECTS**  
contains 2 lbs. of  
Sevin per gallon

**CAUTION: Keep Out of Reach of Children.**

**PROTECT FROM TEMPERATURES BELOW 32°F.**  
Keep cover on opened container to prevent drying.

**DIRECTIONS:** Liquid 2 Sevin is an extremely fine dispersion of Sevin in water. **SHAKE WELL BEFORE USING.** Pour recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.

**LAWNS:** For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Sod Webworms (Lawn Moths) and Millipedes, apply 8 fl. oz. in 15-20 gals. of water evenly over each 500 sq. ft. of established lawn area. For best results, mow the lawn and apply immediately after rain or watering. Repeat in 2-3 weeks if necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES:** For use on Roses, Carnations, Gladioli, Zinnias, Chrysanthemums, Lilacs, Arbor Vites, Junipers, Hydrangeas, Azaleas, Elms, Maples, Oaks, Dogwood, Birch and Pines (DO NOT USE ON BOSTON IVY), use 4 teaspoonsful per gal. of water or 1 pt. to 25 gals. (1 to 200 dilution) to control Apple Aphid, Bagworm, Birchleaf miner, Blister Beetles, Boxelder Bugs, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, June Beetle, Lace Bug, Leafhoppers, Leafrollers, Mealybugs, Mirids, Webworms, Oakleaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Puss Caterpillars, Psyllids, Rose Aphids, Rose Slug, Scale Insects, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Willowleaf Beetles. Apply when insects first appear. Repeat weekly or as necessary throughout the season.

**ADULT MOSQUITO CONTROL** for Home Grounds, Parks and Recreation Areas, use 1 to 200 dilution in hydraulic low pressure sprayers (4 teaspoonsful per gal. or 1 pt. to 25 gals. of water). Spray to thoroughly cover low shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate (DO NOT APPLY TO WATER). Repeat at 7-10 day intervals or as necessary. (DO NOT SPRAY BOSTON IVY).

2/2

ACCEPTED  
JUL 8 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

904-208

RECEIVED  
JUL 10 1973