

BUNDLES OF NURSERY PLANTING STOCK SHOULD BE PERMITTED TO DRY THOROUGHLY BEFORE PLANTING OR STORING.

BRUSH APPLICATION: Apply on lower trunk of fruit trees or ornamental plantings from ground level to a height of approximately 2½ ft. to protect plants from injury to trunk.

PRATT'S ANIMAL REPELLENT should be applied only when surfaces of plants to be treated are dry and application should be made under favorable drying conditions, preferably during periods of moderate temperature and light winds, to allow the repellent film to form and set properly. Spraying under favorable drying conditions is also essential to avoid excessive run-off of the diluted repellent.

BUYER assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION

Harmful if swallowed. May cause skin irritation. Do not inhale spray mist. Do not get in eyes or on skin. Wash thoroughly after handling. Do not use on plant parts that are to be used for food or feed.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Flush thoroughly to clean application equipment immediately after using. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Stir before using and keep container tightly closed to prevent evaporation. Avoid storage at freezing temperature. If frozen, thaw completely and stir well before using.

E.P.A. Reg. No. 904-210

Manufactured for

R. G. PRATT DIVISION

GABRIEL CHEMICALS LTD., PATERSON, N.J. 07509

NET CONTENTS: 1 LIQUID QUART

ON FRUIT TREES USE DORMANT APPLICATION ONLY

PRATT'S ANIMAL REPELLENT is a weather resistant contact repellent containing 20% Thiram. It is a specially formulated stable suspension designed for use on forest nursery planting stock, fruit trees, and ornamental plantings and nursery stock for protection against rabbits, meadow mice, and, to a lesser extent, deer. It may be applied during the dormant or growing season. When applied as directed, a single application will protect treated surfaces of plants from damage up to six months or longer, depending upon weather and other conditions.

DIRECTIONS FOR USE

Dilute **PRATT'S ANIMAL REPELLENT** with equal parts water. A 1:1 dilution contains 10% Thiram. Mix thoroughly before applying and mix only the quantity to be used. Diluted material should be used immediately. The diluted repellent may be applied as a spray, dip or brush application.

SPRAY APPLICATION Apply as a full-coverage spray on forest nursery planting stock, ornamental plantings or other nursery stock. The spray can be applied with a hand sprayer or mechanical sprayer. Apply on forest nursery planting stock prior to planting. Apply on fruit trees, ornamentals and other nursery stock in late summer or early Fall, or when rodent feeding damage is first noticed. In the case of tree seedlings, it is suggested that not more than 3,000 seedlings be treated with each gallon of finished spray.

DIP APPLICATION Apply to bundles of forest nursery or fruit trees planting stock by dipping to thoroughly wet the foliage and other parts to be protected. Excess repellent should be permitted to run off treated plants without coming in contact with the roots.

EMPTIED

911018

BUNDLES OF NURSERY PLANTING STOCK SHOULD BE PERMITTED TO DRY THOROUGHLY BEFORE PLANTING OR STORING.

BRUSH APPLICATION: Apply on lower trunk of fruit trees or ornamental plantings from ground level to a height of approximately 2½ ft. to protect plants from injury to trunk.

PRATT'S ANIMAL REPELLENT should be applied only when surfaces of plants to be treated are dry and application should be made under favorable drying conditions, preferably during periods of moderate temperature and light winds, to allow the repellent film to form and set properly. Spraying under favorable drying conditions is also essential to avoid excessive run-off of the diluted repellent.

BUYER assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION

Harmful if swallowed. May cause skin irritation. Do not inhale spray mist. Do not get in eyes or on skin. Wash thoroughly after handling. Do not use on plant parts that are to be used for food or feed.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Flush thoroughly to clean application equipment immediately after using. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Stir before using and keep container tightly closed to prevent evaporation. Avoid storage at freezing temperature. If frozen, thaw completely and stir well before using.

PRATT

PROTECTS
NURSERY STOCK, FRUIT TREES
AND ORNAMENTALS
AGAINST
feeding damage of
RABBITS
DEER
MEADOW MICE



STIR OR SHAKE WELL BEFORE USING

ON FRUIT TREES USE DORMANT APPLICATION ONLY

PRATT'S ANIMAL REPELLENT is a weather resistant contact repellent containing 20% Thiram. It is a specially formulated stable suspension designed for use on forest nursery planting stock, fruit trees, and ornamental plantings and nursery stock for protection against rabbits, meadow mice, and, to a lesser extent, deer. It may be applied during the dormant or growing season. When applied as directed, a single application will protect treated surfaces of plants from damage up to six months or longer, depending upon weather and other conditions.

DIRECTIONS FOR USE

Dilute PRATT'S ANIMAL REPELLENT with equal parts water. A 1:1 dilution contains 10% Thiram. Mix thoroughly before applying and mix only the quantity to be used. Diluted material should be used immediately. The diluted repellent may be applied as a spray, dip or brush application.

SPRAY APPLICATION: Apply as a full-coverage spray on forest nursery planting stock, ornamental plantings or other nursery stock. The spray can be applied with a hand sprayer or mechanical sprayer. Apply on forest nursery planting stock prior to planting. Apply on fruit trees, ornamentals and other nursery stock in late summer or early Fall, or when rodent feeding damage is first noticed. In the case of tree seedlings, it is suggested that not more than 3,000 seedlings be treated with each gallon of finished spray.

DIP APPLICATION: Apply to bundles of forest nursery or fruit trees planting stock by dipping to thoroughly wet the foliage and other parts to be protected. Excess repellent should be permitted to run off treated plants without coming in contact with the roots.

ACCEPTED

5/26/73

RECEIVED
MAY 26 1973
FEDERAL BUREAU OF INVESTIGATION
U.S. DEPARTMENT OF JUSTICE