

B  
G



SINCE 1904  
TM

# METHOXY-DIAZINON® 20-10 E.C.

For control of insects on alfalfa, clover, vegetables & small fruits.

<b>ACTIVE INGREDIENTS:</b>	
Methoxychlor Technical*	20%
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate**	10%
Xylene	63%
<b>INERT INGREDIENTS</b>	<b>7%</b>
<b>TOTAL</b>	<b>100%</b>

\*Equivalent to 17.6% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.4% other isomers and reaction products.

\*\*Diazinon® - Trademark of CIBA-GEIGY Corp.  
EPA Reg. No. 904-221

### CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. May be absorbed through skin. Wash thoroughly after handling. Avoid contamination of food and feed. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.  
DO NOT RE-USE EMPTY CONTAINER. Destroy by perforating or crushing and burying in a safe place.  
Store at temperatures above 32°F. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NET CONTENTS     GALLONS  
1 5 30 55

B. G. PRATT DIVISION, Gabriel Chemicals Ltd.  
Paterson, N. J. 07509

EPA EST. 904-NJ-1

PROD. NO. 7100

Pratt Methoxy-Diazinon 20-10 E. C. is an emulsifiable solution which is to be diluted with water. When ground applications are to be made, dilute with water according to the Directions for Use section. For application by aircraft, refer to the Aerial Application section of the label.  
Pratt Methoxy-Diazinon 20-10 E. C. is intended for use against insect pests of agricultural crops and should not be used in barns or dwellings. Applications should be repeated as necessary to maintain control.

### DIRECTIONS FOR USE APPLY ONLY AS SPECIFIED ON THIS LABEL

#### FORAGE INSECTS

**ALFALFA AND CLOVER (PURE STANDS AND STANDS CONTAINING GRASS):** Alfalfa Weevil Larvae, Egyptian Alfalfa Weevils, Pea Aphids, Yellow Clover Aphids, Spotted Alfalfa Aphids, Leafhoppers, Plant Bug Nymphs, Mites: Apply 2-3 qts. per acre in sufficient water to achieve thorough coverage. Apply when 30-50% of the growing tips show feeding damage. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.  
**Alfalfa Caterpillars, Armyworms, Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Lygus Bugs, Spittlebugs:** Apply 2½-3 qts. per acre in sufficient water to achieve thorough coverage when insects first appear. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.

**NOTE:** Dairy and beef cattle and sheep 1) may be grazed 7 days following treatment, 2) may be fed green forage or hay provided alfalfa or clover is not cut for 7 days following treatment. Do not spray livestock directly with either air or ground equipment.

#### SMALL FRUIT INSECTS

For the control of the insects on the following crops, apply 2½ qts. per 100 gals. of water per acre as a thorough cover spray when insects first appear. Repeat applications at 7-10 day intervals as infestations warrant.

**CRANBERRIES:** Cranberry Fruitworms, Leafhoppers, Spittlebugs: Do not apply within 14 days of harvest.

**GRAPES:** Flea Beetles, Grape Berry Moths, Grape Leaf Folders, Grape Leaf Skeletonizers, Japanese Beetles, Leafhoppers, Pacific Mites, Rose Chafers: For Grape Berry Moths, apply as a post-bloom cover spray and repeat 10 days later; make third and fourth applications at 10-day intervals if necessary. Do not apply within 14 days of harvest.

**STRAWBERRIES:** Aphids, Flea Beetles, Spittlebugs, Strawberry Leaf Beetles, Strawberry Leaf Rollers, Strawberry Weevils, Two-Spotted Mites: Do not apply within 5 days of harvest.

#### VEGETABLE INSECTS

For the control of the insects on the following crops, use 2½ qts. per acre in sufficient water for thorough coverage. Begin treatment at first sign of insect infestation and repeat at 7-10 day intervals as infestation warrants.

**BROCCOLI:** Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles: Do not apply within 14 days of harvest.

**CABBAGE:** Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles: Do not apply within 7 days of harvest.

**CAULIFLOWER:** Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles: Do not apply within 7 days of harvest.

**CARROTS:** Aphids, Leafhoppers: Do not apply within 10 days of harvest or 14 days if tops are used for food or feed.

**COLLARDS:** Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, Harlequin Cabbage Bugs, Imported Cabbageworms: Do not apply within 14 days of harvest.

**CORN:** Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Rose Chafers: Do not apply within 7 days of harvest.

**CUCUMBERS:** Aphids, Cucumber Beetles, Serpentine Leaf Miners, Thrips: Do not apply within 7 days of harvest.

**KALE:** Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, Harlequin Cabbage Bugs: Do not apply within 14 days of harvest.

**LETTUCE:** Aphids, Dipterous Leaf Miners, Leafhoppers: Do not apply within 14 days of harvest.

**MELONS (Cantaloupes, Muskmelons, Watermelon):** Aphids, Cucumber Beetles, Leafhoppers, Melon Worms, Mites, Serpentine Leaf Miners, Thrips: Do not apply within 3 days of harvest.

**PEAS:** Aphids, Pea Weevils: Do not apply within 7 days of harvest or grazing.

**POTATOES:** Aphids, Banded Cucumber Beetles, Blister Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Flea Beetles, Leafhoppers, Southern Armyworms: Do not apply within 35 days of harvest.

**RADISHES:** Aphids, Dipterous Leaf Miners, Flea Beetles: Do not apply within 10 days of harvest.

**SQUASH (Summer and Winter):** Aphids, Cucumber Beetles, Leafhoppers, Mites, Squash Vine Borers: Do not apply to winter squash within 3 days, or summer squash within 10 days, of harvest.

**TOMATOES:** Aphids, Blister Beetles, Dipterous Leaf Miners, Fall Armyworms, Flea Beetles, Japanese Beetles, Tomato Fruitworms: Do not apply within 1 day of harvest.

**TURNIPS:** Aphids, Dipterous Leaf Miners, Flea Beetles: Do not apply within 10 days of harvest or 14 days if tops are to be used as food or feed.

#### AERIAL APPLICATION

Pratt Methoxy-Diazinon 20-10 E. C. may be effectively applied by aircraft. Aerial applications may be made on those crops where such applications are practical or are accepted practice. Refer to the specific crop and use under the Directions for Use section for the rate and timing of application. Mix the recommended amount of Pratt Methoxy-Diazinon 20-10 E. C. in sufficient water to apply a minimum of 2 gallons of spray per acre.

Observe the Caution statement and all precautions on the label to avoid any effects to application personnel or crops. Flagmen and loaders should wash thoroughly after handling, before eating and at the end of each day's operation.

Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Do not blow spray lines or dump contents while in flight.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

274-LF-7100-3-5/74

904-221

8-1-1974

1/2

2/2

5  
2  
2

**ACCEPTED**  
**AUG 1 1974**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR RECORDING POISON REGISTER  
ED UNDER NO. 904-227