

BEANS (SNAP) - White Rot (Sclerotinia), Gray Mold (Botrytis). Apply 1 1/2 - 2 pounds per acre (2 1/2 - 5 level tablespoons) in 5 gallons water applied to 1000 sq. ft.), at 25% to 50% bloom; repeat at peak bloom. Do not apply within 14 days of harvest; do not graze or feed treated bean vines or hay to livestock.

BULBS (Easter Lily, Tulip, Gladiolus, Daffodil, Iris) - Fusarium and Penicillium Rots: Use 1 1/2 pounds per 100 gallons water (2 tablespoonsful per gallon). Soak cleaned bulbs for 15-30 minutes in warm dip (80° to 85°F), preferably within 48 hours after digging. Dry bulbs after treatment. If bulbs are for forcing, treat after bulbs have been heat-cured.

IMPORTANT: Never allow Benomyl 50W to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of Benomyl 50W as a fungicide. Keep container closed when not in use.

CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Keep away from fire or sparks. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

Do not re-use container. Wrap empty container in newspaper and place it in trash can for safe disposal.

ENVIRONMENTAL CAUTION

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Use this material only in the manner described on this label.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.


Packed by:
B.G. PRATT DIV., GABRIEL CHEMICALS LTD.
Paterson, N.J. 07509

EPA Reg. No. 904-224

EPA Est. 904-NJ-1

Prod. No. 7302

B
·
G



PRATT

SINCE 1904
TM

BENOMYL 50W SYSTEMIC FUNGICIDE

Provides Curative (Eradicant) and
Protective Action for certain diseases of

- TURF • FLOWERS
 - ROSES • ORNAMENTALS
 - BULBS • VEGETABLES
- as listed

ACTIVE INGREDIENT:	BY WEIGHT
Benomyl (Methyl 1-(butylcarbamoyl)- 2-benzimidazolecarbamate).....	50%
INERT INGREDIENTS:.....	50%
TOTAL.....	100%

U.S. Pat. No. 3,631,176 -
E.I. DuPont DeNemours & Company

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Cautions

NET WEIGHT 2 OZ. AV.


Benomyl 50W is re flowers, ornamental fungus diseases, p action. Benomyl 50 application as a spr sprayer tank and ag spraying to assure

FLOWERS, ORNAM
Mold (Botrytis): Use when disease first a

TURF - Dollar spot (Solani), Fusarium P tablespoonsful) per coverage, usually 2- disease and contin conditions are unusu to 5-7 days. Do not

ROSES (Field and tablespoonsful per appears and repeat shorten intervals du

CUCUMBERS, MEL
Blight. Powdery Mi teaspoonsful in 5 g when plants begin to intervals as needed, higher rate and app



BENOMYL 50W SYSTEMIC FUNGICIDE

Provides Curative (Eradicant) and
Protective Action for certain diseases of

- TURF • FLOWERS
 - ROSES • ORNAMENTALS
 - BULBS • VEGETABLES
- as listed

INGREDIENT:	BY WEIGHT
Benomyl (Methyl 1-(butylcarbamoyl)- 1H-imidazolecarbamate)	50%
INERT INGREDIENTS:	50%
.....	100%

No. 3,631,176 -
Mont DeNemours & Company

CAUTION:
KEEP OUT OF REACH OF CHILDREN
See Inside Panel for Additional Cautions
NET WEIGHT 2 OZ. AV.

DIRECTIONS

Benomyl 50W is recommended for the control of certain diseases of roses, flowers, ornamentals and turf. It is highly effective for the control of many fungus diseases, providing both curative (eradicant) action and protective action. Benomyl 50W is a wettable powder to be mixed with water for application as a spray. Add recommended quantity of fungicide to half-filled sprayer tank and agitate. Then add balance of water. Agitate mixture while spraying to assure uniformity.

FLOWERS, ORNAMENTALS (Field and Greenhouse) - Powdery Mildew, Gray Mold (Botrytis): Use one tablespoonful per 2 gallons water. Begin application when disease first appears and repeat at 10-14 day intervals as long as needed.

TURF - Dollar spot (Sclerotinia Homeocarpa), Large Brown Patch (Rhizoctonia Solani), Fusarium Patch (Fusarium Nivale): Apply at the rate of 2 ounces (12 tablespoonful) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2-5 gallons per 1000 sq. ft. Apply at earliest appearance of disease and continue at 10-14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5-7 days. Do not graze or feed clippings from treated areas to livestock.

ROSES (Field and Greenhouse) - Powdery Mildew, Black Spot: Use one tablespoonful per 2 gallons water. Begin application when disease first appears and repeat at 10-14 day intervals throughout the growing season; shorten intervals during humid, rainy weather.

CUCUMBERS, MELONS, SUMMER and WINTER SQUASH - Gummy Stem Blight, Powdery Mildew, Anthracnose: Apply 4-8 ounces per acre (2 level teaspoonful in 5 gallons water applied to 1000 sq. ft.). Begin applications when plants begin to run or when disease first appears, and repeat at 7-14 day intervals as needed. Under conditions of severe disease pressure use the higher rate and apply at 7-day intervals or as needed.

2-13-75

104-224

CAUTION

Harmful if swallowed, inhaled or absorbed through skin or eyes. Do not swallow, inhale or allow contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate food or feed.

Toxic to fish and wildlife, do not contaminate streams, lakes or ponds with this product.

KEEP CONTAINER CLOSED.

DESTROY CONTAINER WHEN EMPTY

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and or damage, resulting from its misuse as such, or in combination with other products.



CREOSOTE A. W. P. A.

COMPOSITION

ACTIVE INGREDIENTS:

Coal Tar Creosote 97.0%

INERT INGREDIENTS:

3.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Caution Statement to Left

NET CONTENTS 1 QUART

PACKAGED BY

HTTTB971

DIRECTIONS

LACCO CREOSOTE A.W.P.A. is prepared for use against certain pests of timber as listed below. This product meets the specifications of the American Wood Preservers Association for No. 1 Creosote; California Department of Agriculture, Economic Poisons Regulation No. 1; and complies with specifications contained in Sec. 106, Building Ordinance No. 287,000, City of Los Angeles, Department of Building and Safety.

Meets Federal Specification TT-C-655

This product is not adulterated and will congeal during cold weather. Congealing may be prevented by storing in a warm place until contents are liquid. It is illegal to open this container for resale of contents. Contents must be sold in original, unopened and labeled container. L.A.F.D. CLASSIFICATION F-2.

TERMITES, WOOD DECAY: Fenceposts, mudsills, bridge timber, telephone poles and other timber which will be in contact with the soil. Immerse dry, barked and well seasoned timber in this product full strength for 12-48 hours.

Treat sufficient length so that when placed in the soil at least 6 inches of treated timber will be above the soil. Structural timber, shingles, fence rails and other timber which will not be in contact with the soil. Immerse dry, well seasoned timber in this product full strength for 3 minutes or apply 3 flowing coats to the surface with a brush or coarse low pressure spray.

LOS ANGELES CHEMICAL COMPANY

LOT NO.



NET WEIGHT
50 POUNDS

COPRO 53

CAUTION
HARMFUL IF SWALLOWED
KEEP OUT OF REACH OF CHILDREN

ACCEPTED
OCT 1 1968
UNDER THE PROVISIONS OF THE
FUNGICIDE ACT OF 1948 AND
THE PESTICIDE ACT OF 1948
ED UNDER FEDERAL REGULATION
TO THE FIP COMMISSION

Harmful if swallowed. Do not swallow or inhale.

KEEP CONTAINER CLOSED

DESTROY CONTAINER WHEN EMPTY

COMPOSITION

ACTIVE INGREDIENT:	% BY WT.
copper oxychloride sulfate*	94.6%
INERT INGREDIENTS:	5.4%

* Equivalent to 53% copper as metallic by wt. this product.

DIRECTIONS FOR USE

To mix this product, fill spray tank 3/4 full, begin agitation, and add required amounts of spray adjuvants. As tank continues to fill add required amount of this product. To prevent curdling of spray mix due to hard water, a water softener should be added during mixing operations. Constant agitation during the spraying operation is necessary to maintain adequate suspension. Do not apply to or allow this product to drift onto crop or non cropland not specifically being treated. Do not apply to any crop not stated on the label. This product is compatible with most other insecticides and/or fungicides, except lime sulfur and dithiocarbamates. For specific crop uses consult rear panel.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of LOS ANGELES CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

MANUFACTURED BY

LOS ANGELES CHEMICAL COMPANY

4545 ARDINE STREET SOUTH GATE, CALIFORNIA

CB 7/68