

PRATT FERTILIZERS
SYSTEMIC FUNGICIDE

Provides Curative and Protective Action for
TURF, FLOWER, VEGETABLE, ORNAMENTAL

ACTIVE INGREDIENT: Benomyl, Methyl 1-(butyl-1H-imidazol-2-yl)-3-methyl-2-pyrrolidone-1-carbamate, 50% INERT INGREDIENTS: 50%

US Pat. Nos. 3,541,213 and 3,631,177

EPA Reg. No. 904-224

100% No. 1100-11-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT: In case of contact, flush skin and eyes with plenty of water. Get medical attention.

NET WEIGHT: 1 L. AV
2 OZ. AV
4 OZ. AV
8 OZ. AV

PACKAGED FOR
PRATT-GABRIEL DIVISION
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331/An Alco Standard Company

100 100

100 100

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS-CAUTION: May irritate eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. ENVIRONMENTAL HAZARD-This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur or when weather conditions favor drift from areas treated. CHEMICAL HAZARD-Keep away from fire or spark. Never allow Benomyl 50w to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of Benomyl 50w as a fungicide.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1-14-54

...store this product in its original container, tightly closed, and in a cool, dry place away from children and out of the reach of animals. Do not use material that is wet, frozen, or has not been contaminated with water. Do not use material that is wet, frozen, or has not been contaminated with water. Do not use material that is wet, frozen, or has not been contaminated with water.

Bermy 50W is recommended for the control of certain diseases of flowers, ornamentals and turf. It is highly effective for the control of many fungi diseases, providing both curative and preventive action and protective action. Bermy 50W is a wettable powder to be mixed with water for application as a spray. Add the proper quantity of fungicide to half-filled sprayer tank and add the proper quantity of water. Agitate mixture while spraying to assure uniformity.

FLOWERS, ORNAMENTS (Field and Greenhouse): Powdery Mildew, Gray Mold (Botrytis). Use one tablespoonful per 2 gallons water. Begin application when disease first appears and repeat at 7-14 day intervals as long as needed.

TURF: Dollar spot (Blizzaria Homeocarpa), Large Brown Spot (Rhizoctonia Scrieri), Fusarium Patch (Fusarium Niveum). Apply at the rate of 2 ounces (12 tablespoonful) per 1000 sq. ft. in 500 gal. water to obtain thorough coverage, usually 2.5 gallons per 1000 sq. ft. Apply at earliest appearance of disease and continue at 7-14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5-7 days. Do not graze or feed clippings from treated areas to livestock.

ROSES (Field and Greenhouse): Powdery Mildew, Black Spot. Use one tablespoonful per 2 gallons water. Begin application when disease first appears and repeat at 7-14 intervals throughout the growing season, shorten intervals during humid, rainy weather.

CUCUMBERS, MELONS, SUMMER and WINTER SQUASH: Gummy Stem Blight, Powdery Mildew, Anthracnose. Apply 4.8 ounces per acre (2 level tablespoonful in 5 gallons water applied to 1000 sq. ft.). Begin applications when plants begin to run or when disease first appears and repeat at 7-14 day intervals as needed. Under conditions of severe disease pressure, use the higher rate and apply at 7 day intervals or as needed.

BEANS (SNAP): White Rot (Sclerotinia), Gray Mold (Botrytis). Apply 1.2 pounds per acre (2 1/2 level tablespoonful in 5 gallons water applied to 1000 sq. ft.) at 25% to 50% bloom, repeat at peak bloom. Do not apply within 14 days of harvest. Do not graze or feed treated bean vines or hay to livestock.

BULBS (Easter Lily, Tulip, Gladiolus, Dahlias, Iris): Fusarium and Penicillium Rots. Use 1 1/2 pounds per 100 gallons water (2 level tablespoonful per gallon). Soak cleaned bulbs for 15-30 minutes in warm dip (80° to 85°F) preferably within 48 hours after digging. Dry bulbs after treatment. If bulbs are for forcing, treat after bulbs have been heat-cured.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

BEST AVAILABLE



904-224
 300 101663
 17 1

H. Pratt
 116 Clinton Place
 Hackensack, NJ 07601

Attention: H. B. Pratt

Gentlemen:

Subject: Pratt Benzyl 507 Systemic Fungicide
 EPA Registration No. 904-224
 Your letter dated July 11, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. Add "And Domestic Animals" after "Hazards to Humans."
2. In the Environmental Hazard Statement change "Keep...water" to "Do not apply directly to water and wetlands."

A stamped copy is enclosed for your records.

Sincerely yours,

Henry M. Jacoby
 Product Manager (21)
 Fungicide-Herbicide Branch
 Registration Division (TS-767)

Enclosure

RD:Jacobys:DCR-10794:WANG-0265C:bjc:Raven:479-2013:7/29/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							