



TURF FUNGICIDE

50% Wettable Powder

CONTAINS: DYRENE®

An Organic Fungicide For: LAWNS & TURF

- CONTROLS: • Helminthosporium Disease (Melting-out, Leaf Spot)
- Copper Spot • Sclerotinia Dollor Spot
- Snow Mold (Typhula Sp.) • RUST • BROWN PATCH

ACTIVE INGREDIENT:	BY WEIGHT
2,4-dichloro-6-(o-chloroanilino)-s-triazine.....	50.0%
INERT INGREDIENTS:.....	50.0%
TOTAL.....	<u>100.0%</u>

DYRENE® - Registered Trademark of Mobay Chemical Corporation

EPA Reg. No. 904-229

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Call a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF INHALED: Remove victim to fresh air.
- IF ON SKIN: Wash immediately with soap and warm water. If irritation occurs, get medical attention.
- IF IN EYES: Immediately flush eyes with plenty of water. Call a physician.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.


Pratt-Gabriel Division
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331

ACCEPTED

JAN 6 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 904-229

NET WEIGHT: ONE POUND
5 POUNDS



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: DANGER--Causes irreversible eye damage. Harmful if swallowed. Avoid contact with skin. Do not get in eyes or on clothing. Wear safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Protective clothing should be worn during periods of exposure, such as during application or when contacting treated foliage. ANILAZINE may cause a temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of after effects or permanent injury.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

PHYSICAL OR CHEMICAL HAZARDS

Protect from rain and moisture. Keep away from space heaters, and walls that sweat. Do not store next to herbicides. Store in a cool dry place.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not graze treated areas. Do not feed clippings to livestock. Do not allow children or pets to play on lawn for 24 hours after treatment. Do not use on seed crops intended to be used for feed or forage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container.

FOR DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS

TURF FUNGICIDE is a wettable powder that mixes with water to make a turfgrass spray to prevent: Brown Patch; Copper Spot; Dollar Spot; Leaf Spot; Melting-out; Rust; and Snow Mold (*Typhula* sp.)

DILUTION RATE: 1/4 - 1/2 pound in 5 gallons of water per 1,000 sq. ft. (50' x 20').

For a Preventive Program: Begin applications before diseases appear; and repeat at 10 to 14 day intervals throughout the spring and summer months - except during extended periods of cool, moist conditions. During these weather conditions, repeated applications at 7-day intervals are advised. Use the lower rate in the preventive program. If disease pressure increases, increase the rate of application of this product to the higher rate of 1/2 pound per 1,000 sq. ft. as noted above.

For Control of Snow Mold: Apply spray at lower specific rate after the first killing frost in the Fall. Make a second application when the turf is free of snow in mid-Winter. A third application should be made after the final snow has melted in the early Spring.

Schedule applications in accordance with local conditions. Consult your County Agricultural Extension Service for specific timing of applications.

LIMITED WARRANTY: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.