

DIRECTIONS FOR USE ON TURF

PESTS OF LAWN AND OTHER ORNAMENTAL AND RECREATIONAL TURFGRASS AREAS Use Dursban 250 EC to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and sod webworms, lawn moths, etc. Mix Dursban 250 EC with water according to directions given in the following table and apply as a coarse low pressure spray using suitable application equipment. For best results, the lawn should be treated at the time of treatment and care should be taken to obtain complete and uniform coverage of pest infested area. Spray when pests first appear, repeat when needed. Keep children and pets off treated areas until spray has dried. Do not graze treated areas.

DOSAGE AND DILUTION RATES FOR USE ON TURF

Areas to be treated	Amount of Dursban 250 EC	Dilute with water to give
100 sq. ft.	2.5 fl. oz.	15 gallons
200 sq. ft.	1.4 fl. oz.	30 gallons
500 sq. ft.	1.4 fl. oz.	150 gallons
1,000 sq. ft.	1.4 fl. oz.	125 gallons
1 acre	1.4 fl. oz.	1,200 gallons

TO CONTROL CERTAIN WHITE GRUBS Larvae of European Chaffer and Japanese beetle. Use twice a year, the amounts of Dursban 250 EC shown in the Turf table, i.e., 1.4 fl. oz. per 1,000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service representative for proper timing to treat in your area.

TO CONTROL TURFGRASS (HYPERODES) WEEVILS Use twice the amount of Dursban 250 EC shown in Turf table, i.e., 1 gallon diluted in 1,000 gallons of water per acre. Spray suspected problem areas in mid April and again in mid May. Consult your local Agricultural Extension Service representative for proper timing to treat in your area.

NOTE: To help prevent infestation of lawns and gardens by chinch bugs, crickets, grasshoppers, earwigs, etc., treat a band of soil 6 to 10 feet wide around and adjacent to the building and the building foundation to a height of 10 to 12 feet. Where pests are active in the soil, use the same dosage per acre as shown in the Turf table. For these pests, the timing shown in the Turf table. Apply as a coarse spray, thoroughly and uniformly, with the band area.

NOTE: Mixtures of Dursban 250 EC and water should be stored in a cool, dry place. Do not use if the mixture has been stored for more than 30 days. Do not use if the mixture has been stored for more than 30 days. Do not use if the mixture has been stored for more than 30 days.

B.G.

PRATT

SINCE 1904
TM

DURSBAN ★ 250 EC

LAWN AND ORNAMENTAL PLANT INSECTICIDE

Controls

- Chinch Bugs
- Certain White Grubs
- Hyperodes Weevils
- Bagworms
- And Other Insects As Listed

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate)
Aromatic petroleum derivative solvent

INERT INGREDIENTS:

BY WEIGHT

24.9%
53.7%
21.4%

TOTAL..... 100.0%

Contains 2 pounds of chlorpyrifos per gallon.

*DURSBAN - - TRADEMARK OF THE DOW CHEMICAL COMPANY
U. S. PATENT NO. 3,244,586

FOR PROFESSIONAL USE ONLY

WARNING:
KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNINGS

B.G. Pratt Division, Gabriel Chemicals Ltd.
204 21st Avenue, Paterson, N.J. 07509

NET CONTENTS: 1 5 30 55 GALLONS

DIRECTIONS

PESTS OF ORNAMENTAL AREAS Use Dursban 250 EC to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and sod webworms, lawn moths, etc. Mix Dursban 250 EC with water according to directions given in the following table and apply as a coarse low pressure spray using suitable application equipment. For best results, the lawn should be treated at the time of treatment and care should be taken to obtain complete and uniform coverage of pest infested area. Spray when pests first appear, repeat when needed. Keep children and pets off treated areas until spray has dried. Do not graze treated areas.

NOTE: Do not use if the mixture has been stored for more than 30 days. Do not use if the mixture has been stored for more than 30 days. Do not use if the mixture has been stored for more than 30 days.

DOSAGE AND DILUTION RATES FOR USE ON TURF

Areas to be treated	Amount of Dursban 250 EC	Dilute with water to give
100 sq. ft.	2.5 fl. oz.	15 gallons
200 sq. ft.	1.4 fl. oz.	30 gallons
500 sq. ft.	1.4 fl. oz.	150 gallons
1,000 sq. ft.	1.4 fl. oz.	125 gallons
1 acre	1.4 fl. oz.	1,200 gallons

1.4 fl. oz. per 1,000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service representative for proper timing to treat in your area.

HUMAN WARNING
SWALLOWED. MAY CAUSE ILLNESS. Wash thoroughly all contaminated eyes; breathing vapors are harmful. Do not use in areas where children and pets are present. Do not use in areas where children and pets are present. Do not use in areas where children and pets are present.

NOTE TO PHYSICIAN: Do not spray in areas where children and pets are present. Do not use in areas where children and pets are present. Do not use in areas where children and pets are present.

FLAMMABILITY: DO NOT USE IN AREAS WHERE FLAMMABLE LIQUIDS OR GASES ARE PRESENT. DO NOT USE IN AREAS WHERE FLAMMABLE LIQUIDS OR GASES ARE PRESENT. DO NOT USE IN AREAS WHERE FLAMMABLE LIQUIDS OR GASES ARE PRESENT.

ENVIRONMENTAL: Keep out of lakes, streams, rivers, etc. Do not apply when weather is windy. Do not apply when weather is windy. Do not apply when weather is windy. Do not apply when weather is windy. Do not apply when weather is windy.

DO NOT REUSE EMPTY CONTAINERS: Do not reuse empty containers for any other purpose. Do not reuse empty containers for any other purpose. Do not reuse empty containers for any other purpose.

NOTICE BUYER AS TO MATERIAL NOT HEREWITH

B.G.

PRATT

SINCE 1904

TM

DURSBAN ★ 250 EC

LAWN AND ORNAMENTAL PLANT INSECTICIDE

Controls

- Chinch Bugs
- Certain White Grubs
- Hyperodes Weevils
- Bagworms
- And Other Insects As Listed

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate]
Aromatic petroleum derivative solvent

INERT INGREDIENTS:

BY WEIGHT

24.9%

53.7%

21.4%

TOTAL..... 100.0%

Contains 2 pounds of chlorpyrifos per gallon.

*DURSBAN - TRADEMARK OF THE DOW CHEMICAL COMPANY
U.S. PATENT NO. 3,244,586

FOR PROFESSIONAL USE ONLY

WARNING:

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNINGS

B.G. Pratt Division, Gabriel Chemicals Ltd.
204 21st Avenue, Paterson, N.J. 07509

NET CONTENTS: GALLONS

1 5 30 55

DIRECTIONS FOR USE ON ORNAMENTAL PLANTS

PESTS OF ORNAMENTALS: Use Dursban 250 EC to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, exposed thrips, and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests.

Dilute Dursban 250 EC with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals if needed.

NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

DOSAGE AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS

Pests Controlled	Amount of Dursban 250 EC in Water to Make		
	3 Gallons	15 Gallons	100 Gallons
Bagworms Grasshoppers Mites Spittlebugs Whiteflies	1 fl. oz.	1.6 fl. oz.	1 pint*
Cutworms Mealybugs Leafhoppers Thrips (exposed)	1 fl. oz.	3.2 fl. oz.	1 quart**

*Equivalent to 4 ounces of Chlorpyrifos per 100 gallons

**Equivalent to 8 ounces of Chlorpyrifos per 100 gallons

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN. MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH THE SKIN. MAY BE INJURIOUS TO EYES AND SKIN. COMBUSTIBLE. Do not take internally. Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of water and get prompt medical attention. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs and domestic water supplies. Handle concentrate in a ventilated area. Keep children and pets off treated areas until spray has dried. Do not spray vegetable gardens or other plants used for food purposes.

Do not spray vegetable gardens or other plants used for food purposes.
NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treating symptomatically. Atropine is an antidote.

FLAMMABILITY WARNING

DO NOT USE, POUR, FILL OR STORE NEAR HEAT OR OPEN FLAMES. KEEP CONTAINER CLOSED.

ENVIRONMENTAL CAUTION: This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on plants. For more information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Promptly crush empty containers and bury them and wastes in locations away from domestic water supplies. Do not reuse empty drums. Return to a drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTICE: BUYER ASSUMES ALL RISKS OF USE, STORAGE, OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS GIVEN HEREWITH.

EPA Reg.
No. 904-234

EPA Est.
904-NJ-1