

B. G. PRATT DIVISION

GABRIEL CHEMICALS, LTD. • 204 Twenty-First Avenue • Paterson, New Jersey 07624
(201) 684-4400

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not breathe spray mist or take internally. Avoid contact with skin. Wear gloves and long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after handling. Do not contaminate food or feed. Avoid storage near food or feed products.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. For protection of bees, avoid use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers to locate hives at a safe distance until one week after application. Further protective information may be obtained from your Cooperative Agricultural Extension Service. Do not allow drifting to adjoining food or forage crops.

Do not reuse empty container. Wrap in newspaper and place in trash can for safe disposal or bury in a safe place away from water supplies.


Do not reuse empty drum. Return to a drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

PROTECT FROM TEMPERATURES BELOW 32° F. Keep cover on opened container to prevent drying.

DIRECTIONS: L 2 Sevin is an extremely fine dispersion of Sevin in water. SHAKE WELL BEFORE USING. Pour recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.

LAWNS: For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Sod Webworms (Lawn Moths) and Millipedes, apply 8 fl. oz. in 15-20 gallons of water evenly over each 500 sq. ft. of established lawn area. For best results, mow the lawn and apply immediately after rain or watering. Do not water for at least 2 days after application. Repeat in 2-3 weeks if necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: For use on Roses, Zinnias, Gladioli, Carnations, Lilacs, Arborvitae, Junipers, Hydrangeas, Azaleas, Elms, Chrysanthemums, Maples, Oaks, Dog



PRATT

SINCE 1904
T.M.

L 2 SEVIN® PLANT INSECTICIDE

- FOR
- LAWNS
 - ORNAMENTAL PLANTS
 - FLOWERS
 - SHRUBS
 - SHADE TREES

KILLS
• CHINCH BUGS
• JAPANESE BEETLES
• MOSQUITOES
AND CERTAIN
OTHER INSECTS
as listed

ACTIVE INGREDIENTS	BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)	21.5%
INERT INGREDIENTS	78.5%
TOTAL	100.0%

Registered Trademark of Union Carbide Corporation U.S. Patent No. 2,903,478

Contains 2 lbs. of Sevin per gallon.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

NET CONTENTS:
GALLONS

1 5 30 55

EPA
Est. 904-NJ-1
EPA
Reg. No. 904-241
PROD NO

wood, Birch and Pines (DO NOT USE ON BOSTON IVY OR VIRGINIA CREEPER). Use 4 teaspoonful per gallon of water or 1 pint in 25 gallons (1:200 dilution) to control: Apple Aphid, Bagworm, Birchleaf Miner, Bristle Beetles, Boxelder Bugs, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, June Beetle, Lace Bug, Leafhoppers, Leafrollers, Mealy bugs, Mimosa Webworms, Oakleaf Miner, Orange Tortrix, Periodical Cicada, Plant Bugs, Puss Caterpillars, Psyllids, Rose Aphids, Rose Slug, Scale Insects, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Willowleaf Beetles. Apply when insects first appear. Repeat weekly or as necessary throughout the season.

ADULT MOSQUITO CONTROL: For Home Grounds, Parks and Recreation Areas, use 1:200 dilution in hydraulic low pressure sprayers (4 teaspoonful per 1 gallon or 1 pint to 25 gallons of water). Apply in early morning or evening when air is calm and adults are active. Spray to thoroughly cover low shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate (DO NOT APPLY TO WATER). Repeat at 7-10 day intervals or as necessary (DO NOT SPRAY BOSTON IVY OR VIRGINIA CREEPER). May kill shrimp and crabs. Do not apply where these are important resources.

VEGETABLE GARDEN SPRAY: Mix 1 fl. oz. in 3 gallons of water to 650 sq. ft. (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7-10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with the directions:

BEANS (Green, Lima, and Snap), Japanese and Mexican Bean Beetle, Tarnished Plant Bug, TOMATOES, EGG-PLANTS and PEPPERS, Leafhopper, Lace Bug, Tomato Fruitworm, PUMPKINS, SQUASH, CUCUMBERS, MELONS, Flea Beetle, Melonworms, Cucumber Beetle, Squash Bug, SWEET CORN, Corn Earworm, Japanese Beetle and Leafhoppers. Mix 1 fl. oz. in 3 gallons of water to cover 650 sq. ft. (2 qts. in sufficient water for 1 acre). Application of this product to the tassel region of corn during the pollen shed period will severely reduce bee population. For European Corn Borer, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Do not apply within 7 days of harvest if used for feed or forage. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS GIVEN HEREWITH.

B.G. Pratt Division
Gabriel Chemicals Ltd.
204 21st Avenue
Paterson, N.J. 07509

SPRAYS FUNGICIDES HERBICIDES INSECTICIDES

SINCE 1904