

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION--Harmful if inhaled or swallowed. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. Do not take internally. Wear long-sleeves. Change clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Keep children and pets off treated areas until spray has dried. ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of waste. BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

PROTECT FROM TEMPERATURES BELOW 32° F. CLOSE CONTAINER IMMEDIATELY AFTER USING TO PREVENT EVAPORATION OR DRYING. SHAKE WELL BEFORE USING.

READ ENTIRE LABEL BEFORE USING--OBSERVE ALL CAUTIONS--USE ONLY AS DIRECTED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (8 OZ. - 1 GAL.): Do not contaminate water, feed or food by storage or disposal. Store in the closed, original container in a locked area away from children and domestic animals. For disposal, securely wrap closed container in several layers of newspaper and place in trash collection.

STORAGE AND DISPOSAL (5 GAL. - 55 GAL.): Do not contaminate water, food or feed by storage or disposal. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or reuse and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic containers may be incinerated, or if allowed by state and local authorities, burned. If burned, stay out of smoke.

MIXING DIRECTIONS: L2 Sevin is an extremely fine dispersion of carbaryl in water. Stir thoroughly or shake before measuring. Pour recommended amount of concentrate into nearly full spray tank or premix material with water in another container and add to half full spray tank. Add balance of water with agitation. Agitate or stir well frequently while making application.

SPRAY THOROUGHLY TO COVER ALL PLANT SURFACES.

BEST DOCUMENT AVAILABLE

LAWNS: For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Sod Webworms (Lawn Moths), and Millipedes, apply 8 fl. oz. in 15-20 gallons of water evenly over each 500 sq. ft. of established lawn area. For best results, mow the lawn and apply immediately after rain or watering. Do not water for at least 7 days after application. Repeat in 2-3 weeks if necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: For use on Roses, Zinnias, Gladioli, Carnations, Lilacs, Arborvitae, Junipers, Hydrangeas, Azaleas, Elms, Chrysanthemums, Maples, Oaks, Dogwood, Birch and Pines. (DO NOT USE ON BOSTON IVY OR VIRGINIA CREEPER), use 4 teaspoonsful per gallon of water or 1 pint in 25 gallons (1-200 dilution) to control Apple Aphid, Bagworm, Birchleaf Miner, Blister Beetles, Boxelder Bugs, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, June Beetle, Lace Bug, Leafhoppers, Leafrollers, Mealybugs, Mimosa Webworms, Oakleaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Puss Caterpillars, Psyllids, Rose Slug, Scale Insects, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Willowleaf Beetles. Apply when insects first appear. Repeat weekly or as necessary throughout the season.

ADULT MOSQUITO CONTROL: For Home Grounds, Parks and Recreational Areas, use 1-200 dilution in hydraulic low pressure sprayers (4 teaspoonsful per 1 gallon or 1 pint to 25 gallons of water). Apply in early morning or evening when air is calm and adults are active. Spray to thoroughly cover low shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate (DO NOT APPLY TO WATER). Repeat at 7-10 day intervals or as necessary. (DO NOT SPRAY BOSTON IVY OR VIRGINIA CREEPER). May kill shrimp and crabs. Do not apply where these are important resources.

VEGETABLE GARDEN SPRAY: Mix 1 fl. oz. in 3 gallons of water to 650 sq. ft. (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7-10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with the directions.

BEANS (Green, Lima, and Snap): Japanese and Mexican Bean Beetle, Tarnished Plant Bug;

TOMATOES, EGGPLANTS AND PEPPERS: Leafhopper, Lace Bug, Tomato Fruitworm; **PUMPKINS,**

SQUASH, CUCUMBERS MELONS: Flea Beetle, Melonworms, Cucumber Beetle, Squash Bug;

SWEET CORN: Corn Earworm, Japanese Beetle and Leafhoppers: Mix 1½ fl. oz. in 3 gallons of water to cover 650 sq. ft. (2 qts. in sufficient water for 1 acre).

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. For European Corn Borer, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Do not apply within 7 days of harvest if used for feed or forage. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS HEREWITH.

BEST DOCUMENT AVAILABLE

Mr. H. B. Pratt
Pratt-Gabriel Division
Hillier Chemical & Fertilizer Co.,
116 Clinton Place
Hackensack, NJ 07601

Dear Mr. Pratt:

Subject: General Review of Label
Pratt 42 Sevin
EPA Registration No. 10-221
Application Date October 29, 1963

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Make the labeling changes listed below before you release the product for shipment.

1. Add the statement listed below to your Environmental Hazards paragraph:

This product is toxic to fish.

2. In the statement about leaf injury due to high humidity in your Use Directions note the spelling of "the" and "presence."

3. Revise your Disposal Directions to read as follows:

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. Rinse thoroughly before disposal in trash.

Ray A. Releand
Product Buyer (12)
Insecticide-Rodenticide Branch
Registration Division (13-767)

ENCLOSURE

13-767:Ellenberger:1008-20430: A 05-179:101:150: DIV: 13-767:11/8/67

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

EPA Form 1320-1 (4-81)

OFFICIAL FILE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION--Harmful if inhaled or swallowed. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. Do not take internally. Wear long-sleeves. Change clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Keep children and pets off treated areas until spray has dried. ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of waste. BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

PROTECT FROM TEMPERATURES BELOW 32° F. CLOSE CONTAINER IMMEDIATELY AFTER USING TO PREVENT EVAPORATION OR DRYING. SHAKE WELL BEFORE USING.

READ ENTIRE LABEL BEFORE USING--OBSERVE ALL CAUTIONS--USE ONLY AS DIRECTED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (8 OZ. - 1 GAL.): Do not contaminate water, feed or food by storage or disposal. Store in the closed, original container in a locked area away from children and domestic animals. For disposal, securely wrap closed container in several layers of newspaper and place in trash collection.

STORAGE AND DISPOSAL (5 GAL. - 55 GAL.): Do not contaminate water, food or feed by storage or disposal. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or store and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic containers may be incinerated, or if allowed by state and local authorities, burned. If burned, stay out of smoke.

MIXING DIRECTIONS: L2 Sevin is an extremely fine dispersion of carbaryl in water. Stir thoroughly or shake before measuring. Pour recommended amount of concentrate into nearly full spray tank or premix material with water in another container and add to half full spray tank. Add balance of water with agitation. Agitate or stir well frequently while making application.

SPRAY THOROUGHLY TO COVER ALL PLANT SURFACES.

BEST DOCUMENT AVAILABLE

LAWNS: For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Sod Webworms (Lawn Moths), and Millipedes, apply 8 fl. oz. in 15-20 gallons of water evenly over each 500 sq. ft. of established lawn area. For best results, mow the lawn and apply immediately after rain or watering. Do not water for at least 2 days after application. Repeat in 2-3 weeks if necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: For use on Roses, Zinnias, Gladioli, Carnations, Lilacs, Arborvitae, Junipers, Hydrangeas, Azaleas, Elms, C. santhemums, Maples, Oaks, Dogwood, Birch and Pines. (DO NOT USE ON BOSTON IVY OR VIRGINIA CREEPER), use 4 teaspoonsful per gallon of water or 1 pint in 25 gallons (1-200 dilution) to control Apple Aphid, Bagworm, Birchleaf Miner, Blister Beetles, Boxelder Bugs, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, June Beetle, Lace Bug, Leafhoppers, Leafrollers, Mealybugs, Mimosa Webworms, Oakleaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Puss Caterpillars, Psyllids, Rose Slug, Scale Insects, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Willowleaf Beetles. Apply when insects first appear. Repeat weekly or as necessary throughout the season.

ADULT MOSQUITO CONTROL: For Home Grounds, Parks and Recreational Areas, use 1-200 dilution in hydraulic low pressure sprayers (4 teaspoonsful per 1 gallon or 1 pint to 25 gallons of water). Apply in early morning or evening when air is calm and adults are active. Spray to thoroughly cover low shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate (DO NOT APPLY TO WATER). Repeat at 7-10 day intervals or as necessary. (DO NOT SPRAY BOSTON IVY OR VIRGINIA CREEPER). May kill shrimp and crabs. Do not apply where these are important resources.

VEGETABLE GARDEN SPRAY: Mix 1 fl. oz. in 3 gallons of water to 650 sq. ft. (2 qts. in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7-10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with the directions.

BEANS (Green, Lima, and Snap): Japanese and Mexican Bean Beetle, Tarnished Plant Bug;

TOMATOES, EGGPLANTS AND PEPPERS: Leafhopper, Lace Bug, Tomato Fruitworm; **PUMPKINS,**

SQUASH, CUCUMBERS MELONS: Flea Beetle, Melonworms, Cucumber Beetle, Squash Bug;

SWEET CORN: Corn Earworm, Japanese Beetle and Leafhoppers: Mix 1½ fl. oz. in 3 gallons of water to cover 650 sq. ft. (2 qts. in sufficient water for 1 acre).

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. For European Corn Borer, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Do not apply within 7 days of harvest if used for feed or forage. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS HEREWITH.

NOV 18 1983

Mr. H. B. Pratt
Pratt-Gabriel Division
Hillar Chemical & Fertilizer Corp.
116 Clinton Place
Hackensack, NJ 07601

Dear Mr. Pratt:

Subject: General Revision Of Label
Pratt L2 Sevin
EPA Registration No. 90-1-241
Application Date October 20, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Make the labeling changes listed below before you release the product for shipment.

1. Add the statement listed below to your Environmental Hazards paragraph:

This product is toxic to fish.

2. In the statement about leaf injury due to high humidity in your Use Directions note the spelling of "the" and "presence."
3. Revise your Disposal directions to read as follows:

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:Ellenberger:DCR-28430:WANG-1212C:lsa:Raven:479-2013:11/8/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

